

NOT FOR SALE

CHAPTER P Preparation for Calculus

Section P.1	Graphs and Models.....	2
Section P.2	Linear Models and Rates of Change.....	11
Section P.3	Functions and Their Graphs.....	23
Section P.4	Fitting Models to Data.....	33
Review Exercises	35
Problem Solving	41

INSTRUCTOR USE ONLY

CHAPTER P Preparation for Calculus

Section P.1 Graphs and Models

1. $y = -\frac{3}{2}x + 3$

x -intercept: (2, 0)

y -intercept: (0, 3)

Matches graph (b).

2. $y = \sqrt{9 - x^2}$

x -intercepts: (-3, 0), (3, 0)

y -intercept: (0, 3)

Matches graph (d).

3. $y = 3 - x^2$

x -intercepts: $(\sqrt{3}, 0), (-\sqrt{3}, 0)$

y -intercept: (0, 3)

Matches graph (a).

4. $y = x^3 - x$


x -intercepts: (0, 0), (-1, 0), (1, 0)

y -intercept: (0, 0)

Matches graph (c).


5. $y = \frac{1}{2}x + 2$

x	-4	-2	0	2	4
y	0	1	2	3	4




6. $y = 5 - 2x$

x	-1	0	1	2	$\frac{5}{2}$	3	4
y	7	5	3	1	0	-1	-3




7. $y = 4 - x^2$

x	-3	-2	0	2	3
y	-5	0	4	0	-5




8. $y = (x - 3)^2$

x	0	1	2	3	4	5	6
y	9	4	1	0	1	4	9




9. $y = |x + 2|$

x	-5	-4	-3	-2	-1	0	1
y	3	2	1	0	1	2	3




10. $y = |x| - 1$

x	-3	-2	-1	0	1	2	3
y	2	1	0	-1	0	1	2




11. $y = \sqrt{x} - 6$

x	0	1	4	9	16
y	-6	-5	-4	-3	-2




12. $y = \sqrt{x + 2}$

x	-2	-1	0	2	7	14
y	0	1	$\sqrt{2}$	2	3	4




13. $y = \frac{3}{x}$

x	-3	-2	-1	0	1	2	3
y	-1	$-\frac{3}{2}$	-3	Undef.	3	$\frac{3}{2}$	1



14. $y = \frac{1}{x + 2}$

x	-6	-4	-3	-2	-1	0	2
y	$-\frac{1}{4}$	$-\frac{1}{2}$	-1	Undef.	1	$\frac{1}{2}$	$\frac{1}{4}$




15. $\begin{matrix} \text{Xmin} = -5 \\ \text{Xmax} = 4 \\ \text{Xscl} = 1 \\ \text{Ymin} = -5 \\ \text{Ymax} = 8 \\ \text{Yscl} = 1 \end{matrix}$

Note that $y = -3$ when $x = 0$ and $y = 0$ when $x = -1$.

16. $\begin{matrix} \text{Xmin} = -20 \\ \text{Xmax} = 30 \\ \text{Xscl} = 5 \\ \text{Ymin} = -10 \\ \text{Ymax} = 40 \\ \text{Yscl} = 5 \end{matrix}$

Note that $y = 16$ when $x = 0$ or 16 .


17. $y = \sqrt{5 - x}$



(a) $(2, y) = (2, 1.73)$ ($y = \sqrt{5 - 2} = \sqrt{3} \approx 1.73$)

(b) $(x, 3) = (-4, 3)$ ($3 = \sqrt{5 - (-4)}$)

18. $y = x^5 - 5x$



(a) $(-0.5, y) = (-0.5, 2.47)$

(b) $(x, -4) = (-1.65, -4)$ and $(x, -4) = (1, -4)$

19. $y = 2x - 5$

y-intercept: $y = 2(0) - 5 = -5$; $(0, -5)$

x-intercept: $0 = 2x - 5$

$5 = 2x$

$x = \frac{5}{2}$; $(\frac{5}{2}, 0)$

20. $y = 4x^2 + 3$

y-intercept: $y = 4(0)^2 + 3 = 3$; $(0, 3)$

x-intercept: $0 = 4x^2 + 3$

$-3 = 4x^2$

None. y cannot equal 0.

21. $y = x^2 + x - 2$

y-intercept: $y = 0^2 + 0 - 2$

$y = -2$; $(0, -2)$

x-intercepts: $0 = x^2 + x - 2$

$0 = (x + 2)(x - 1)$

$x = -2, 1$; $(-2, 0), (1, 0)$

22. $y^2 = x^3 - 4x$

y-intercept: $y^2 = 0^3 - 4(0)$

$y = 0$; $(0, 0)$

x-intercepts: $0 = x^3 - 4x$

$0 = x(x - 2)(x + 2)$

$x = 0, \pm 2$; $(0, 0), (\pm 2, 0)$

23. $y = x\sqrt{16 - x^2}$

y-intercept: $y = 0\sqrt{16 - 0^2} = 0$; $(0, 0)$

x-intercepts: $0 = x\sqrt{16 - x^2}$

$0 = x\sqrt{(4 - x)(4 + x)}$

$x = 0, 4, -4$; $(0, 0), (4, 0), (-4, 0)$

24. $y = (x - 1)\sqrt{x^2 + 1}$

y-intercept: $y = (0 - 1)\sqrt{0^2 + 1}$

$y = -1$; $(0, -1)$

x-intercept: $0 = (x - 1)\sqrt{x^2 + 1}$

$x = 1$; $(1, 0)$

25. $y = \frac{2 - \sqrt{x}}{5x}$

y-intercept: None. x cannot equal 0.

x-intercept: $0 = \frac{2 - \sqrt{x}}{5x}$

$0 = 2 - \sqrt{x}$

$x = 4$; $(4, 0)$

26. $y = \frac{x^2 + 3x}{(3x + 1)^2}$

y-intercept: $y = \frac{0^2 + 3(0)}{[3(0) + 1]^2}$

$y = 0$; $(0, 0)$

x-intercepts: $0 = \frac{x^2 + 3x}{(3x + 1)^2}$

$0 = \frac{x(x + 3)}{(3x + 1)^2}$

$x = 0, -3$; $(0, 0), (-3, 0)$

27. $x^2y - x^2 + 4y = 0$

y-intercept: $0^2(y) - 0^2 + 4y = 0$

$y = 0$; $(0, 0)$

x-intercept: $x^2(0) - x^2 + 4(0) = 0$

$x = 0$; $(0, 0)$

28. $y = 2x - \sqrt{x^2 + 1}$

y-intercept: $y = 2(0) - \sqrt{0^2 + 1}$

$y = -1$; $(0, -1)$

x-intercept: $0 = 2x - \sqrt{x^2 + 1}$

$2x = \sqrt{x^2 + 1}$

$4x^2 = x^2 + 1$

$3x^2 = 1$

$x^2 = \frac{1}{3}$

$x = \pm \frac{\sqrt{3}}{3}$

$x = \frac{\sqrt{3}}{3}$; $(\frac{\sqrt{3}}{3}, 0)$

Note: $x = -\sqrt{3}/3$ is an extraneous solution.29. Symmetric with respect to the y -axis because

$y = (-x)^2 - 6 = x^2 - 6$.

30. $y = x^2 - x$

No symmetry with respect to either axis or the origin.

31. Symmetric with respect to the x -axis because

$$(-y)^2 = y^2 = x^3 - 8x.$$

32. Symmetric with respect to the origin because

$$\begin{aligned} (-y) &= (-x)^3 + (-x) \\ -y &= -x^3 - x \\ y &= x^3 + x. \end{aligned}$$

33. Symmetric with respect to the origin because

$$(-x)(-y) = xy = 4.$$

34. Symmetric with respect to the x -axis because

$$x(-y)^2 = xy^2 = -10.$$

35. $y = 4 - \sqrt{x + 3}$

No symmetry with respect to either axis or the origin.

36. Symmetric with respect to the origin because

$$\begin{aligned} (-x)(-y) - \sqrt{4 - (-x)^2} &= 0 \\ xy - \sqrt{4 - x^2} &= 0. \end{aligned}$$

37. Symmetric with respect to the origin because

$$\begin{aligned} -y &= \frac{-x}{(-x)^2 + 1} \\ y &= \frac{x}{x^2 + 1}. \end{aligned}$$

38. $y = \frac{x^2}{x^2 + 1}$ is symmetric with respect to the y -axis

$$\text{because } y = \frac{(-x)^2}{(-x)^2 + 1} = \frac{x^2}{x^2 + 1}.$$

39. $y = |x^3 + x|$ is symmetric with respect to the y -axis

$$\text{because } y = |(-x)^3 + (-x)| = |-(x^3 + x)| = |x^3 + x|.$$


40. $|y| - x = 3$ is symmetric with respect to the x -axis

$$\begin{aligned} \text{because} \\ |-y| - x &= 3 \\ |y| - x &= 3. \end{aligned}$$

41. $y = 2 - 3x$

Intercepts: $(0, 2), (\frac{2}{3}, 0)$


Symmetry: None



42. $y = -\frac{3}{2}x + 6$

Intercepts: $(0, 6), (4, 0)$


Symmetry: None



43. $y = \frac{1}{2}x - 4$

Intercepts: $(8, 0), (0, -4)$


Symmetry: none




44. $y = \frac{2}{3}x + 1$

Intercepts: $(0, 1), (-\frac{3}{2}, 0)$

Symmetry: none




45. $y = 9 - x^2$

Intercepts: $(0, 9)$, $(3, 0)$, $(-3, 0)$ Symmetry: y -axis


49. $y = x^3 + 2$

Intercepts: $(-\sqrt[3]{2}, 0)$, $(0, 2)$

Symmetry: none




46. $y = x^2 + 3$

Intercept: $(0, 3)$ Symmetry: y -axis

50. $y = x^3 - 4x$

Intercepts: $(0, 0)$, $(2, 0)$, $(-2, 0)$


Symmetry: origin



47. $y = (x + 3)^2$

Intercepts: $(-3, 0)$, $(0, 9)$


Symmetry: none



51. $y = x\sqrt{x + 5}$

Intercepts: $(0, 0)$, $(-5, 0)$


Symmetry: none




48. $y = 2x^2 + x = x(2x + 1)$

Intercepts: $(0, 0)$, $(-\frac{1}{2}, 0)$

Symmetry: none




52. $y = \sqrt{25 - x^2}$

Intercepts: $(0, 5)$, $(5, 0)$, $(-5, 0)$ Symmetry: y -axis

53. $x = y^3$

Intercept: (0, 0)


Symmetry: origin



54. $x = y^2 - 4$

Intercepts: (0, 2), (0, -2), (-4, 0)


Symmetry: x-axis



55. $y = \frac{8}{x}$

Intercepts: none


Symmetry: origin



56. $y = \frac{10}{x^2 + 1}$

Intercept: (0, 10)


Symmetry: y-axis



57. $y = 6 - |x|$

Intercepts: (0, 6), (-6, 0), (6, 0)


Symmetry: y-axis



58. $y = |6 - x|$

Intercepts: (0, 6), (6, 0)

Symmetry: none




59. $y^2 - x = 9$

$$y^2 = x + 9$$

$$y = \pm\sqrt{x + 9}$$

Intercepts: (0, 3), (0, -3), (-9, 0)

Symmetry: x-axis




60. $x^2 + 4y^2 = 4 \Rightarrow y = \pm\frac{\sqrt{4 - x^2}}{2}$

Intercepts: (-2, 0), (2, 0), (0, -1), (0, 1)

Symmetry: origin and both axes


Domain: [-2, 2]



61. $x + 3y^2 = 6$

$$3y^2 = 6 - x$$


$$y = \pm \sqrt{\frac{6-x}{3}}$$

Intercepts: $(6, 0)$, $(0, \sqrt{2})$, $(0, -\sqrt{2})$ Symmetry: x -axis

62. $3x - 4y^2 = 8$

$$4y^2 = 3x - 8$$

$$y = \pm \sqrt{\frac{3}{4}x - 2}$$

Intercept: $(\frac{8}{3}, 0)$ Symmetry: x -axis

63. $x + y = 8 \Rightarrow y = 8 - x$

$$4x - y = 7 \Rightarrow y = 4x - 7$$

$$8 - x = 4x - 7$$

$$15 = 5x$$

$$3 = x$$

The corresponding y -value is $y = 5$.Point of intersection: $(3, 5)$

64. $3x - 2y = -4 \Rightarrow y = \frac{3x + 4}{2}$

$$4x + 2y = -10 \Rightarrow y = \frac{-4x - 10}{2}$$

$$\frac{3x + 4}{2} = \frac{-4x - 10}{2}$$

$$3x + 4 = -4x - 10$$

$$7x = -14$$

$$x = -2$$

The corresponding y -value is $y = -1$.Point of intersection: $(-2, -1)$

65. $x^2 + y = 6 \Rightarrow y = 6 - x^2$

$$x + y = 4 \Rightarrow y = 4 - x$$

$$6 - x^2 = 4 - x$$

$$0 = x^2 - x - 2$$

$$0 = (x - 2)(x + 1)$$

$$x = 2, -1$$

The corresponding y -values are $y = 2$ (for $x = 2$) and $y = 5$ (for $x = -1$).Points of intersection: $(2, 2)$, $(-1, 5)$

66. $x = 3 - y^2 \Rightarrow y^2 = 3 - x$

$$y = x - 1$$

$$3 - x = (x - 1)^2$$

$$3 - x = x^2 - 2x + 1$$

$$0 = x^2 - x - 2 = (x + 1)(x - 2)$$

$$x = -1 \text{ or } x = 2$$

The corresponding y -values are $y = -2$ (for $x = -1$) and $y = 1$ (for $x = 2$).Points of intersection: $(-1, -2)$, $(2, 1)$

67. $x^2 + y^2 = 5 \Rightarrow y^2 = 5 - x^2$

$$x - y = 1 \Rightarrow y = x - 1$$

$$5 - x^2 = (x - 1)^2$$

$$5 - x^2 = x^2 - 2x + 1$$

$$0 = 2x^2 - 2x - 4 = 2(x + 1)(x - 2)$$

$$x = -1 \text{ or } x = 2$$

The corresponding y -values are $y = -2$ (for $x = -1$) and $y = 1$ (for $x = 2$).Points of intersection: $(-1, -2)$, $(2, 1)$

68. $x^2 + y^2 = 25 \Rightarrow y^2 = 25 - x^2$

$$-3x + y = 15 \Rightarrow y = 3x + 15$$

$$25 - x^2 = (3x + 15)^2$$

$$25 - x^2 = 9x^2 + 90x + 225$$

$$0 = 10x^2 + 90x + 200$$

$$0 = x^2 + 9x + 20$$

$$0 = (x + 5)(x + 4)$$

$$x = -4 \text{ or } x = -5$$

The corresponding y -values are $y = 3$ (for $x = -4$) and $y = 0$ (for $x = -5$).Points of intersection: $(-4, 3)$, $(-5, 0)$

69. $y = x^3$

$y = x$

$x^3 = x$

$x^3 - x = 0$

$x(x + 1)(x - 1) = 0$

$x = 0, x = -1, \text{ or } x = 1$

The corresponding y -values are

$y = 0$ (for $x = 0$), $y = -1$ (for $x = -1$), and

$y = 1$ (for $x = 1$).

Points of intersection: $(0, 0), (-1, -1), (1, 1)$

70. $y = x^3 - 4x$

$y = -(x + 2)$

$x^3 - 4x = -(x + 2)$

$x^3 - 3x + 2 = 0$

$(x - 1)^2(x + 2) = 0$

$x = 1 \text{ or } x = -2$

The corresponding y -values are

$y = -3$ (for $x = 1$) and $y = 0$ (for $x = -2$).

Points of intersection: $(1, -3), (-2, 0)$

71. Analytically,

$y = x^3 - 2x^2 + x - 1$

$y = -x^2 + 3x - 1$


$x^3 - 2x^2 + x - 1 = -x^2 + 3x - 1$

$x^3 - x^2 - 2x = 0$

$x(x - 2)(x + 1) = 0$

$x = -1, 0, 2.$

Points of intersection: $(-1, -5), (0, -1), (2, 1)$



72. Analytically,

$y = x^4 - 2x^2 + 1$

$y = 1 - x^2$


$1 - x^2 = x^4 - 2x^2 + 1$

$0 = x^4 - x^2$

$0 = x^2(x + 1)(x - 1)$


$x = -1, 0, 1.$

Points of intersection: $(-1, 0), (0, 1), (1, 0)$



73. $y = \sqrt{x + 6}$

$y = \sqrt{-x^2 - 4x}$



Points of intersection: $(-2, 2), (-3, \sqrt{3}) \approx (-3, 1.732)$

Analytically, $\sqrt{x + 6} = \sqrt{-x^2 - 4x}$

$x + 6 = -x^2 - 4x$


$x^2 + 5x + 6 = 0$

$(x + 3)(x + 2) = 0$

$x = -3, -2.$

74. $y = -|2x - 3| + 6$

$y = 6 - x$



Points of intersection: $(3, 3), (1, 5)$

Analytically, $-|2x - 3| + 6 = 6 - x$


$|2x - 3| = x$

$2x - 3 = x \text{ or } 2x - 3 = -x$

$x = 3 \text{ or } x = 1.$

75. (a) Using a graphing utility, you obtain

$$y = -0.027t^2 + 5.73t + 26.9.$$




The model is a good fit for the data.

(c) For 2010, $t = 40$ and $y = 212.9$.

76. (a) Using a graphing utility, you obtain

$$y = 0.77t^2 + 2.1t + 4$$



The model is a good fit for the data.

(c) For 2015, $t = 25$ and $y \approx 538$ million subscribers.

77. $C = R$

$$5.5\sqrt{x} + 10,000 = 3.29x$$

$$(5.5\sqrt{x})^2 = (3.29x - 10,000)^2$$

$$30.25x = 10.8241x^2 - 65,800x + 100,000,000$$


$$0 = 10.8241x^2 - 65,830.25x + 100,000,000 \quad \text{Use the Quadratic Formula.}$$

$$x \approx 3133 \text{ units}$$

The other root, $x \approx 2949$, does not satisfy the equation $R = C$.

This problem can also be solved by using a graphing utility and finding the intersection of the graphs of C and R .

78. $y = \frac{10,770}{x^2} - 0.37$



If the diameter is doubled, the resistance is changed by approximately a factor of $\frac{1}{4}$. For instance,

$$y(20) \approx 26.555 \text{ and } y(40) \approx 6.36125.$$

79. Answers may vary. *Sample answer:*

$$y = (x + 4)(x - 3)(x - 8) \text{ has intercepts at } x = -4, x = 3, \text{ and } x = 8.$$

80. Answers may vary. *Sample answer:*

$$y = \left(x + \frac{3}{2}\right)(x - 4)\left(x - \frac{5}{2}\right) \text{ has intercepts at } x = -\frac{3}{2}, x = 4, \text{ and } x = \frac{5}{2}.$$

82. (a) v [Because $y = 3(-x)^2 + 3 = 3x^2 + 3$]

(b) i [Because $y = 3x^3 - 3x = 3x(x - 1)(x + 1)$ has x -intercepts $(0, 0)$, $(1, 0)$, $(-1, 0)$]

(c) None of the equations are symmetric with respect to the x -axis

(d) ii [Because $(-2 + 3)^2 = 1$] and vi [Because $\sqrt{-2 + 3} = 1$]

(e) i [Because $3(-x)^3 - 3(-x) = -3x^3 + 3x = -y$] and iv [Because $\sqrt[3]{-x} = -\sqrt[3]{x} = -y$]

(f) i [Because $3(0)^3 - 3(0) = 0$] and iv [Because $\sqrt[3]{0} = 0$]

81. (a) If (x, y) is on the graph, then so is $(-x, y)$ by y -axis symmetry. Because $(-x, y)$ is on the graph, then so is $(-x, -y)$ by x -axis symmetry. So, the graph is symmetric with respect to the origin. The converse is not true. For example, $y = x^3$ has origin symmetry but is not symmetric with respect to either the x -axis or the y -axis.

(b) Assume that the graph has x -axis and origin symmetry. If (x, y) is on the graph, so is $(x, -y)$ by x -axis symmetry. Because $(x, -y)$ is on the graph, then so is $(-x, -(-y)) = (-x, y)$ by origin symmetry. Therefore, the graph is symmetric with respect to the y -axis. The argument is similar for y -axis and origin symmetry.


83. False. x -axis symmetry means that if $(-4, -5)$ is on the graph, then $(-4, 5)$ is also on the graph. For example, $(4, -5)$ is not on the graph of $x = y^2 - 29$, whereas $(-4, -5)$ is on the graph.

84. True. $f(4) = f(-4)$.

85. True. The x -intercepts are $\left(\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}, 0\right)$.

86. True. The x -intercept is $\left(-\frac{b}{2a}, 0\right)$.

87.



$$2\sqrt{(x-0)^2 + (y-3)^2} = \sqrt{(x-0)^2 + (y-0)^2}$$

$$4[x^2 + (y-3)^2] = x^2 + y^2$$

$$4x^2 + 4y^2 - 24y + 36 = x^2 + y^2$$

$$3x^2 + 3y^2 - 24y + 36 = 0$$

$$x^2 + y^2 - 8y + 12 = 0$$

$$x^2 + (y-4)^2 = 4$$

Circle of radius 2 and center $(0, 4)$.

88. Distance from the origin = $K \times$ Distance from $(2, 0)$

$$\sqrt{x^2 + y^2} = K\sqrt{(x-2)^2 + y^2}, K \neq 1$$

$$x^2 + y^2 = K^2(x^2 - 4x + 4 + y^2)$$

$$(1 - K^2)x^2 + (1 - K^2)y^2 + 4K^2x - 4K^2 = 0$$

Note: This is the equation of a circle!

Section P.2 Linear Models and Rates of Change

1. $m = 1$


2. $m = 2$

3. $m = 0$


4. $m = -1$

5. $m = -12$


6. $m = \frac{40}{3}$




8.



9. $m = \frac{2 - (-4)}{5 - 3} = \frac{6}{2} = 3$




10. $m = \frac{7-1}{-2-1} = \frac{6}{-3} = -2$




11. $m = \frac{1-6}{4-4} = \frac{-5}{0}$, undefined.

The line is vertical




12. $m = \frac{-5 - (-5)}{5 - 3} = \frac{0}{2} = 0$


The line is horizontal



13. $m = \frac{\frac{2}{3} - \frac{1}{6}}{-\frac{1}{2} - (-\frac{3}{4})} = \frac{\frac{1}{2}}{\frac{1}{4}} = 2$



14. $m = \frac{(\frac{3}{4}) - (-\frac{1}{4})}{(\frac{7}{8}) - (\frac{5}{4})} = \frac{1}{-\frac{3}{8}} = -\frac{8}{3}$



15. Because the slope is 0, the line is horizontal and its equation is $y = 2$. Therefore, three additional points are $(0, 2)$, $(1, 2)$, $(5, 2)$.

16. Because the slope is undefined, the line is vertical and its equation is $x = -4$. Therefore, three additional points are $(-4, 0)$, $(-4, 1)$, $(-4, 2)$.

17. The equation of this line is

$$y - 7 = -3(x - 1)$$

$$y = -3x + 10.$$

Therefore, three additional points are $(0, 10)$, $(2, 4)$, and $(3, 1)$.


18. The equation of this line is

$$y + 2 = 2(x + 2)$$

$$y = 2x + 2.$$

Therefore, three additional points are $(-3, -4)$, $(-1, 0)$, and $(0, 2)$.

19. (a) Slope = $\frac{\Delta y}{\Delta x} = \frac{1}{3}$




By the Pythagorean Theorem,

$$x^2 = 30^2 + 10^2 = 1000$$

$$x = 10\sqrt{10} \approx 31.623 \text{ feet.}$$

20. (a) $m = 800$ indicates that the revenues increase by 800 in one day.
 (b) $m = 250$ indicates that the revenues increase by 250 in one day.
 (c) $m = 0$ indicates that the revenues do not change from one day to the next.

21. (a)



(b) The slopes are:

$$\frac{285.3 - 282.4}{1 - 0} = 2.9$$

$$\frac{288.2 - 285.3}{2 - 1} = 2.9$$


$$\frac{291.1 - 288.2}{3 - 2} = 2.9$$

$$\frac{293.9 - 291.1}{4 - 3} = 2.8$$

$$\frac{296.6 - 293.9}{5 - 4} = 2.7$$

The population increased least rapidly from 2004 to 2005.

22. (a)



(b) The slopes are:

$$\frac{74 - 57}{10 - 5} = 3.4$$

$$\frac{85 - 74}{15 - 10} = 2.2$$

$$\frac{84 - 85}{20 - 15} = -0.2$$

$$\frac{61 - 84}{25 - 20} = -4.6$$

$$\frac{43 - 61}{30 - 25} = -3.6$$

The rate changed most rapidly between 20 and 25 seconds. The change is -4.6 mi/h/sec.

23. $y = 4x - 3$

the slope is $m = 4$ and the y -intercept is $(0, -3)$.

24. $-x + y = 1$

$$y = x + 1$$

The slope is $m = 1$ and the y -intercept is $(0, 1)$.

25. $x + 5y = 20$

$$y = -\frac{1}{5}x + 4$$

Therefore, the slope is $m = -\frac{1}{5}$ and the y -intercept is $(0, 4)$.

26. $6x - 5y = 15$

$$y = \frac{6}{5}x - 3$$

Therefore, the slope is $m = \frac{6}{5}$ and the y -intercept is $(0, -3)$.

27. $x = 4$

The line is vertical. Therefore, the slope is undefined and there is no y -intercept.


28. $y = -1$

The line is horizontal. Therefore, the slope is $m = 0$ and the y -intercept is $(0, -1)$.

29. $y = \frac{3}{4}x + 3$

$$4y = 3x + 12$$


$$0 = 3x - 4y + 12$$



30. The slope is undefined so the line is vertical.

$$x = -5$$


$$x + 5 = 0$$



31. $y = \frac{2}{3}x$


$3y = 2x$

$0 = 2x - 3y$



32. $y = 4$

$y - 4 = 0$




33. $y + 2 = 3(x - 3)$

$y + 2 = 3x - 9$

$y = 3x - 11$


$0 = 3x - y - 11$



34. $y - 4 = -\frac{3}{5}(x + 2)$

$5y - 20 = -3x - 6$

$3x + 5y - 14 = 0$




35. $m = \frac{8 - 0}{4 - 0} = 2$

$y - 0 = 2(x - 0)$

$y = 2x$

$0 = 2x - y$




36. $m = \frac{5 - 0}{-1 - 0} = -5$

$y - 0 = -5(x - 0)$

$y = -5x$

$5x + y = 0$




37. $m = \frac{1 - (-3)}{2 - 0} = 2$

$y - 1 = 2(x - 2)$

$y - 1 = 2x - 4$

$0 = 2x - y - 3$




38. $m = \frac{7 - (-2)}{1 - (-2)} = \frac{9}{3} = 3$

$$y - (-2) = 3(x - (-2))$$

$$y + 2 = 3(x + 2)$$

$$y = 3x + 4$$

$$0 = 3x - y + 4$$




39. $m = \frac{8 - 0}{2 - 5} = -\frac{8}{3}$

$$y - 0 = -\frac{8}{3}(x - 5)$$

$$y = -\frac{8}{3}x + \frac{40}{3}$$

$$8x + 3y - 40 = 0$$




40. $m = \frac{6 - 2}{-3 - 1} = \frac{4}{-4} = -1$

$$y - 2 = -1(x - 1)$$

$$y - 2 = -x + 1$$

$$x + y - 3 = 0$$




41. $m = \frac{8 - 3}{6 - 6} = \frac{5}{0}$, undefined

The line is horizontal.

$$x = 6$$


$$x - 6 = 0$$



42. $m = \frac{-2 - (-2)}{3 - 1} = \frac{0}{2} = 0$

$$y = -2$$

$$y + 2 = 0$$




43. $m = \frac{\frac{7}{2} - \frac{3}{4}}{\frac{1}{2} - 0} = \frac{\frac{11}{4}}{\frac{1}{2}} = \frac{11}{2}$

$$y - \frac{3}{4} = \frac{11}{2}(x - 0)$$

$$y = \frac{11}{2}x + \frac{3}{4}$$

$$0 = 22x - 4y + 3$$




$$44. m = \frac{\left(\frac{3}{4}\right) - \left(-\frac{1}{4}\right)}{\left(\frac{7}{8}\right) - \left(\frac{5}{4}\right)} = \frac{1}{-\frac{3}{8}} = -\frac{8}{3}$$

$$y + \frac{1}{4} = -\frac{8}{3}\left(x - \frac{5}{4}\right)$$


$$12y + 3 = -32x + 40$$

$$32x + 12y - 37 = 0$$



$$45. x = 3$$

$$x - 3 = 0$$




$$46. m = -\frac{b}{a}$$

$$y = -\frac{b}{a}x + b$$

$$\frac{b}{a}x + y = b$$

$$\frac{x}{a} + \frac{y}{b} = 1$$



$$47. \frac{x}{2} + \frac{y}{3} = 1$$

$$3x + 2y - 6 = 0$$

$$48. \frac{x}{2} + \frac{y}{-2} = 1$$

$$\frac{-3x}{2} - \frac{y}{2} = 1$$

$$3x + y = -2$$

$$3x + y + 2 = 0$$

$$49. \frac{x}{a} + \frac{y}{a} = 1$$

$$\frac{1}{a} + \frac{2}{a} = 1$$

$$\frac{3}{a} = 1$$

$$a = 3 \Rightarrow x + y = 3$$

$$x + y - 3 = 0$$

$$50. \frac{x}{a} + \frac{y}{a} = 1$$


$$\frac{-3}{a} + \frac{4}{a} = 1$$

$$\frac{1}{a} = 1$$


$$a = 1 \Rightarrow x + y = 1$$

$$x + y - 1 = 0$$


$$51. y = -3$$




$$52. x = 4$$



$$53. y = -2x + 1$$




54. $y = \frac{1}{3}x - 1$




55. $y - 2 = \frac{3}{2}(x - 1)$

$$y = \frac{3}{2}x + \frac{1}{2}$$




56. $y - 1 = 3(x + 4)$

$$y = 3x + 13$$




57. $2x - y - 3 = 0$

$$y = 2x - 3$$




58. $x + 2y + 6 = 0$

$$y = -\frac{1}{2}x - 3$$




59. (a)



The lines do not appear perpendicular.

(b)



The lines appear perpendicular.

The lines are perpendicular because their slopes 2 and $-\frac{1}{2}$ are negative reciprocals of each other. You must use a square setting in order for perpendicular lines to appear perpendicular. Answers depend on calculator used.

60. $ax + by = 4$

(a) The line is parallel to the x -axis if $a = 0$ and $b \neq 0$.

(b) The line is parallel to the y -axis if $b = 0$ and $a \neq 0$.

(c) Answers will vary. *Sample answer:* $a = -5$ and $b = 8$.

$$-5x + 8y = 4$$

$$y = \frac{1}{8}(5x + 4) = \frac{5}{8}x + \frac{1}{2}$$

(d) The slope must be $-\frac{5}{2}$.

Answers will vary. *Sample answer:* $a = 5$ and $b = 2$.

$$5x + 2y = 4$$

$$y = \frac{1}{2}(-5x + 4) = -\frac{5}{2}x + 2$$

(e) $a = \frac{5}{2}$ and $b = 3$.

$$\frac{5}{2}x + 3y = 4$$

$$5x + 6y = 8$$

61. The given line is vertical.

(a) $x = -7$, or $x + 7 = 0$

(b) $y = -2$, or $y + 2 = 0$

62. The given line is horizontal.

(a) $y = 0$

(b) $x = -1$, or $x + 1 = 0$

63. $4x - 2y = 3$

$$y = 2x - \frac{3}{2}$$

$$m = 2$$

(a) $y - 1 = 2(x - 2)$

$$y - 1 = 2x - 4$$

$$0 = 2x - y - 3$$

(b) $y - 1 = -\frac{1}{2}(x - 2)$

$$2y - 2 = -x + 2$$

$$x + 2y - 4 = 0$$

64. $x + y = 7$

$$y = -x + 7$$

$$m = -1$$

(a) $y - 2 = -1(x + 3)$

$$y - 2 = -x - 3$$

$$x + y + 1 = 0$$

(b) $y - 2 = 1(x + 3)$

$$y - 2 = x + 3$$

$$0 = x - y + 5$$

65. $5x - 3y = 0$

$$y = \frac{5}{3}x$$

$$m = \frac{5}{3}$$

(a) $y - \frac{7}{8} = \frac{5}{3}(x - \frac{3}{4})$

$$24y - 21 = 40x - 30$$

$$0 = 40x - 24y - 9$$

(b) $y - \frac{7}{8} = -\frac{3}{5}(x - \frac{3}{4})$

$$40y - 35 = -24x + 18$$

$$24x + 40y - 53 = 0$$

66. $3x + 4y = 7$

$$4y = -3x + 7$$

$$y = -\frac{3}{4}x + \frac{7}{4}$$

$$m = -\frac{3}{4}$$

(a) $y - (-5) = -\frac{3}{4}(x - 4)$

$$y + 5 = -\frac{3}{4}x + 3$$

$$4y + 20 = -3x + 12$$

$$3x + 4y + 8 = 0$$

(b) $y - (-5) = \frac{4}{3}(x - 4)$

$$y + 5 = \frac{4}{3}x - \frac{16}{3}$$

$$3y + 15 = 4x - 16$$

$$0 = 4x - 3y - 31$$

67. The slope is 250. $V = 1850$ when $t = 8$.

$$V = 250(t - 8) + 1850 = 250t - 150.$$

68. The slope is 4.5. $V = 156$ when $t = 4$.

$$V = 4.5(t - 4) + 156 = 4.5t + 138$$


69. The slope is -1600. $V = 17,200$ when $t = 8$.

$$V = -1600(t - 8) + 17,200 = -1600t + 30,000$$

70. The slope is -5600. $V = 245,000$ when $t = 4$.

$$V = -5600(t - 4) + 245,000 = -5600t + 267,400$$

71.



You can use the graphing utility to determine that the points of intersection are $(0, 0)$ and $(2, 4)$. Analytically,

$$x^2 = 4x - x^2$$

$$2x^2 - 4x = 0$$

$$2x(x - 2) = 0$$

$$x = 0 \Rightarrow y = 0 \Rightarrow (0, 0)$$

$$x = 2 \Rightarrow y = 4 \Rightarrow (2, 4).$$


The slope of the line joining $(0, 0)$ and $(2, 4)$ is

$$m = (4 - 0)/(2 - 0) = 2. \text{ So, an equation of the line is}$$

$$y - 0 = 2(x - 0)$$

$$y = 2x.$$

72. $y = x^2 - 4x + 3$, $y = -x^2 + 2x + 3$



You can use the graphing utility to determine that the points of intersection are $(0, 3)$ and $(3, 0)$. Analytically,

$$x^2 - 4x + 3 = -x^2 + 2x + 3$$

$$2x^2 - 6x = 0$$

$$2x(x - 3) = 0$$

$$x = 0 \Rightarrow y = 3 \Rightarrow (0, 3)$$

$$x = 3 \Rightarrow y = 0 \Rightarrow (3, 0).$$

The slope of the line joining $(0, 3)$ and $(3, 0)$ is

$$m = (0 - 3)/(3 - 0) = -1. \text{ So, an equation of the line is}$$

$$y - 3 = -1(x - 0)$$

$$y = -x + 3.$$

73. $m_1 = \frac{1 - 0}{-2 - (-1)} = -1$

$m_2 = \frac{-2 - 0}{2 - (-1)} = -\frac{2}{3}$

$m_1 \neq m_2$

The points are not collinear.

74. $m_1 = \frac{-6 - 4}{7 - 0} = -\frac{10}{7}$

$m_2 = \frac{11 - 4}{-5 - 0} = -\frac{7}{5}$

$m_1 \neq m_2$

The points are not collinear.

75. Equations of perpendicular bisectors:

$$y - \frac{c}{2} = \frac{a - b}{c} \left(x - \frac{a + b}{2} \right)$$


$$y - \frac{c}{2} = \frac{a + b}{-c} \left(x - \frac{b - a}{2} \right)$$

Setting the right-hand sides of the two equations equal and solving for x yields $x = 0$.

Letting $x = 0$ in either equation gives the point of intersection:

$$\left(0, \frac{-a^2 + b^2 + c^2}{2c} \right)$$

This point lies on the third perpendicular bisector, $x = 0$.




76. Equations of medians:

$$y = \frac{c}{b}x$$

$$y = \frac{c}{3a + b}(x + a)$$

$$y = \frac{c}{-3a + b}(x - a)$$

Solving simultaneously, the point of intersection is $\left(\frac{b}{3}, \frac{c}{3} \right)$.



77. Equations of altitudes:


$$y = \frac{a - b}{c}(x + a)$$

$$x = b$$

$$y = -\frac{a + b}{c}(x - a)$$

Solving simultaneously, the point of intersection is

$$\left(b, \frac{a^2 - b^2}{c} \right)$$



78. The slope of the line segment from $\left(\frac{b}{3}, \frac{c}{3} \right)$ to

$\left(b, \frac{a^2 - b^2}{c} \right)$ is:

$$m_1 = \frac{\left[\frac{a^2 - b^2}{c} \right] - (c/3)}{b - (b/3)} = \frac{(3a^2 - 3b^2 - c^2)/(3c)}{(2b)/3} = \frac{3a^2 - 3b^2 - c^2}{2bc}$$

The slope of the line segment from $\left(\frac{b}{3}, \frac{c}{3} \right)$ to

$\left(0, \frac{-a^2 + b^2 + c^2}{2c} \right)$ is:

$$m_2 = \frac{\left[\frac{-a^2 + b^2 + c^2}{2c} \right] - (c/3)}{0 - (b/3)} = \frac{(-3a^2 + 3b^2 + 3c^2 - 2c^2)/(6c)}{-b/3} = \frac{3a^2 - 3b^2 - c^2}{2bc}$$

$m_1 = m_2$

Therefore, the points are collinear.

79. Find the equation of the line through the points (0, 32) and (100, 212).

$$m = \frac{180}{100} = \frac{9}{5}$$

$$F - 32 = \frac{9}{5}(C - 0)$$

$$F = \frac{9}{5}C + 32$$

or

$$C = \frac{1}{9}(5F - 160)$$


$$5F - 9C - 160 = 0$$

For $F = 72^\circ$, $C \approx 22.2^\circ$.

80. $C = 0.48x + 175$

For $x = 137$, $C = 0.48(137) + 175 = \$240.76$.

81. (a) $W_1 = 0.75x + 14.50$
 $W_2 = 1.30x + 11.20$



Using a graphing utility, the point of intersection is (6, 19)

Analytically,

$$W_1 = W_2$$

$$0.75x + 14.50 = 1.30x + 11.20$$

$$3.3 = 0.55x$$

$$6 = x$$

$$y = 1.30(6) + 11.20 = 19.$$


(c) When six units are produced, the wage for both options is \$19.00 per hour. Choose option 1 if you think you will produce less than six units per hour, and choose option 2 if you think you will produce more than six.

82. (a) Depreciation per year:

$$\frac{875}{5} = \$175$$

$$y = 875 - 175x$$

where $0 \leq x \leq 5$.



(b) $y = 875 - 175(2) = \$525$

(c) $200 = 875 - 175x$
 $175x = 675$
 $x \approx 3.86$ years

83. (a) Two points are (50, 780) and (47, 825).

The slope is


$$m = \frac{825 - 780}{47 - 50} = \frac{45}{-3} = -15.$$

$$p - 780 = -15(x - 50)$$

$$p = -15x + 750 + 780 = -15x + 1530$$

or


$$x = \frac{1}{15}(1530 - p)$$



If $p = 855$, then $x = 45$ units.

(c) If $p = 795$, then $x = \frac{1}{15}(1530 - 795) = 49$ units.

84. (a) $y = 18.91 + 3.97x$
 ($x =$ quiz score, $y =$ test score)




(c) If $x = 17$, $y = 18.91 + 3.97(17) = 86.4$.

(d) The slope shows the average increase in exam score for each unit increase in quiz score.

(e) The points would shift vertically upward 4 units. The new regression line would have a y -intercept 4 greater than before: $y = 22.91 + 3.97x$.

85. The tangent line is perpendicular to the line joining the point (5, 12) and the center (0, 0).



Slope of the line joining (5, 12) and (0, 0) is $\frac{12}{5}$.


The equation of the tangent line is

$$y - 12 = \frac{-5}{12}(x - 5)$$

$$y = \frac{-5}{12}x + \frac{169}{12}$$

$$5x + 12y - 169 = 0.$$

86. The tangent line is perpendicular to the line joining the point (4, -3) and the center of the circle, (1, 1).



Slope of the line joining (1, 1) and (4, -3) is $\frac{1 + 3}{1 - 4} = \frac{-4}{3}$.

Tangent line:

$$y + 3 = \frac{3}{4}(x - 4)$$

$$y = \frac{3}{4}x - 6$$

$$0 = 3x - 4y - 24$$

87. $4x + 3y - 10 = 0 \Rightarrow d = \frac{|4(0) + 3(0) - 10|}{\sqrt{4^2 + 3^2}}$

$$= \frac{10}{5} = 2$$

88. $4x + 3y - 10 = 0 \Rightarrow d = \frac{|4(2) + 3(3) - 10|}{\sqrt{4^2 + 3^2}} = \frac{7}{5}$

89. $x - y - 2 = 0 \Rightarrow d = \frac{|1(-2) + (-1)(1) - 2|}{\sqrt{1^2 + 1^2}} = \frac{5}{\sqrt{2}} = \frac{5\sqrt{2}}{2}$

90. $x + 1 = 0 \Rightarrow d = \frac{|1(6) + (0)(2) + 1|}{\sqrt{1^2 + 0^2}} = 7$

91. A point on the line $x + y = 1$ is $(0, 1)$. The distance from the point $(0, 1)$ to $x + y - 5 = 0$ is

$$d = \frac{|1(0) + 1(1) - 5|}{\sqrt{1^2 + 1^2}} = \frac{|1 - 5|}{\sqrt{2}} = \frac{4}{\sqrt{2}} = 2\sqrt{2}.$$

92. A point on the line $3x - 4y = 1$ is $(-1, -1)$. The distance from the point $(-1, -1)$ to $3x - 4y - 10 = 0$ is

$$d = \frac{|3(-1) - 4(-1) - 10|}{\sqrt{3^2 + (-4)^2}} = \frac{|-3 + 4 - 10|}{5} = \frac{9}{5}.$$

93. If $A = 0$, then $By + C = 0$ is the horizontal line $y = -C/B$. The distance to (x_1, y_1) is

$$d = \left| y_1 - \left(\frac{-C}{B} \right) \right| = \frac{|By_1 + C|}{|B|} = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}}.$$

If $B = 0$, then $Ax + C = 0$ is the vertical line $x = -C/A$. The distance to (x_1, y_1) is

$$d = \left| x_1 - \left(\frac{-C}{A} \right) \right| = \frac{|Ax_1 + C|}{|A|} = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}}.$$

(Note that A and B cannot both be zero.) The slope of the line $Ax + By + C = 0$ is $-A/B$.

The equation of the line through (x_1, y_1) perpendicular to $Ax + By + C = 0$ is:

$$y - y_1 = \frac{B}{A}(x - x_1)$$

$$Ay - Ay_1 = Bx - Bx_1$$

$$Bx_1 - Ay_1 = Bx - Ay$$

The point of intersection of these two lines is:

$$Ax + By = -C \quad \Rightarrow \quad A^2x + ABY = -AC \quad (1)$$

$$Bx - Ay = Bx_1 - Ay_1 \Rightarrow B^2x - ABY = B^2x_1 - ABY_1 \quad (2)$$

$$(A^2 + B^2)x = -AC + B^2x_1 - ABY_1 \quad (\text{By adding equations (1) and (2)})$$

$$x = \frac{-AC + B^2x_1 - ABY_1}{A^2 + B^2}$$

$$Ax + By = -C \quad \Rightarrow \quad ABx + B^2y = -BC \quad (3)$$

$$Bx - Ay = Bx_1 - Ay_1 \Rightarrow -ABx + A^2y = -ABx_1 + A^2y_1 \quad (4)$$

$$(A^2 + B^2)y = -BC - ABx_1 + A^2y_1 \quad (\text{By adding equations (3) and (4)})$$

$$y = \frac{-BC - ABx_1 + A^2y_1}{A^2 + B^2}$$

$$\left(\frac{-AC + B^2x_1 - ABY_1}{A^2 + B^2}, \frac{-BC - ABx_1 + A^2y_1}{A^2 + B^2} \right) \text{ point of intersection}$$


The distance between (x_1, y_1) and this point gives you the distance between (x_1, y_1) and the line $Ax + By + C = 0$.

$$\begin{aligned} d &= \sqrt{\left[\frac{-AC + B^2x_1 - ABY_1}{A^2 + B^2} - x_1 \right]^2 + \left[\frac{-BC - ABx_1 + A^2y_1}{A^2 + B^2} - y_1 \right]^2} \\ &= \sqrt{\left[\frac{-AC - ABY_1 - A^2x_1}{A^2 + B^2} \right]^2 + \left[\frac{-BC - ABx_1 - B^2y_1}{A^2 + B^2} \right]^2} \\ &= \sqrt{\left[\frac{-A(C + By_1 + Ax_1)}{A^2 + B^2} \right]^2 + \left[\frac{-B(C + Ax_1 + By_1)}{A^2 + B^2} \right]^2} = \sqrt{\frac{(A^2 + B^2)(C + Ax_1 + By_1)^2}{(A^2 + B^2)^2}} = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}} \end{aligned}$$

94. $y = mx + 4 \Rightarrow mx + (-1)y + 4 = 0$

$$d = \frac{|Ax_1 + By_1 + C|}{\sqrt{A^2 + B^2}} = \frac{|m(3) + (-1)(1) + 4|}{\sqrt{m^2 + (-1)^2}} = \frac{|3m + 3|}{\sqrt{m^2 + 1}}$$

The distance is 0 when $m = -1$. In this case, the line $y = -x + 4$ contains the point $(3, 1)$.



95. For simplicity, let the vertices of the rhombus be $(0, 0)$, $(a, 0)$, (b, c) , and $(a + b, c)$, as shown in the figure. The


slopes of the diagonals are then $m_1 = \frac{c}{a + b}$ and

$m_2 = \frac{c}{b - a}$. Because the sides of the rhombus are

equal, $a^2 = b^2 + c^2$, and you have

$$m_1 m_2 = \frac{c}{a + b} \cdot \frac{c}{b - a} = \frac{c^2}{b^2 - a^2} = \frac{c^2}{-c^2} = -1.$$

Therefore, the diagonals are perpendicular.



96. For simplicity, let the vertices of the quadrilateral be $(0, 0)$, $(a, 0)$, (b, c) , and (d, e) , as shown in the figure. The midpoints of the sides are


$$\left(\frac{a}{2}, 0\right), \left(\frac{a+b}{2}, \frac{c}{2}\right), \left(\frac{b+d}{2}, \frac{c+e}{2}\right), \text{ and } \left(\frac{d}{2}, \frac{e}{2}\right).$$

The slope of the opposite sides are equal:

$$\frac{\frac{c}{2} - 0}{\frac{a+b}{2} - \frac{a}{2}} = \frac{\frac{c+e}{2} - \frac{e}{2}}{\frac{b+d}{2} - \frac{d}{2}} = \frac{c}{b}$$


$$\frac{0 - \frac{e}{2}}{\frac{a}{2} - \frac{d}{2}} = \frac{\frac{c}{2} - \frac{c+e}{2}}{\frac{a+b}{2} - \frac{b+d}{2}} = -\frac{e}{a-d}$$

Therefore, the figure is a parallelogram.



97. Consider the figure below in which the four points are collinear. Because the triangles are similar, the result immediately follows.

$$\frac{y_2^* - y_1^*}{x_2^* - x_1^*} = \frac{y_2 - y_1}{x_2 - x_1}$$



98. If $m_1 = -1/m_2$, then $m_1 m_2 = -1$. Let L_3 be a line with slope m_3 that is perpendicular to L_1 . Then $m_1 m_3 = -1$.

So, $m_2 = m_3 \Rightarrow L_2$ and L_3 are parallel. Therefore, L_2 and L_1 are also perpendicular.

99. True.

$$ax + by = c_1 \Rightarrow y = -\frac{a}{b}x + \frac{c_1}{b} \Rightarrow m_1 = -\frac{a}{b}$$

$$bx - ay = c_2 \Rightarrow y = \frac{b}{a}x - \frac{c_2}{a} \Rightarrow m_2 = \frac{b}{a}$$

$$m_2 = -\frac{1}{m_1}$$

100. False; if m_1 is positive, then $m_2 = -1/m_1$ is negative.

Section P.3 Functions and Their Graphs

1. (a) Domain of f : $-4 \leq x \leq 4 \Rightarrow [-4, 4]$
 Range of f : $-3 \leq y \leq 5 \Rightarrow [-3, 5]$
 Domain of g : $-3 \leq x \leq 3 \Rightarrow [-3, 3]$
 Range of g : $-4 \leq y \leq 4 \Rightarrow [-4, 4]$
 - (b) $f(-2) = -1$
 $g(3) = -4$
 - (c) $f(x) = g(x)$ for $x = -1$
 - (d) $f(x) = 2$ for $x = 1$
 - (e) $g(x) = 0$ for $x = -1, 1$ and 2
2. (a) Domain of f : $-5 \leq x \leq 5 \Rightarrow [-5, 5]$
 Range of f : $-4 \leq y \leq 4 \Rightarrow [-4, 4]$
 Domain of g : $-4 \leq x \leq 5 \Rightarrow [-4, 5]$
 Range of g : $-4 \leq y \leq 2 \Rightarrow [-4, 2]$
 - (b) $f(-2) = -2$
 $g(3) = 2$
 - (c) $f(x) = g(x)$ for $x = -2$ and $x = 4$
 - (d) $f(x) = 2$ for $x = -4, 4$
 - (e) $g(x) = 0$ for $x = -1$
3. (a) $f(0) = 7(0) - 4 = -4$
 (b) $f(-3) = 7(-3) - 4 = -25$
 (c) $f(b) = 7(b) - 4 = 7b - 4$
 (d) $f(x - 1) = 7(x - 1) - 4 = 7x - 11$
4. (a) $f(-4) = \sqrt{-4 + 5} = \sqrt{1} = 1$
 (b) $f(11) = \sqrt{11 + 5} = \sqrt{16} = 4$
 (c) $f(-8) = \sqrt{-8 + 5} = \sqrt{-3}$, undefined
 (d) $f(x + \Delta x) = \sqrt{x + \Delta x + 5}$
5. (a) $g(0) = 5 - 0^2 = 5$
 (b) $g(\sqrt{5}) = 5 - (\sqrt{5})^2 = 5 - 5 = 0$
 (c) $g(-2) = 5 - (-2)^2 = 5 - 4 = 1$
 (d) $g(t - 1) = 5 - (t - 1)^2 = 5 - (t^2 - 2t + 1)$
 $= 4 + 2t - t^2$
6. (a) $g(4) = 4^2(4 - 4) = 0$
 (b) $g\left(\frac{3}{2}\right) = \left(\frac{3}{2}\right)^2\left(\frac{3}{2} - 4\right) = \frac{9}{4}\left(-\frac{5}{2}\right) = -\frac{45}{8}$
 (c) $g(c) = c^2(c - 4) = c^3 - 4c^2$
 (d) $g(t + 4) = (t + 4)^2(t + 4 - 4)$
 $= (t + 4)^2 t = t^3 + 8t^2 + 16t$
7. (a) $f(0) = \cos(2(0)) = \cos 0 = 1$
 (b) $f\left(-\frac{\pi}{4}\right) = \cos\left(2\left(-\frac{\pi}{4}\right)\right) = \cos\left(-\frac{\pi}{2}\right) = 0$
 (c) $f\left(\frac{\pi}{3}\right) = \cos\left(2\left(\frac{\pi}{3}\right)\right) = \cos\frac{2\pi}{3} = -\frac{1}{2}$
8. (a) $f(\pi) = \sin \pi = 0$
 (b) $f\left(\frac{5\pi}{4}\right) = \sin\left(\frac{5\pi}{4}\right) = \frac{-\sqrt{2}}{2}$
 (c) $f\left(\frac{2\pi}{3}\right) = \sin\left(\frac{2\pi}{3}\right) = \frac{\sqrt{3}}{2}$
9. $\frac{f(x + \Delta x) - f(x)}{\Delta x} = \frac{(x + \Delta x)^3 - x^3}{\Delta x} = \frac{x^3 + 3x^2\Delta x + 3x^2(\Delta x)^2 + (\Delta x)^3 - x^3}{\Delta x} = 3x^2 + 3x\Delta x + (\Delta x)^2, \Delta x \neq 0$
10. $\frac{f(x) - f(1)}{x - 1} = \frac{3x - 1 - (3 - 1)}{x - 1} = \frac{3(x - 1)}{x - 1} = 3, x \neq 1$
11. $\frac{f(x) - f(2)}{x - 2} = \frac{(1/\sqrt{x-1}) - 1}{x - 2}$
 $= \frac{1 - \sqrt{x-1}}{(x-2)\sqrt{x-1}} \cdot \frac{1 + \sqrt{x-1}}{1 + \sqrt{x-1}} = \frac{2 - x}{(x-2)\sqrt{x-1}(1 + \sqrt{x-1})} = \frac{-1}{\sqrt{x-1}(1 + \sqrt{x-1})}, x \neq 2$
12. $\frac{f(x) - f(1)}{x - 1} = \frac{x^3 - x - 0}{x - 1} = \frac{x(x+1)(x-1)}{x - 1} = x(x+1), x \neq 1$

13. $f(x) = 4x^2$

Domain: $(-\infty, \infty)$

Range: $[0, \infty)$

14. $g(x) = x^2 - 5$

Domain: $(-\infty, \infty)$

Range: $[-5, \infty)$

15. $g(x) = \sqrt{6x}$

Domain: $6x \geq 0$

$x \geq 0 \Rightarrow [0, \infty)$

Range: $[0, \infty)$

16. $h(x) = -\sqrt{x+3}$

Domain: $x+3 \geq 0 \Rightarrow [-3, \infty)$

Range: $(-\infty, 0]$

17. $f(t) = \sec \frac{\pi t}{4}$

$\frac{\pi t}{4} \neq \frac{(2n+1)\pi}{2} \Rightarrow t \neq 4n+2$

Domain: all $t \neq 4n+2$, n an integer

Range: $(-\infty, -1] \cup [1, \infty)$

18. $h(t) = \cot t$

Domain: all $t = n\pi$, n an integer

Range: $(-\infty, \infty)$

19. $f(x) = \frac{3}{x}$

Domain: all $x \neq 0 \Rightarrow (-\infty, 0) \cup (0, \infty)$

Range: $(-\infty, 0) \cup (0, \infty)$

20. $g(x) = \frac{2}{x-1}$

Domain: $(-\infty, 1) \cup (1, \infty)$

Range: $(-\infty, 0) \cup (0, \infty)$

21. $f(x) = \sqrt{x} + \sqrt{1-x}$

$x \geq 0$ and $1-x \geq 0$

$x \geq 0$ and $x \leq 1$

Domain: $0 \leq x \leq 1 \Rightarrow [0, 1]$

22. $f(x) = \sqrt{x^2 - 3x + 2}$

$x^2 - 3x + 2 \geq 0$

$(x-2)(x-1) \geq 0$

Domain: $x \geq 2$ or $x \leq 1$

Domain: $(-\infty, 1] \cup [2, \infty)$

23. $g(x) = \frac{2}{1 - \cos x}$

$1 - \cos x \neq 0$

$\cos x \neq 1$

Domain: all $x \neq 2n\pi$, n an integer

24. $h(x) = \frac{1}{\sin x - (1/2)}$

$\sin x - \frac{1}{2} \neq 0$

$\sin x \neq \frac{1}{2}$

Domain: all $x \neq \frac{\pi}{6} + 2n\pi, \frac{5\pi}{6} + 2n\pi$, n integer

25. $f(x) = \frac{1}{|x+3|}$

$|x+3| \neq 0$

$x+3 \neq 0$

Domain: all $x \neq -3$

Domain: $(-\infty, -3) \cup (-3, \infty)$

26. $g(x) = \frac{1}{|x^2 - 4|}$

$|x^2 - 4| \neq 0$

$(x-2)(x+2) \neq 0$

Domain: all $x \neq \pm 2$

Domain: $(-\infty, -2) \cup (-2, \infty)$

27. $f(x) = \begin{cases} 2x+1, & x < 0 \\ 2x+2, & x \geq 0 \end{cases}$

(a) $f(-1) = 2(-1) + 1 = -1$

(b) $f(0) = 2(0) + 2 = 2$

(c) $f(2) = 2(2) + 2 = 6$

(d) $f(t^2 + 1) = 2(t^2 + 1) + 2 = 2t^2 + 4$

(Note: $t^2 + 1 \geq 0$ for all t)

Domain: $(-\infty, \infty)$

Range: $(-\infty, 1) \cup [2, \infty)$

28. $f(x) = \begin{cases} x^2 + 2, & x \leq 1 \\ 2x^2 + 2, & x > 1 \end{cases}$

- (a) $f(-2) = (-2)^2 + 2 = 6$
- (b) $f(0) = 0^2 + 2 = 2$
- (c) $f(1) = 1^2 + 2 = 3$
- (d) $f(s^2 + 2) = 2(s^2 + 2)^2 + 2 = 2s^4 + 8s^2 + 10$

(Note: $s^2 + 2 > 1$ for all s)

Domain: $(-\infty, \infty)$

Range: $[2, \infty)$

29. $f(x) = \begin{cases} |x| + 1, & x < 1 \\ -x + 1, & x \geq 1 \end{cases}$

- (a) $f(-3) = |-3| + 1 = 4$
- (b) $f(1) = -1 + 1 = 0$
- (c) $f(3) = -3 + 1 = -2$
- (d) $f(b^2 + 1) = -(b^2 + 1) + 1 = -b^2$

Domain: $(-\infty, \infty)$

Range: $(-\infty, 0] \cup [1, \infty)$

30. $f(x) = \begin{cases} \sqrt{x+4}, & x \leq 5 \\ (x-5)^2, & x > 5 \end{cases}$

- (a) $f(-3) = \sqrt{-3+4} = \sqrt{1} = 1$
- (b) $f(0) = \sqrt{0+4} = 2$
- (c) $f(5) = \sqrt{5+4} = 3$
- (d) $f(10) = (10-5)^2 = 25$


Domain: $[-4, \infty)$

Range: $[0, \infty)$

31. $f(x) = 4 - x$

Domain: $(-\infty, \infty)$


Range: $(-\infty, \infty)$




32. $g(x) = \frac{4}{x}$

Domain: $(-\infty, 0) \cup (0, \infty)$

Range: $(-\infty, 0) \cup (0, \infty)$



33. $h(x) = \sqrt{x-6}$




Domain: $x - 6 \geq 0$

$x \geq 6 \Rightarrow [6, \infty)$

Range: $[0, \infty)$

34. $f(x) = \frac{1}{4}x^3 + 3$




Domain: $(-\infty, \infty)$

Range: $(-\infty, \infty)$


35. $f(x) = \sqrt{9 - x^2}$

Domain: $[-3, 3]$


Range: $[0, 3]$




36. $f(x) = x + \sqrt{4 - x^2}$

Domain: $[-2, 2]$ Range: $[-2, 2\sqrt{2}] \approx [-2, 2.83]$ y -intercept: $(0, 2)$ x -intercept: $(-\sqrt{2}, 0)$ 


37. $g(t) = 3 \sin \pi t$

Domain: $(-\infty, \infty)$ Range: $[-3, 3]$

38. $h(\theta) = -5 \cos \frac{\theta}{2}$

Domain: $(-\infty, \infty)$ Range: $[-5, 5]$ 

39. The student travels $\frac{2-0}{4-0} = \frac{1}{2}$ mi/min during the first 4 minutes. The student is stationary for the next 2 minutes. Finally, the student travels $\frac{6-2}{10-6} = 1$ mi/min during the final 4 minutes.



40. $x - y^2 = 0 \Rightarrow y = \pm\sqrt{x}$

y is not a function of x . Some vertical lines intersect the graph twice.

42. $\sqrt{x^2 - 4} - y = 0 \Rightarrow y = \sqrt{x^2 - 4}$

y is a function of x . Vertical lines intersect the graph at most once.

- 43.
- y
- is a function of
- x
- . Vertical lines intersect the graph at most once.

44. $x^2 + y^2 = 4$

$$y = \pm\sqrt{4 - x^2}$$

y is not a function of x . Some vertical lines intersect the graph twice.

45. $x^2 + y^2 = 16 \Rightarrow y = \pm\sqrt{16 - x^2}$

y is not a function of x because there are two values of y for some x .

46. $x^2 + y = 16 \Rightarrow y = 16 - x^2$

y is a function of x because there is one value of y for each x .

47. $y^2 = x^2 - 1 \Rightarrow y = \pm\sqrt{x^2 - 1}$

y is not a function of x because there are two values of y for some x .

48. $x^2y - x^2 + 4y = 0 \Rightarrow y = \frac{x^2}{x^2 + 4}$

y is a function of x because there is one value of y for each x .

- 49.
- $y = f(x + 5)$
- is a horizontal shift 5 units to the left. Matches d.

- 50.
- $y = f(x) - 5$
- is a vertical shift 5 units downward. Matches b.


- 51.
- $y = -f(-x) - 2$
- is a reflection in the
- y
- axis, a reflection in the
- x
- axis, and a vertical shift downward 2 units. Matches c.

- 52.
- $y = -f(x - 4)$
- is a horizontal shift 4 units to the right, followed by a reflection in the
- x
- axis. Matches a.


53. $y = f(x + 6) + 2$ is a horizontal shift to the left 6 units, and a vertical shift upward 2 units. Matches e.

54. $y = f(x - 1) + 3$ is a horizontal shift to the right 1 unit, and a vertical shift upward 3 units. Matches g.


55. (a) the graph is shifted 3 units to the left.




(b) The graph is shifted 1 unit to the right.




(c) The graph is shifted 2 units upward.




(d) The graph is shifted 4 units downward.



(e) The graph is stretched vertically by a factor of 3.



(f) The graph is stretched vertically by a factor of $\frac{1}{4}$.




56. (a) $g(x) = f(x - 4)$

$$g(6) = f(2) = 1$$


$$g(0) = f(-4) = -3$$

Shift f right 4 units




(b) $g(x) = f(x + 2)$

Shift f left 2 units




(c) $g(x) = f(x) + 4$

Vertical shift upwards 4 units




(d) $g(x) = f(x) - 1$


Vertical shift down 1 unit




(e) $g(x) = 2f(x)$
 $g(2) = 2f(2) = 2$
 $g(-4) = 2f(-4) = -6$



(f) $g(x) = \frac{1}{2}f(x)$
 $g(2) = \frac{1}{2}f(2) = \frac{1}{2}$
 $g(-4) = \frac{1}{2}f(-4) = -\frac{3}{2}$




57. (a) $y = \sqrt{x} + 2$




Vertical shift 2 units upward

(b) $y = -\sqrt{x}$



Reflection about the x -axis

(c) $y = \sqrt{x - 2}$



Horizontal shift 2 units to the right

58. (a) $h(x) = \sin(x + (\pi/2)) + 1$ is a horizontal shift $\pi/2$ units to the left, followed by a vertical shift 1 unit upwards.

(b) $h(x) = -\sin(x - 1)$ is a horizontal shift 1 unit to the right followed by a reflection about the x -axis.

59. (a) $f(g(1)) = f(0) = 0$

(b) $g(f(1)) = g(1) = 0$

(c) $g(f(0)) = g(0) = -1$

(d) $f(g(-4)) = f(15) = \sqrt{15}$

(e) $f(g(x)) = f(x^2 - 1) = \sqrt{x^2 - 1}$

(f) $g(f(x)) = g(\sqrt{x}) = (\sqrt{x})^2 - 1 = x - 1, (x \geq 0)$

60. $f(x) = \sin x, g(x) = \pi x$

(a) $f(g(2)) = f(2\pi) = \sin(2\pi) = 0$

(b) $f\left(g\left(\frac{1}{2}\right)\right) = f\left(\frac{\pi}{2}\right) = \sin\left(\frac{\pi}{2}\right) = 1$

(c) $g(f(0)) = g(0) = 0$

(d) $g\left(f\left(\frac{\pi}{4}\right)\right) = g\left(\sin\left(\frac{\pi}{4}\right)\right)$
 $= g\left(\frac{\sqrt{2}}{2}\right) = \pi\left(\frac{\sqrt{2}}{2}\right) = \frac{\pi\sqrt{2}}{2}$

(e) $f(g(x)) = f(\pi x) = \sin(\pi x)$

(f) $g(f(x)) = g(\sin x) = \pi \sin x$

61. $f(x) = x^2, g(x) = \sqrt{x}$

$(f \circ g)(x) = f(g(x))$
 $= f(\sqrt{x}) = (\sqrt{x})^2 = x, x \geq 0$

Domain: $[0, \infty)$

$(g \circ f)(x) = g(f(x)) = g(x^2) = \sqrt{x^2} = |x|$

Domain: $(-\infty, \infty)$

No. Their domains are different. $(f \circ g) = (g \circ f)$ for $x \geq 0$.

62. $f(x) = x^2 - 1$, $g(x) = \cos x$
 $(f \circ g)(x) = f(g(x)) = f(\cos x) = \cos^2 x - 1$
 Domain: $(-\infty, \infty)$
 $(g \circ f)(x) = g(x^2 - 1) = \cos(x^2 - 1)$
 Domain: $(-\infty, \infty)$
 No, $f \circ g \neq g \circ f$.


63. $f(x) = \frac{3}{x}$, $g(x) = x^2 - 1$
 $(f \circ g)(x) = f(g(x)) = f(x^2 - 1) = \frac{3}{x^2 - 1}$
 Domain: all $x \neq \pm 1 \Rightarrow (-\infty, -1) \cup (-1, 1) \cup (1, \infty)$
 $(g \circ f)(x) = g(f(x)) = g\left(\frac{3}{x}\right) = \left(\frac{3}{x}\right)^2 - 1 = \frac{9}{x^2} - 1 = \frac{9 - x^2}{x^2}$
 Domain: all $x \neq 0 \Rightarrow (-\infty, 0) \cup (0, \infty)$
 No, $f \circ g \neq g \circ f$.

64. $(f \circ g)(x) = f(\sqrt{x+2}) = \frac{1}{\sqrt{x+2}}$

Domain: $(-2, \infty)$

$(g \circ f)(x) = g\left(\frac{1}{x}\right) = \sqrt{\frac{1}{x} + 2} = \sqrt{\frac{1 + 2x}{x}}$

You can find the domain of $g \circ f$ by determining the intervals where $(1 + 2x)$ and x are both positive, or both negative.



Domain: $(-\infty, -\frac{1}{2}] \cup (0, \infty)$

65. (a) $(f \circ g)(3) = f(g(3)) = f(-1) = 4$
 (b) $g(f(2)) = g(1) = -2$
 (c) $g(f(5)) = g(-5)$, which is undefined
 (d) $(f \circ g)(-3) = f(g(-3)) = f(-2) = 3$
 (e) $(g \circ f)(-1) = g(f(-1)) = g(4) = 2$
 (f) $f(g(-1)) = f(-4)$, which is undefined

66. $(A \circ r)(t) = A(r(t)) = A(0.6t) = \pi(0.6t)^2 = 0.36\pi t^2$
 $(A \circ r)(t)$ represents the area of the circle at time t .

67. $F(x) = \sqrt{2x - 2}$
 Let $h(x) = 2x$, $g(x) = x - 2$ and $f(x) = \sqrt{x}$.
 Then, $(f \circ g \circ h)(x) = f(g(2x)) = f((2x) - 2) = \sqrt{(2x) - 2} = \sqrt{2x - 2} = F(x)$.
 [Other answers possible]

68. $F(x) = -4 \sin(1 - x)$
 Let $f(x) = -4x$, $g(x) = \sin x$ and $h(x) = 1 - x$. Then,
 $(f \circ g \circ h)(x) = f(g(1 - x)) = f(\sin(1 - x)) = -4 \sin(1 - x) = F(x)$.
 [Other answers possible]

69. $f(-x) = (-x)^2(4 - (-x)^2) = x^2(4 - x^2) = f(x)$
 Even

70. $f(-x) = \sqrt[3]{-x} = -\sqrt[3]{x} = -f(x)$
 Odd

71. $f(-x) = (-x) \cos(-x) = -x \cos x = -f(x)$
 Odd

72. $f(-x) = \sin^2(-x) = \sin(-x)\sin(-x) = (-\sin x)(-\sin x) = \sin^2 x$
 Even

73. (a) If f is even, then $(\frac{3}{2}, 4)$ is on the graph.


(b) If f is odd, then $(\frac{3}{2}, -4)$ is on the graph.

74. (a) If f is even, then $(-4, 9)$ is on the graph.


(b) If f is odd, then $(-4, -9)$ is on the graph.

75. f is even because the graph is symmetric about the y -axis. g is neither even nor odd. h is odd because the graph is symmetric about the origin.

76. (a) If f is even, then the graph is symmetric about the y -axis.



(b) If f is odd, then the graph is symmetric about the origin.



77. Slope = $\frac{4 - (-6)}{-2 - 0} = \frac{10}{-2} = -5$


$$y - 4 = -5(x - (-2))$$

$$y - 4 = -5x - 10$$

$$y = -5x - 6$$

For the line segment, you must restrict the domain.

$$f(x) = -5x - 6, -2 \leq x \leq 0$$



78. Slope = $\frac{8 - 1}{5 - 3} = \frac{7}{2}$


$$y - 1 = \frac{7}{2}(x - 3)$$

$$y - 1 = \frac{7}{2}x - \frac{21}{2}$$

$$y = \frac{7}{2}x - \frac{19}{2}$$

For the line segment, you must restrict the domain.

$$f(x) = \frac{7}{2}x - \frac{19}{2}, 3 \leq x \leq 5$$




79. $x + y^2 = 0$

$$y^2 = -x$$

$$y = -\sqrt{-x}$$


$$f(x) = -\sqrt{-x}, x \leq 0$$




80. $x^2 + y^2 = 36$

$$y^2 = 36 - x^2$$


$$y = -\sqrt{36 - x^2}, -6 \leq x \leq 6$$



81. Answers will vary. *Sample answer:* Speed begins and ends at 0. The speed might be constant in the middle:




82. Answers will vary. *Sample answer:* Height begins a few feet above 0, and ends at 0.



83. Answers will vary. *Sample answer:* In general, as the price decreases, the store will sell more.



84. Answers will vary. *Sample answer:* As time goes on, the value of the car will decrease



85.
$$y = \sqrt{c - x^2}$$

$$y^2 = c - x^2$$

$$x^2 + y^2 = c, \text{ a circle.}$$

For the domain to be $[-5, 5]$, $c = 25$.

86. For the domain to be the set of all real numbers, you must require that $x^2 + 3cx + 6 \neq 0$. So, the discriminant must be less than zero:

$$(3c)^2 - 4(6) < 0$$

$$9c^2 < 24$$

$$c^2 < \frac{8}{3}$$

$$-\sqrt{\frac{8}{3}} < c < \sqrt{\frac{8}{3}}$$

$$-\frac{2}{3}\sqrt{6} < c < \frac{2}{3}\sqrt{6}$$


87. (a) $T(4) = 16^\circ$, $T(15) \approx 23^\circ$

(b) If $H(t) = T(t - 1)$, then the changes in temperature will occur 1 hour later.

(c) If $H(t) = T(t) - 1$, then the overall temperature would be 1 degree lower.


88. (a) For each time t , there corresponds a depth d .
 (b) Domain: $0 \leq t \leq 5$

Range: $0 \leq d \leq 30$




- (d) $d(4) \approx 18$. At time 4 seconds, the depth is approximately 18 cm.

89. (a)



- (b) $A(20) \approx 384$ acres/farm

90. (a)



(b)
$$H\left(\frac{x}{1.6}\right) = 0.002\left(\frac{x}{1.6}\right)^2 + 0.005\left(\frac{x}{1.6}\right) - 0.029$$

$$= 0.00078125x^2 + 0.003125x - 0.029$$

91. $f(x) = |x| + |x - 2|$

If $x < 0$, then $f(x) = -x - (x - 2) = -2x + 2$.

If $0 \leq x < 2$, then $f(x) = x - (x - 2) = 2$.

If $x \geq 2$, then $f(x) = x + (x - 2) = 2x - 2$.

So,

$$f(x) = \begin{cases} -2x + 2, & x \leq 0 \\ 2, & 0 < x < 2. \\ 2x - 2, & x \geq 2 \end{cases}$$

92. $p_1(x) = x^3 - x + 1$ has one zero. $p_2(x) = x^3 - x$ has three zeros. Every cubic polynomial has at least one zero. Given $p(x) = Ax^3 + Bx^2 + Cx + D$, you have $p \rightarrow -\infty$ as $x \rightarrow -\infty$ and $p \rightarrow \infty$ as $x \rightarrow \infty$ if $A > 0$. Furthermore, $p \rightarrow \infty$ as $x \rightarrow -\infty$ and $p \rightarrow -\infty$ as $x \rightarrow \infty$ if $A < 0$. Because the graph has no breaks, the graph must cross the x -axis at least one time.

$$\begin{aligned} 94. f(-x) &= a_{2n}(-x)^{2n} + a_{2n-2}(-x)^{2n-2} + \cdots + a_2(-x)^2 + a_0 \\ &= a_{2n}x^{2n} + a_{2n-2}x^{2n-2} + \cdots + a_2x^2 + a_0 \\ &= f(x) \end{aligned}$$

Even

95. Let $F(x) = f(x)g(x)$ where f and g are even. Then $F(-x) = f(-x)g(-x) = f(x)g(x) = F(x)$.

So, $F(x)$ is even. Let $F(x) = f(x)g(x)$ where f and g are odd. Then

$$F(-x) = f(-x)g(-x) = [-f(x)][-g(x)] = f(x)g(x) = F(x).$$


So, $F(x)$ is even.

96. Let $F(x) = f(x)g(x)$ where f is even and g is odd. Then

$$F(-x) = f(-x)g(-x) = f(x)[-g(x)] = -f(x)g(x) = -F(x).$$

So, $F(x)$ is odd.

97. (a) $V = x(24 - 2x)^2$

Domain: $0 < x < 12$ Maximum volume occurs at $x = 4$. So, the dimensions of the box would be $4 \times 16 \times 16$ cm.

(c)

x	length and width	volume
1	$24 - 2(1)$	$1[24 - 2(1)]^2 = 484$
2	$24 - 2(2)$	$2[24 - 2(2)]^2 = 800$
3	$24 - 2(3)$	$3[24 - 2(3)]^2 = 972$
4	$24 - 2(4)$	$4[24 - 2(4)]^2 = 1024$
5	$24 - 2(5)$	$5[24 - 2(5)]^2 = 980$
6	$24 - 2(6)$	$6[24 - 2(6)]^2 = 864$

The dimensions of the box that yield a maximum volume appear to be $4 \times 16 \times 16$ cm.

$$\begin{aligned} 93. f(-x) &= a_{2n+1}(-x)^{2n+1} + \cdots + a_3(-x)^3 + a_1(-x) \\ &= -[a_{2n+1}x^{2n+1} + \cdots + a_3x^3 + a_1x] \\ &= -f(x) \end{aligned}$$

Odd

98. By equating slopes, $\frac{y-2}{0-3} = \frac{0-2}{x-3}$

$$y - 2 = \frac{6}{x - 3}$$

$$y = \frac{6}{x - 3} + 2 = \frac{2x}{x - 3},$$

$$L = \sqrt{x^2 + y^2} = \sqrt{x^2 + \left(\frac{2x}{x - 3}\right)^2}.$$


99. False. If $f(x) = x^2$, then $f(-3) = f(3) = 9$, but $-3 \neq 3$.

100. True

101. True. The function is even.


102. False. If $f(x) = x^2$ then, $f(3x) = (3x)^2 = 9x^2$ and $3f(x) = 3x^2$. So, $3f(x) \neq f(3x)$.

103. First consider the portion of R in the first quadrant: $x \geq 0, 0 \leq y \leq 1$ and $x - y \leq 1$; shown below.



The area of this region is $1 + \frac{1}{2} = \frac{3}{2}$.

By symmetry, you obtain the entire region R :



The area of R is $4\left(\frac{3}{2}\right) = 6$.

104. Let $g(x) = c$ be constant polynomial.


Then $f(g(x)) = f(c)$ and $g(f(x)) = c$.

So, $f(c) = c$. Because this is true for all real numbers c , f is the identity function: $f(x) = x$.

Section P.4 Fitting Models to Data

1. Trigonometric function
2. Quadratic function
3. No relationship
4. Linear function


5. (a), (b)



Yes. The cancer mortality increases linearly with increased exposure to the carcinogenic substance.

(c) If $x = 3$, then $y \approx 136$.

6. (a)




No, the relationship does not appear to be linear.

(b) Quiz scores are dependent on several variables such as study time, class attendance, etc. These variables may change from one quiz to the next.

7. (a) $d = 0.066F$


(b)



The model fits the data well.

(c) If $F = 55$, then $d \approx 0.066(55) = 3.63$ cm.

8. (a) $s = 9.7t + 0.4$




The model fits the data well.

(c) If $t = 2.5$, $s = 24.65$ meters/second.

9. (a) Using a graphing utility,
 $y = 0.151x + 0.10$


The correlation coefficient is $r \approx 0.880$.



(c) Greater per capita energy consumption by a country tends to correspond to greater per capita gross national product. The four countries that differ most from the linear model are Venezuela, South Korea, Hong Kong and United Kingdom.

(d) Using a graphing utility,
 $y = 0.155x + 0.22$ and $r \approx 0.984$.


10. (a) Linear model: $H = -0.3323t + 612.9333$



The model fits the data well.


(c) When $t = 500$,
 $H = -0.3323(500) + 612.9333 \approx 446.78$.

11. (a) $y_1 = 0.04040t^3 - 0.3695t^2 + 1.123t + 5.88$
 $y_2 = 0.264t + 3.35$
 $y_3 = 0.01439t^3 - 0.1886t^2 + 0.476t + 1.59$




For year 12, $y_1 + y_2 + y_3 \approx 47.5$ cents/mile.

12. (a) $S = 180.89x^2 - 205.79x + 272$



(c) When $x = 2$, $S \approx 583.98$ pounds.

13. (a) $t = 0.002s^2 - 0.04s + 1.9$



(c) According to the model, the times required to attain speeds of less than 20 miles per hour are all about the same.


(d) Adding $(0, 0)$ to the data produces

$$t = 0.002s^2 + 0.02s + 0.1$$

(e) No. From the graph in part (b), you can see that the model from part (a) follows the data more closely than the model from part (d).


14. (a) $N_1 = 3.72t + 31.6$

$$N_3 = -0.0932t^3 + 1.735t^2 - 3.77t + 35.1$$



(c) The cubic model is better.

(d) $N_2 = -0.221t^2 + 6.8t + 24.9$




The model does not fit the data well.

(e) For 2007, $t = 17$, and $N_1 \approx 94.8$ million and $N_3 \approx 14.5$ million. Neither seem accurate. The linear model's estimate is too high and the cubic model's estimate is too low.


(f) Answers will vary

15. (a) $y = -1.806x^3 + 14.58x^2 + 16.4x + 10$



(c) If $x = 4.5$, $y \approx 214$ horsepower.

16. (a) $T = 2.9856 \times 10^{-4} p^3 - 0.0641 p^2 + 5.282 p + 143.1$




- (c) For $T = 300^\circ F$, $p \approx 68.29$ lb/in.².
 (d) The model is based on data up to 100 pounds per square inch.

17. (a) Yes, y is a function of t . At each time t , there is one and only one displacement y .

(b) The amplitude is approximately $(2.35 - 1.65)/2 = 0.35$.


The period is approximately $2(0.375 - 0.125) = 0.5$.

(c) One model is $y = 0.35 \sin(4\pi t) + 2$.




The model appears to fit the data.

18. (a) $S(t) = 56.37 + 25.47 \sin(0.5080t - 2.07)$



The model is a good fit.



The model is a good fit.

(d) The average is the constant term in each model. $83.70^\circ F$ for Miami and $56.37^\circ F$ for Syracuse.


(e) The period for Miami is $2\pi/0.4912 \approx 12.8$. The period for Syracuse is $2\pi/0.5080 \approx 12.4$. In both cases the period is approximately 12, or one year.

(f) Syracuse has greater variability because $25.47 > 7.46$.

19. Answers will vary.

20. Answers will vary.

21. Yes, $A_1 \leq A_2$. To see this, consider the two triangles of areas A_1 and A_2 :



For $i = 1, 2$, the angles satisfy $\alpha_i + \beta_i + \gamma_i = \pi$. At least one of $\alpha_1 \leq \alpha_2$, $\beta_1 \leq \beta_2$, $\gamma_1 \leq \gamma_2$ must hold.

Assume $\alpha_1 \leq \alpha_2$. Because $\alpha_2 \leq \pi/2$ (acute triangle), and the sine function increases on $[0, \pi/2]$, you have

$$\begin{aligned} A_1 &= \frac{1}{2} b_1 c_1 \sin \alpha_1 \leq \frac{1}{2} b_2 c_2 \sin \alpha_1 \\ &\leq \frac{1}{2} b_2 c_2 \sin \alpha_2 = A_2 \end{aligned}$$

Review Exercises for Chapter P

1. $y = 5x - 8$

$x = 0: y = 5(0) - 8 = -8 \Rightarrow (0, -8)$ y-intercept

$y = 0: 0 = 5x - 8 \Rightarrow x = \frac{8}{5} \Rightarrow (\frac{8}{5}, 0)$ x-intercept

2. $y = (x - 2)(x - 6)$

$x = 0: y = (0 - 2)(0 - 6) = 12 \Rightarrow (0, 12)$ y-intercept

$y = 0: 0 = (x - 2)(x - 6) \Rightarrow x = 2, 6 \Rightarrow (2, 0), (6, 0)$ x-intercepts

3. $y = \frac{x-3}{x-4}$

$x = 0: y = \frac{0-3}{0-4} = \frac{3}{4} \Rightarrow \left(0, \frac{3}{4}\right)$ y-intercept

$y = 0: 0 = \frac{x-3}{x-4} \Rightarrow x = 3 \Rightarrow (3, 0)$ x-intercept.

4. $xy = 4$

$x = 0$ and $y = 0$ are both impossible. No intercepts.

5. Symmetric with respect to y-axis because

$(-x)^2 y - (-x)^2 + 4y = 0$

$x^2 y - x^2 + 4y = 0.$

6. Symmetric with respect to y-axis because


$y = (-x)^4 - (-x)^2 + 3$

$y = x^4 - x^2 + 3.$

7. $y = -\frac{1}{2}x + \frac{3}{2}$

Slope: $-\frac{1}{2}$

y-intercept: $\frac{3}{2}$




8. $6x - 3y = 12$

$-3y = -6x + 12$

$y = 2x - 4$

Slope: 2

y-intercept: $(0, -4)$




9. $-\frac{1}{3}x + \frac{5}{6}y = 1$

$-\frac{2}{5}x + y = \frac{6}{5}$

$y = \frac{2}{5}x + \frac{6}{5}$

Slope: $\frac{2}{5}$

y-intercept: $\frac{6}{5}$




10. $0.02x + 0.15y = 0.25$

$2x + 15y = 25$

$y = -\frac{2}{15}x + \frac{5}{3}$

Slope: $-\frac{2}{15}$


y-intercept: $\left(0, \frac{5}{3}\right)$



11. $y = 9 - 8x - x^2 = -(x-1)(x+9)$

y-intercept: $(0, 9)$


x-intercepts: $(1, 0), (-9, 0)$




12. $y = x(6-x)$

y-intercept: $(0, 0)$


x-intercepts: $(0, 0), (6, 0)$



13. $y = 2\sqrt{4-x}$
 Domain: $(-\infty, 4]$



14. $y = |x-4| - 4$




15. $y = 4x^2 - 25$

Xmin = -5
 Xmax = 5
 Xscl = 1
 Ymin = -30
 Ymax = 10
 Yscl = 5

16. $y = 8\sqrt[3]{x-6}$

Xmin = -40
 Xmax = 40
 Xscl = 10
 Ymin = -40
 Ymax = 40
 Yscl = 10

17. $5x + 3y = -1 \Rightarrow y = \frac{1}{3}(-5x - 1)$
 $x - y = -5 \Rightarrow y = x + 5$




Using a graphing utility, the lines intersect at $(-2, 3)$. Analytically,

$$\begin{aligned} \frac{1}{3}(-5x - 1) &= x + 5 \\ -5x - 1 &= 3x + 15 \\ -16 &= 8x \\ -2 &= x. \end{aligned}$$

For $x = -2$, $y = x + 5 = -2 + 5 = 3$.

18. $x - y + 1 = 0 \Rightarrow y = x + 1$
 $y - x^2 = 7 \Rightarrow y = x^2 + 7$



$$\begin{aligned} y &= x + 1 \\ (x + 1) - x^2 &= 7 \\ 0 &= x^2 - x + 6 \end{aligned}$$

No real solution.

No points of intersection.

The graphs of $y = x + 1$ and $y = x^2 + 7$ do not intersect.

19. Answers will vary. *Sample answer:*

You need factors $(x + 4)$ and $(x - 4)$.

Multiply by x to obtain origin symmetry.

$$y = x(x + 4)(x - 4) = x^3 - 16x$$


20. $y = kx^3$

(a) $4 = k(1)^3 \Rightarrow k = 4$ and $y = 4x^3$

(b) $1 = k(-2)^3 \Rightarrow k = -\frac{1}{8}$ and $y = -\frac{1}{8}x^3$


(c) $0 = k(0)^3 \Rightarrow$ and k will do!

(d) $-1 = k(-1)^3 \Rightarrow k = 1 \Rightarrow y = x^3$



$$\text{Slope} = \frac{\left(\frac{5}{2}\right) - 1}{5 - \left(\frac{3}{2}\right)} = \frac{\frac{3}{2}}{\frac{7}{2}} = \frac{3}{7}$$

22. The line is horizontal and has slope 0.



$$23. \frac{t-5}{0-(-8)} = \frac{-1-5}{2-(-8)}$$

$$\frac{t-5}{8} = \frac{-6}{10}$$

$$\frac{t-5}{8} = \frac{3}{5}$$

$$5t - 25 = -24$$

$$5t = 1$$

$$t = \frac{1}{5}$$

$$24. \frac{3-(-1)}{-3-t} = \frac{3-6}{-3-8}$$

$$\frac{4}{-3-t} = \frac{-3}{-11}$$

$$-44 = 9 + 3t$$

$$-53 = 3t$$


$$t = -\frac{53}{3}$$

$$25. y - (-5) = \frac{7}{4}(x - 3)$$

$$y + 5 = \frac{7}{4}x - \frac{21}{4}$$


$$4y + 20 = 7x - 21$$

$$0 = 7x - 4y - 41$$



26. Because m is undefined the line is vertical.


$$x = -8 \text{ or } x + 8 = 0$$



$$27. y - 0 = -\frac{2}{3}(x - (-3))$$

$$y = -\frac{2}{3}x - 2$$


$$2x + 3y + 6 = 0$$



28. Because $m = 0$, the line is horizontal.

$$y - 4 = 0(x - 5)$$

$$y = 4 \text{ or } y - 4 = 0$$



$$29. (a) y - 5 = \frac{7}{16}(x + 3)$$

$$16y - 80 = 7x + 21$$

$$0 = 7x - 16y + 101$$

(b) $5x - 3y = 3$ has slope $\frac{5}{3}$.

$$y - 5 = \frac{5}{3}(x + 3)$$

$$3y - 15 = 5x + 15$$

$$0 = 5x - 3y + 30$$

$$(c) m = \frac{5-0}{-3-0} = -\frac{5}{3}$$

$$y - 5 = -\frac{5}{3}(x + 3)$$

$$3y - 15 = -5x - 15$$

$$5x + 3y = 0$$

(d) Slope is undefined so the line is vertical.

$$x = -3$$

$$x + 3 = 0$$

$$30. (a) y - 4 = -\frac{2}{3}(x - 2)$$

$$3y - 12 = -2x + 4$$

$$2x + 3y - 16 = 0$$

(b) $x + y = 0$ has slope -1 . Slope of the perpendicular line is 1 .

$$y - 4 = 1(x - 2)$$

$$y = x + 2$$

$$0 = x - y + 2$$

$$(c) m = \frac{4-1}{2-6} = -\frac{3}{4}$$

$$y - 4 = -\frac{3}{4}(x - 2)$$

$$4y - 16 = -3x + 6$$

$$3x + 4y - 22 = 0$$

(d) Because the line is horizontal the slope is 0 .

$$y = 4$$

$$y - 4 = 0$$

31. The slope is -850 .

$$V = -850t + 12,500.$$

$$V(3) = -850(3) + 12,500 = \$9950$$

32. (a) $C = 9.25t + 13.50t + 36,500 = 22.75t + 36,500$

(b) $R = 30t$

(c) $30t = 22.75t + 36,500$


$$7.25t = 36,500$$

$$t \approx 5034.48 \text{ hours to break even}$$

33. $x - y^2 = 6$


$$y = \pm\sqrt{x-6}$$

Not a function because there are two values of y for some x .




34. $x^2 - y = 0$

Function of x because there is one value for y for each x .




35. $y = \frac{|x-2|}{x-2}$

y is a function of x because there is one value of y for each x .



36. $x = 9 - y^2$

Not a function of x since there are two values of y for some x .



37. $f(x) = \frac{1}{x}$

(a) $f(0)$ does not exist.

$$\begin{aligned} \text{(b)} \quad \frac{f(1+\Delta x) - f(1)}{\Delta x} &= \frac{\frac{1}{1+\Delta x} - \frac{1}{1}}{\Delta x} = \frac{1-1-\Delta x}{(1+\Delta x)\Delta x} \\ &= \frac{-1}{1+\Delta x}, \Delta x \neq -1, 0 \end{aligned}$$

38. $f(x) = \begin{cases} x^2 + 2, & x < 0 \\ |x - 2|, & x \geq 0 \end{cases}$

(a) $f(-4) = (-4)^2 + 2 = 18$ (because $-4 < 0$)

(b) $f(0) = |0 - 2| = 2$

(c) $f(1) = |1 - 2| = 1$

39. (a) Domain: $36 - x^2 \geq 0 \Rightarrow -6 \leq x \leq 6$ or $[-6, 6]$

Range: $[0, 6]$

(b) Domain: all $x \neq 5$ or $(-\infty, 5) \cup (5, \infty)$

Range: all $y \neq 0$ or $(-\infty, 0) \cup (0, \infty)$

(c) Domain: all x or $(-\infty, \infty)$

Range: all y or $(-\infty, \infty)$


40. $f(x) = 1 - x^2$ and $g(x) = 2x + 1$

(a) $f(x) - g(x) = (1 - x^2) - (2x + 1) = -x^2 - 2x$


(b) $f(x)g(x) = (1 - x^2)(2x + 1) = -2x^3 - x^2 + 2x + 1$

(c) $g(f(x)) = g(1 - x^2) = 2(1 - x^2) + 1 = 3 - 2x^2$


41. (a) $f(x) = x^3 + c, c = -2, 0, 2$




(b) $f(x) = (x - c)^3, c = -2, 0, 2$




(c) $f(x) = (x - 2)^3 + c, c = -2, 0, 2$



(d) $f(x) = cx^3, c = -2, 0, 2$



42. $f(x) = x^3 - 3x^2$




(a) The graph of g is obtained from f by a vertical shift down 1 unit, followed by a reflection in the x -axis:

$$g(x) = -[f(x) - 1] = -x^3 + 3x^2 + 1$$

(b) The graph of g is obtained from f by a vertical shift upwards of 1 and a horizontal shift of 2 to the right.


$$g(x) = f(x - 2) + 1 = (x - 2)^3 - 3(x - 2)^2 + 1$$

43. (a) Odd powers: $f(x) = x, g(x) = x^3, h(x) = x^5$



The graphs of $f, g,$ and h all rise to the right and fall to the left. As the degree increases, the graph rises and falls more steeply. All three graphs pass through the points $(0, 0), (1, 1),$ and $(-1, -1)$ and are symmetric with respect to the origin.

Even powers: $f(x) = x^2, g(x) = x^4, h(x) = x^6$




The graphs of $f, g,$ and h all rise to the left and to the right. As the degree increases, the graph rises more steeply. All three graphs pass through the points $(0, 0), (1, 1),$ and $(-1, 1)$ and are symmetric with respect to the y -axis.

All of the graphs, even and odd, pass through the origin. As the powers increase, the graphs become flatter in the interval $-1 < x < 1$.

(b) $y = x^7$ will look like $h(x) = x^5$, but rise and fall even more steeply. $y = x^8$ will look like $h(x) = x^6$, but rise even more steeply.


44. (a) $f(x) = x^2(x - 6)^2$

The leading coefficient is positive and the degree is even so the graph will rise to the left and to the right.




(b) $g(x) = x^3(x - 6)^2$

The leading coefficient is positive and the degree is odd so the graph will rise to the right and fall to the left.




(c) $h(x) = x^3(x - 6)^3$

The leading coefficient is positive and the degree is even so the graph will rise to the left and to the right.



45. (a)




$$2x + 2y = 24$$

$$y = 12 - x$$

$$A = xy = x(12 - x)$$

(b) Domain: $0 < x < 12$ or $(0, 12)$




(c) Maximum area is $A = 36 \text{ in.}^2$. In general, the maximum area is attained when the rectangle is a square. In this case, $x = 6$.

46. For company (a) the profit rose rapidly for the first year, and then leveled off. For the second company (b), the profit dropped, and then rose again later.

47. (a) 3 (cubic), negative leading coefficient
 (b) 4 (quartic), positive leading coefficient
 (c) 2 (quadratic), negative leading coefficient
 (d) 5, positive leading coefficient


48. (a) $y = -1.204x + 64.2667$



(c) The data point $(27, 44)$ is probably an error. Without this point, the new model is $y = -1.4344x + 66.4387$.

49. (a) Yes, y is a function of t . At each time t , there is one and only one displacement y .
 (b) The amplitude is approximately $(0.25 - (-0.25))/2 = 0.25$. The period is approximately 1.1.

(c) One model is $y = \frac{1}{4} \cos\left(\frac{2\pi}{1.1}t\right) \approx \frac{1}{4} \cos(5.7t)$



The model appears to fit the data.

Problem Solving for Chapter P

1. (a) $x^2 - 6x + y^2 - 8y = 0$
 $(x^2 - 6x + 9) + (y^2 - 8y + 16) = 9 + 16$
 $(x - 3)^2 + (y - 4)^2 = 25$

Center: $(3, 4)$; Radius: 5

(b) Slope of line from $(0, 0)$ to $(3, 4)$ is $\frac{4}{3}$. Slope of tangent line is $-\frac{3}{4}$. So, $y - 0 = -\frac{3}{4}(x - 0) \Rightarrow y = -\frac{3}{4}x$ Tangent line

(c) Slope of line from $(6, 0)$ to $(3, 4)$ is $\frac{4 - 0}{3 - 6} = -\frac{4}{3}$.

Slope of tangent line is $\frac{3}{4}$. So, $y - 0 = \frac{3}{4}(x - 6) \Rightarrow y = \frac{3}{4}x - \frac{9}{2}$ Tangent line

(d) $-\frac{3}{4}x = \frac{3}{4}x - \frac{9}{2}$

$$\frac{3}{2}x = \frac{9}{2}$$

$$x = 3$$

Intersection: $\left(3, -\frac{9}{4}\right)$

2. Let $y = mx + 1$ be a tangent line to the circle from the point $(0, 1)$. Because the center of the circle is at $(0, -1)$ and the radius is 1 you have the following.

$$x^2 + (y + 1)^2 = 1$$

$$x^2 + (mx + 1 + 1)^2 = 1$$

$$(m^2 + 1)x^2 + 4mx + 3 = 0$$

Setting the discriminant $b^2 - 4ac$ equal to zero,

$$16m^2 - 4(m^2 + 1)(3) = 0$$


$$16m^2 - 12m^2 = 12$$

$$4m^2 = 12$$


$$m = \pm\sqrt{3}$$

Tangent lines: $y = \sqrt{3}x + 1$ and $y = -\sqrt{3}x + 1$.


3. $H(x) = \begin{cases} 1, & x \geq 0 \\ 0, & x < 0 \end{cases}$




(d) $H(-x) = \begin{cases} 1, & x \leq 0 \\ 0, & x > 0 \end{cases}$




(a) $H(x) - 2 = \begin{cases} -1, & x \geq 0 \\ -2, & x < 0 \end{cases}$




(e) $\frac{1}{2}H(x) = \begin{cases} \frac{1}{2}, & x \geq 0 \\ 0, & x < 0 \end{cases}$




(b) $H(x - 2) = \begin{cases} 1, & x \geq 2 \\ 0, & x < 2 \end{cases}$




(f) $-H(x - 2) + 2 = \begin{cases} 1, & x \geq 2 \\ 2, & x < 2 \end{cases}$




(c) $-H(x) = \begin{cases} -1, & x \geq 0 \\ 0, & x < 0 \end{cases}$




4. (a) $f(x + 1)$




(b) $f(x) + 1$




(c) $2f(x)$




(d) $f(-x)$




(e) $-f(x)$



(f) $|f(x)|$




(g) $f(|x|)$



5. (a) $x + 2y = 100 \Rightarrow y = \frac{100 - x}{2}$

$$A(x) = xy = x\left(\frac{100 - x}{2}\right) = -\frac{x^2}{2} + 50x$$

Domain: $0 < x < 100$ or $(0, 100)$



Maximum of 1250 m^2 at $x = 50 \text{ m}$, $y = 25 \text{ m}$.

(c) $A(x) = -\frac{1}{2}(x^2 - 100x)$

$$= -\frac{1}{2}(x^2 - 100x + 2500) + 1250$$

$$= -\frac{1}{2}(x - 50)^2 + 1250$$


$A(50) = 1250 \text{ m}^2$ is the maximum.

$x = 50 \text{ m}$, $y = 25 \text{ m}$

6. (a) $4y + 3x = 300 \Rightarrow y = \frac{300 - 3x}{4}$

$$A(x) = x(2y) = x\left(\frac{300 - 3x}{2}\right) = \frac{-3x^2 + 300x}{2}$$

Domain: $0 < x < 100$



Maximum of 3750 ft^2 at $x = 50 \text{ ft}$, $y = 37.5 \text{ ft}$.

(c) $A(x) = -\frac{3}{2}(x^2 - 100x)$

$$= -\frac{3}{2}(x^2 - 100x + 2500) + 3750$$

$$= -\frac{3}{2}(x - 50)^2 + 3750$$

$A(50) = 3750$ square feet is the maximum area,

where $x = 50 \text{ ft}$ and $y = 37.5 \text{ ft}$.

7. The length of the trip in the water is $\sqrt{2^2 + x^2}$, and the length of the trip over land is $\sqrt{1 + (3 - x)^2}$. So, the total time is $T = \frac{\sqrt{4 + x^2}}{2} + \frac{\sqrt{1 + (3 - x)^2}}{4}$ hours.

8. Let d be the distance from the starting point to the beach.

$$\begin{aligned} \text{Average speed} &= \frac{\text{distance}}{\text{time}} \\ &= \frac{2d}{\frac{d}{120} + \frac{d}{60}} \\ &= \frac{2}{\frac{1}{120} + \frac{1}{60}} \\ &= 80 \text{ km/h} \end{aligned}$$


9. (a) Slope = $\frac{9 - 4}{3 - 2} = 5$. Slope of tangent line is less than 5.

(b) Slope = $\frac{4 - 1}{2 - 1} = 3$. Slope of tangent line is greater than 3.

(c) Slope = $\frac{4.41 - 4}{2.1 - 2} = 4.1$. Slope of tangent line is less than 4.1.

$$\begin{aligned} \text{(d) Slope} &= \frac{f(2 + h) - f(2)}{(2 + h) - 2} \\ &= \frac{(2 + h)^2 - 4}{h} \\ &= \frac{4h + h^2}{h} \\ &= 4 + h, h \neq 0 \end{aligned}$$

(e) Letting h get closer and closer to 0, the slope approaches 4. So, the slope at $(2, 4)$ is 4.



(a) Slope = $\frac{3 - 2}{9 - 4} = \frac{1}{5}$. Slope of tangent line is greater than $\frac{1}{5}$.

(b) Slope = $\frac{2 - 1}{4 - 1} = \frac{1}{3}$. Slope of tangent line is less than $\frac{1}{3}$.

(c) Slope = $\frac{2.1 - 2}{4.41 - 4} = \frac{10}{41}$. Slope of tangent line is greater than $\frac{10}{41}$.

$$\text{(d) Slope} = \frac{f(4 + h) - f(4)}{(4 + h) - 4} = \frac{\sqrt{4 + h} - 2}{h}$$

$$\begin{aligned} \text{(e) } \frac{\sqrt{4 + h} - 2}{h} &= \frac{\sqrt{4 + h} - 2}{h} \cdot \frac{\sqrt{4 + h} + 2}{\sqrt{4 + h} + 2} \\ &= \frac{(4 + h) - 4}{h(\sqrt{4 + h} + 2)} \\ &= \frac{1}{\sqrt{4 + h} + 2}, h \neq 0 \end{aligned}$$

As h gets closer to 0, the slope gets closer to $\frac{1}{4}$. The slope is $\frac{1}{4}$ at the point $(4, 2)$.


11. Using the definition of absolute value, you can rewrite the equation.

$$y + |y| = x + |x|$$

$$\begin{cases} 2y, & y > 0 \\ 0, & y \leq 0 \end{cases} = \begin{cases} 2x, & x > 0 \\ 0, & x \leq 0 \end{cases}$$

For $x > 0$ and $y > 0$, you have $2y = 2x \Rightarrow y = x$.

For any $x \leq 0$, y is any $y \leq 0$. So, the graph of $y + |y| = x + |x|$ is as follows.




12. (a) $\frac{I}{x^2} = \frac{2I}{(x-3)^2}$

$$x^2 - 6x + 9 = 2x^2$$

$$x^2 + 6x - 9 = 0$$

$$x = \frac{-6 \pm \sqrt{36 + 36}}{2} = -3 \pm \sqrt{18} \approx 1.2426, -7.2426$$



(b) $\frac{I}{x^2 + y^2} = \frac{2I}{(x-3)^2 + y^2}$


$$(x-3)^2 + y^2 = 2(x^2 + y^2)$$

$$x^2 - 6x + 9 + y^2 = 2x^2 + 2y^2$$

$$x^2 + y^2 + 6x - 9 = 0$$

$$(x+3)^2 + y^2 = 18$$

Circle of radius $\sqrt{18}$ and center $(-3, 0)$.



13. (a) $\frac{I}{x^2 + y^2} = \frac{kI}{(x-4)^2 + y^2}$

$$(x-4)^2 + y^2 = k(x^2 + y^2)$$

$$(k-1)x^2 + 8x + (k-1)y^2 = 16$$


If $k = 1$, then $x = 2$ is a vertical line. Assume $k \neq 1$.

$$x^2 + \frac{8x}{k-1} + y^2 = \frac{16}{k-1}$$

$$x^2 + \frac{8x}{k-1} + \frac{16}{(k-1)^2} + y^2 = \frac{16}{k-1} + \frac{16}{(k-1)^2}$$

$$\left(x + \frac{4}{k-1}\right)^2 + y^2 = \frac{16k}{(k-1)^2}, \text{ Circle}$$

(b) If $k = 3$, $(x+2)^2 + y^2 = 12$



(c) As k becomes very large, $\frac{4}{k-1} \rightarrow 0$ and $\frac{16k}{(k-1)^2} \rightarrow 0$.

The center of the circle gets closer to $(0, 0)$, and its radius approaches 0.

14.

$$d_1 d_2 = 1$$

$$\left[(x+1)^2 + y^2 \right] \left[(x-1)^2 + y^2 \right] = 1$$


$$(x+1)^2(x-1)^2 + y^2 \left[(x+1)^2 + (x-1)^2 \right] + y^4 = 1$$

$$(x^2 - 1)^2 + y^2 [2x^2 + 2] + y^4 = 1$$

$$x^4 - 2x^2 + 1 + 2x^2 y^2 + 2y^2 + y^4 = 1$$

$$(x^4 + 2x^2 y^2 + y^4) - 2x^2 + 2y^2 = 0$$

$$(x^2 + y^2)^2 = 2(x^2 - y^2)$$



Let $y = 0$. Then $x^4 = 2x^2 \Rightarrow x = 0$ or $x^2 = 2$.

So, $(0, 0)$, $(\sqrt{2}, 0)$ and $(-\sqrt{2}, 0)$ are on the curve.

15. $f(x) = y = \frac{1}{1-x}$

(a) Domain: all $x \neq 1$ or $(-\infty, 1) \cup (1, \infty)$

Range: all $y \neq 0$ or $(-\infty, 0) \cup (0, \infty)$


(b) $f(f(x)) = f\left(\frac{1}{1-x}\right) = \frac{1}{1 - \left(\frac{1}{1-x}\right)} = \frac{1}{\frac{1-x-1}{1-x}} = \frac{1-x}{-x} = \frac{x-1}{x}$

Domain: all $x \neq 0, 1$ or $(-\infty, 0) \cup (0, 1) \cup (1, \infty)$

(c) $f(f(f(x))) = f\left(\frac{x-1}{x}\right) = \frac{1}{1 - \left(\frac{x-1}{x}\right)} = \frac{1}{\frac{x-x+1}{x}} = \frac{1}{\frac{1}{x}} = x$

Domain: all $x \neq 0, 1$ or $(-\infty, 0) \cup (0, 1) \cup (1, \infty)$

(d) The graph is not a line. it has holes at $(0, 0)$ and $(1, 1)$.



NOT FOR SALE

CHAPTER 1 Limits and Their Properties

Section 1.1	A Preview of Calculus.....	48
Section 1.2	Finding Limits Graphically and Numerically	49
Section 1.3	Evaluating Limits Analytically	61
Section 1.4	Continuity and One-Sided Limits.....	73
Section 1.5	Infinite Limits	84
Review Exercises	92
Problem Solving	99


INSTRUCTOR USE ONLY

CHAPTER 1 Limits and Their Properties

Section 1.1 A Preview of Calculus

- Precalculus: $(20 \text{ ft/sec})(15 \text{ sec}) = 300 \text{ ft}$
- Calculus required: Velocity is not constant.
Distance $\approx (20 \text{ ft/sec})(15 \text{ sec}) = 300 \text{ ft}$
- Calculus required: Slope of the tangent line at $x = 2$ is the rate of change, and equals about 0.16.
- Precalculus: rate of change = slope = 0.08
- (a) Precalculus: Area = $\frac{1}{2}bh = \frac{1}{2}(5)(4) = 10 \text{ sq. units}$
(b) Calculus required: Area = bh
 $\approx 2(2.5)$
 $= 5 \text{ sq. units}$

6. $f(x) = \sqrt{x}$



(b) slope = $m = \frac{\sqrt{x} - 2}{x - 4}$
 $= \frac{\sqrt{x} - 2}{(\sqrt{x} + 2)(\sqrt{x} - 2)}$
 $= \frac{1}{\sqrt{x} + 2}, x \neq 4$

$x = 1: m = \frac{1}{\sqrt{1} + 2} = \frac{1}{3}$

$x = 3: m = \frac{1}{\sqrt{3} + 2} \approx 0.2679$


$x = 5: m = \frac{1}{\sqrt{5} + 2} \approx 0.2361$

(c) At $P(4, 2)$ the slope is $\frac{1}{\sqrt{4} + 2} = \frac{1}{4} = 0.25$.

You can improve your approximation of the slope at $x = 4$ by considering x -values very close to 4.

8. (a) For the figure on the left, each rectangle has width $\frac{\pi}{4}$.

7. $f(x) = 6x - x^2$



(b) slope = $m = \frac{(6x - x^2) - 8}{x - 2} = \frac{(x - 2)(4 - x)}{x - 2}$
 $= (4 - x), x \neq 2$

For $x = 3, m = 4 - 3 = 1$

For $x = 2.5, m = 4 - 2.5 = 1.5 = \frac{3}{2}$

For $x = 1.5, m = 4 - 1.5 = 2.5 = \frac{5}{2}$

- (c) At $P(2, 8)$, the slope is 2. You can improve your approximation by considering values of x close to 2.

$$\begin{aligned} \text{Area} &\approx \frac{\pi}{4} \left[\sin \frac{\pi}{4} + \sin \frac{\pi}{2} + \sin \frac{3\pi}{4} + \sin \pi \right] \\ &= \frac{\pi}{4} \left[\frac{\sqrt{2}}{2} + 1 + \frac{\sqrt{2}}{2} \right] \\ &= \frac{\sqrt{2} + 1}{4} \pi \approx 1.8961 \end{aligned}$$

For the figure on the right, each rectangle has width $\frac{\pi}{6}$.

$$\begin{aligned} \text{Area} &\approx \frac{\pi}{6} \left[\sin \frac{\pi}{6} + \sin \frac{\pi}{3} + \sin \frac{\pi}{2} + \sin \frac{2\pi}{3} + \sin \frac{5\pi}{6} + \sin \pi \right] \\ &= \frac{\pi}{6} \left[\frac{1}{2} + \frac{\sqrt{3}}{2} + 1 + \frac{\sqrt{3}}{2} + \frac{1}{2} \right] \\ &= \frac{\sqrt{3} + 2}{6} \pi \approx 1.9541 \end{aligned}$$

(b) You could obtain a more accurate approximation by using more rectangles. You will learn later that the exact area is 2.

9. (a) $\text{Area} \approx 5 + \frac{5}{2} + \frac{5}{3} + \frac{5}{4} \approx 10.417$

$$\text{Area} \approx \frac{1}{2} \left(5 + \frac{5}{1.5} + \frac{5}{2} + \frac{5}{2.5} + \frac{5}{3} + \frac{5}{3.5} + \frac{5}{4} + \frac{5}{4.5} \right) \approx 9.145$$

(b) You could improve the approximation by using more rectangles.

10. Answers will vary. Sample *answer*:

The instantaneous rate of change of an automobile's position is the velocity of the automobile, and can be determined by the speedometer.

11. (a) $D_1 = \sqrt{(5-1)^2 + (1-5)^2} = \sqrt{16+16} \approx 5.66$

(b) $D_2 = \sqrt{1 + \left(\frac{5}{2}\right)^2} + \sqrt{1 + \left(\frac{5}{2} - \frac{5}{3}\right)^2} + \sqrt{1 + \left(\frac{5}{3} - \frac{5}{4}\right)^2} + \sqrt{1 + \left(\frac{5}{4} - 1\right)^2}$
 $\approx 2.693 + 1.302 + 1.083 + 1.031 \approx 6.11$

(c) Increase the number of line segments.

Section 1.2 Finding Limits Graphically and Numerically

1.

x	3.9	3.99	3.999	4.001	4.01	4.1
$f(x)$	0.2041	0.2004	0.2000	0.2000	0.1996	0.1961

$$\lim_{x \rightarrow 4} \frac{x-4}{x^2-3x-4} \approx 0.2000 \quad \left(\text{Actual limit is } \frac{1}{5} \right)$$

2.

x	1.9	1.99	1.999	2.001	2.01	2.1
$f(x)$	0.2564	0.2506	0.2501	0.2499	0.2494	0.2439

$$\lim_{x \rightarrow 2} \frac{x-2}{x^2-4} \approx 0.25 \quad \left(\text{Actual limit is } \frac{1}{4} \right)$$

3.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	0.2050	0.2042	0.2041	0.2041	0.2040	0.2033

$$\lim_{x \rightarrow 0} \frac{\sqrt{x+6} - \sqrt{6}}{x} \approx 0.2041 \quad \left(\text{Actual limit is } \frac{1}{2\sqrt{6}} \right)$$

4.

x	-5.1	-5.01	-5.001	-4.999	-4.99	-4.9
$f(x)$	-0.1662	-0.1666	-0.1667	-0.1667	-0.1667	-0.1671

$$\lim_{x \rightarrow -5} \frac{\sqrt{4-x} - 3}{x+5} \approx -0.1667 \quad \left(\text{Actual limit is } -\frac{1}{6} \right)$$

5.

x	2.9	2.99	2.999	3.001	3.01	3.1
$f(x)$	-0.0641	-0.0627	-0.0625	-0.0625	-0.0623	-0.0610

$$\lim_{x \rightarrow 3} \frac{\left[\frac{1}{x+1} \right] - (1/4)}{x-3} \approx -0.0625 \quad \left(\text{Actual limit is } -\frac{1}{16} \right)$$

6.

x	3.9	3.99	3.999	4.001	4.01	4.1
$f(x)$	0.0408	0.0401	0.0400	0.0400	0.0399	0.0392

$$\lim_{x \rightarrow 4} \frac{\left[\frac{x}{x+1} \right] - (4/5)}{x-4} \approx 0.04 \quad \left(\text{Actual limit is } \frac{1}{25} \right)$$

7.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	0.9983	0.99998	1.0000	1.0000	0.99998	0.9983

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} \approx 1.0000 \quad (\text{Actual limit is 1.}) \quad (\text{Make sure you use radian mode.})$$

8.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	0.0500	0.0050	0.0005	-0.0005	-0.0050	-0.0500

$$\lim_{x \rightarrow 0} \frac{\cos x - 1}{x} \approx 0.0000 \quad (\text{Actual limit is 0.}) \quad (\text{Make sure you use radian mode.})$$

9.

x	0.9	0.99	0.999	1.001	1.01	1.1
$f(x)$	0.2564	0.2506	0.2501	0.2499	0.2494	0.2439

$$\lim_{x \rightarrow 1} \frac{x-2}{x^2+x-6} \approx 0.2500 \quad \left(\text{Actual limit is } \frac{1}{4} \right)$$

10.

x	-3.1	-3.01	-3.001	-2.999	-2.99	-2.9
$f(x)$	1.1111	1.0101	1.0010	0.9990	0.9901	0.9091

$$\lim_{x \rightarrow -3} \frac{x+3}{x^2+7x+12} \approx 1.0000 \quad (\text{Actual limit is 1.})$$

11.

x	0.9	0.99	0.999	1.001	1.01	1.1
$f(x)$	0.7340	0.6733	0.6673	0.6660	0.6600	0.6015

$$\lim_{x \rightarrow 1} \frac{x^4 - 1}{x^6 - 1} \approx 0.6666 \quad \left(\text{Actual limit is } \frac{2}{3} \right)$$

12.

x	-2.1	-2.01	-2.001	-1.999	-1.99	-1.9
$f(x)$	12.6100	12.0601	12.0060	11.9940	11.9401	11.4100

$$\lim_{x \rightarrow -2} \frac{x^3 + 8}{x + 2} \approx 12.0000 \quad (\text{Actual limit is } 12.)$$

13.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	1.9867	1.9999	2.0000	2.0000	1.9999	1.9867

$$\lim_{x \rightarrow 0} \frac{\sin 2x}{x} \approx 2.0000 \quad (\text{Actual limit is } 2.) \text{ (Make sure you use radian mode.)}$$

14.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	0.4950	0.5000	0.5000	0.5000	0.5000	0.4950

$$\lim_{x \rightarrow 0} \frac{\tan x}{\tan 2x} \approx 0.5000 \quad \left(\text{Actual limit is } \frac{1}{2} \right)$$

15. $\lim_{x \rightarrow 3} (4 - x) = 1$

16. $\lim_{x \rightarrow 1} (x^2 + 3) = 4$

17. $\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} (4 - x) = 2$

18. $\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} (x^2 + 3) = 4$

19. $\lim_{x \rightarrow 2} \frac{|x - 2|}{x - 2}$ does not exist.

For values of x to the left of 2, $\frac{|x - 2|}{x - 2} = -1$, whereas

for values of x to the right of 2, $\frac{|x - 2|}{x - 2} = 1$.

20. $\lim_{x \rightarrow 5} \frac{2}{x - 5}$ does not exist because the function increases and decreases without bound as x approaches 5.

21. $\lim_{x \rightarrow 1} \sin \pi x = 0$

22. $\lim_{x \rightarrow 0} \sec x = 1$

23. $\lim_{x \rightarrow 0} \cos(1/x)$ does not exist because the function oscillates between -1 and 1 as x approaches 0.

24. $\lim_{x \rightarrow \pi/2} \tan x$ does not exist because the function increases

without bound as x approaches $\frac{\pi}{2}$ from the left and

decreases without bound as x approaches $\frac{\pi}{2}$ from the right.

25. (a) $f(1)$ exists. The black dot at $(1, 2)$ indicates that $f(1) = 2$.

(b) $\lim_{x \rightarrow 1} f(x)$ does not exist. As x approaches 1 from the left, $f(x)$ approaches 3.5, whereas as x approaches 1 from the right, $f(x)$ approaches 1.


(c) $f(4)$ does not exist. The hollow circle at $(4, 2)$ indicates that f is not defined at 4.

(d) $\lim_{x \rightarrow 4} f(x)$ exists. As x approaches 4, $f(x)$ approaches 2: $\lim_{x \rightarrow 4} f(x) = 2$.

26. (a) $f(-2)$ does not exist. The vertical dotted line indicates that f is not defined at -2 .
- (b) $\lim_{x \rightarrow -2} f(x)$ does not exist. As x approaches -2 , the values of $f(x)$ do not approach a specific number.
- (c) $f(0)$ exists. The black dot at $(0, 4)$ indicates that $f(0) = 4$.
- (d) $\lim_{x \rightarrow 0} f(x)$ does not exist. As x approaches 0 from the left, $f(x)$ approaches $\frac{1}{2}$, whereas as x approaches 0 from the right, $f(x)$ approaches 4 .
- (e) $f(2)$ does not exist. The hollow circle at $(2, \frac{1}{2})$ indicates that $f(2)$ is not defined.
- (f) $\lim_{x \rightarrow 2} f(x)$ exists. As x approaches 2 , $f(x)$ approaches $\frac{1}{2}$: $\lim_{x \rightarrow 2} f(x) = \frac{1}{2}$.
- (g) $f(4)$ exists. The black dot at $(4, 2)$ indicates that $f(4) = 2$.
- (h) $\lim_{x \rightarrow 4} f(x)$ does not exist. As x approaches 4 , the values of $f(x)$ do not approach a specific number.


27. $\lim_{x \rightarrow c} f(x)$ exists for all $c \neq -3$.

28. $\lim_{x \rightarrow c} f(x)$ exists for all $c \neq -2, 0$.



$\lim_{x \rightarrow c} f(x)$ exists for all values of $c \neq 4$.


33. $C(t) = 9.99 - 0.79[[-(t - 1)]]$



(b)


t	3	3.3	3.4	3.5	3.6	3.7	4
C	11.57	12.36	12.36	12.36	12.36	12.36	12.36

$\lim_{t \rightarrow 3.5} C(t) = 12.36$




$\lim_{x \rightarrow c} f(x)$ exists for all values of $c \neq \pi$.

31. One possible answer is



32. One possible answer is




(c)

t	2	2.5	2.9	3	3.1	3.5	4
C	10.78	11.57	11.57	11.57	12.36	12.36	12.36

The $\lim_{t \rightarrow 3} C(t)$ does not exist because the values of C approach different values as t approaches 3 from both sides.

34. $C(t) = 5.79 - 0.99[[-(t - 1)]]$



(b)

t	3	3.3	3.4	3.5	3.6	3.7	4
C	7.77	8.76	8.76	8.76	8.76	8.76	8.76

$$\lim_{t \rightarrow 3.5} C(t) = 8.76$$

(c)

t	2	2.5	2.9	3	3.1	3.5	4
C	6.78	7.77	7.77	7.77	8.76	8.76	8.76

The limit $\lim_{t \rightarrow 3} C(t)$ does not exist because the values of C approach different values as t approaches 3 from both sides.

35. You need $|f(x) - 3| = |(x + 1) - 3| = |x - 2| < 0.4$. So, take $\delta = 0.4$. If $0 < |x - 2| < 0.4$, then $|x - 2| = |(x + 1) - 3| = |f(x) - 3| < 0.4$, as desired.

36. You need $|f(x) - 1| = \left| \frac{1}{x-1} - 1 \right| = \left| \frac{2-x}{x-1} \right| < 0.01$. Let $\delta = \frac{1}{101}$. If $0 < |x - 2| < \frac{1}{101}$, then

$$\begin{aligned} -\frac{1}{101} < x - 2 < \frac{1}{101} &\Rightarrow 1 - \frac{1}{101} < x - 1 < 1 + \frac{1}{101} \\ &\Rightarrow \frac{100}{101} < x - 1 < \frac{102}{101} \\ &\Rightarrow |x - 1| > \frac{100}{101} \end{aligned}$$

and you have

$$|f(x) - 1| = \left| \frac{1}{x-1} - 1 \right| = \left| \frac{2-x}{x-1} \right| < \frac{1/101}{100/101} = \frac{1}{100} = 0.01.$$

37. You need to find δ such that $0 < |x - 1| < \delta$ implies

$$|f(x) - 1| = \left| \frac{1}{x} - 1 \right| < 0.1. \text{ That is,}$$

$$-0.1 < \frac{1}{x} - 1 < 0.1$$

$$1 - 0.1 < \frac{1}{x} < 1 + 0.1$$

$$\frac{9}{10} < \frac{1}{x} < \frac{11}{10}$$

$$\frac{10}{9} > x > \frac{10}{11}$$

$$\frac{10}{9} - 1 > x - 1 > \frac{10}{11} - 1$$

$$\frac{1}{9} > x - 1 > -\frac{1}{11}$$

So take $\delta = \frac{1}{11}$. Then $0 < |x - 1| < \delta$ implies

$$-\frac{1}{11} < x - 1 < \frac{1}{11}$$

$$-\frac{1}{11} < x - 1 < \frac{1}{9}$$

Using the first series of equivalent inequalities, you obtain

$$|f(x) - 1| = \left| \frac{1}{x} - 1 \right| < 0.1.$$

38. You need to find δ such that $0 < |x - 2| < \delta$ implies

$$|f(x) - 3| = |x^2 - 1 - 3| = |x^2 - 4| < 0.2. \text{ That is,}$$

$$-0.2 < x^2 - 4 < 0.2$$

$$4 - 0.2 < x^2 < 4 + 0.2$$

$$3.8 < x^2 < 4.2$$

$$\sqrt{3.8} < x < \sqrt{4.2}$$

$$\sqrt{3.8} - 2 < x - 2 < \sqrt{4.2} - 2$$

So take $\delta = \sqrt{4.2} - 2 \approx 0.0494$.

Then $0 < |x - 2| < \delta$ implies

$$-(\sqrt{4.2} - 2) < x - 2 < \sqrt{4.2} - 2$$

$$\sqrt{3.8} - 2 < x - 2 < \sqrt{4.2} - 2.$$

Using the first series of equivalent inequalities, you obtain

$$|f(x) - 3| = |x^2 - 4| < 0.2.$$

39. $\lim_{x \rightarrow 2} (3x + 2) = 8 = L$

$$|(3x + 2) - 8| < 0.01$$

$$|3x - 6| < 0.01$$

$$3|x - 2| < 0.01$$

$$0 < |x - 2| < \frac{0.01}{3} \approx 0.0033 = \delta$$

So, if $0 < |x - 2| < \delta = \frac{0.01}{3}$, you have

$$3|x - 2| < 0.01$$

$$|3x - 6| < 0.01$$

$$|(3x + 2) - 8| < 0.01$$

$$|f(x) - L| < 0.01.$$

40. $\lim_{x \rightarrow 4} \left(4 - \frac{x}{2} \right) = 2 = L$

$$\left| \left(4 - \frac{x}{2} \right) - 2 \right| < 0.01$$

$$\left| 2 - \frac{x}{2} \right| < 0.01$$

$$\left| -\frac{1}{2}(x - 4) \right| < 0.01$$

$$0 < |x - 4| < 0.02 = \delta$$

Hence, if $0 < |x - 4| < \delta = 0.02$, you have

$$\left| -\frac{1}{2}(x - 4) \right| < 0.01$$

$$\left| 2 - \frac{x}{2} \right| < 0.01$$

$$\left| \left(4 - \frac{x}{2} \right) - 2 \right| < 0.01$$

$$|f(x) - L| < 0.01.$$

41. $\lim_{x \rightarrow 2} (x^2 - 3) = 1 = L$

$$|(x^2 - 3) - 1| < 0.01$$

$$|x^2 - 4| < 0.01$$

$$|(x + 2)(x - 2)| < 0.01$$

$$|x + 2||x - 2| < 0.01$$

$$|x - 2| < \frac{0.01}{|x + 2|}$$

If you assume $1 < x < 3$, then $\delta = 0.01/5 = 0.002$.

So, if $0 < |x - 2| < \delta = 0.002$, you have

$$|x - 2| < 0.002 = \frac{1}{5}(0.01) < \frac{1}{|x + 2|}(0.01)$$

$$|x + 2||x - 2| < 0.01$$

$$|x^2 - 4| < 0.01$$

$$|(x^2 - 3) - 1| < 0.01$$

$$|f(x) - L| < 0.01.$$

42. $\lim_{x \rightarrow 5} (x^2 + 4) = 29 = L$

$$|(x^2 + 4) - 29| < 0.01$$

$$|x^2 - 25| < 0.01$$

$$|(x + 5)(x - 5)| < 0.01$$

$$|x - 5| < \frac{0.01}{|x + 5|}$$

If you assume $4 < x < 6$, then $\delta = 0.01/11 \approx 0.0009$.

So, if $0 < |x - 5| < \delta = \frac{0.01}{11}$, you have

$$|x - 5| < \frac{0.01}{11} < \frac{1}{|x + 5|}(0.01)$$

$$|x - 5||x + 5| < 0.01$$

$$|x^2 - 25| < 0.01$$

$$|(x^2 + 4) - 29| < 0.01$$

$$|f(x) - L| < 0.01.$$

43. $\lim_{x \rightarrow 4} (x + 2) = 6$

Given $\varepsilon > 0$:

$$|(x + 2) - 6| < \varepsilon$$

$$|x - 4| < \varepsilon = \delta$$

So, let $\delta = \varepsilon$. So, if $0 < |x - 4| < \delta = \varepsilon$, you have

$$|x - 4| < \varepsilon$$

$$|(x + 2) - 6| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

44. $\lim_{x \rightarrow -3} (2x + 5) = -1$

Given $\varepsilon > 0$:

$$|(2x + 5) - (-1)| < \varepsilon$$

$$|2x + 6| < \varepsilon$$

$$2|x + 3| < \varepsilon$$

$$|x + 3| < \frac{\varepsilon}{2} = \delta$$

So, let $\delta = \varepsilon/2$.

So, if $0 < |x + 3| < \delta = \frac{\varepsilon}{2}$, you have

$$|x + 3| < \frac{\varepsilon}{2}$$

$$|2x + 6| < \varepsilon$$

$$|(2x + 5) - (-1)| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

45. $\lim_{x \rightarrow -4} (\frac{1}{2}x - 1) = \frac{1}{2}(-4) - 1 = -3$

Given $\varepsilon > 0$:

$$|(\frac{1}{2}x - 1) - (-3)| < \varepsilon$$

$$|\frac{1}{2}x + 2| < \varepsilon$$

$$\frac{1}{2}|x - (-4)| < \varepsilon$$

$$|x - (-4)| < 2\varepsilon$$

So, let $\delta = 2\varepsilon$.

So, if $0 < |x - (-4)| < \delta = 2\varepsilon$, you have

$$|x - (-4)| < 2\varepsilon$$

$$|\frac{1}{2}x + 2| < \varepsilon$$

$$|(\frac{1}{2}x - 1) + 3| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

$$46. \lim_{x \rightarrow 1} \left(\frac{2}{5}x + 7 \right) = \frac{2}{5}(1) + 7 = \frac{37}{5}$$

Given $\varepsilon > 0$:

$$\left| \left(\frac{2}{5}x + 7 \right) - \frac{37}{5} \right| = \left| \frac{2}{5}x - \frac{2}{5} \right| < \varepsilon$$

$$\frac{2}{5}|x - 1| < \varepsilon$$

$$|x - 1| < \frac{5}{2}\varepsilon$$

So, let $\delta = \frac{5}{2}\varepsilon$.

So, if $0 < |x - 1| < \delta = \frac{5}{2}\varepsilon$, you have

$$|x - 1| < \frac{5}{2}\varepsilon$$

$$\left| \frac{2}{5}x - \frac{2}{5} \right| < \varepsilon$$

$$\left| \left(\frac{2}{5}x + 7 \right) - \frac{37}{5} \right| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

$$47. \lim_{x \rightarrow 6} 3 = 3$$

Given $\varepsilon > 0$:

$$|3 - 3| < \varepsilon$$

$$0 < \varepsilon$$

So, any $\delta > 0$ will work.

So, for any $\delta > 0$, you have

$$|3 - 3| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

$$48. \lim_{x \rightarrow 2} (-1) = -1$$

$$\text{Given } \varepsilon > 0: |-1 - (-1)| < \varepsilon$$

$$0 < \varepsilon$$

So, any $\delta > 0$ will work.

So, for any $\delta > 0$, you have

$$|(-1) - (-1)| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

$$49. \lim_{x \rightarrow 0} \sqrt[3]{x} = 0$$

$$\text{Given } \varepsilon > 0: \left| \sqrt[3]{x} - 0 \right| < \varepsilon$$

$$\left| \sqrt[3]{x} \right| < \varepsilon$$

$$|x| < \varepsilon^3 = \delta$$

So, let $\delta = \varepsilon^3$.

So, for $0 < |x - 0| < \delta = \varepsilon^3$, you have

$$|x| < \varepsilon^3$$

$$\left| \sqrt[3]{x} \right| < \varepsilon$$

$$\left| \sqrt[3]{x} - 0 \right| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

$$50. \lim_{x \rightarrow 4} \sqrt{x} = \sqrt{4} = 2$$

$$\text{Given } \varepsilon > 0: \left| \sqrt{x} - 2 \right| < \varepsilon$$

$$\left| \sqrt{x} - 2 \right| \left| \sqrt{x} + 2 \right| < \varepsilon \left| \sqrt{x} + 2 \right|$$

$$|x - 4| < \varepsilon \left| \sqrt{x} + 2 \right|$$

Assuming $1 < x < 9$, you can choose $\delta = 3\varepsilon$. Then,

$$0 < |x - 4| < \delta = 3\varepsilon \Rightarrow |x - 4| < \varepsilon \left| \sqrt{x} + 2 \right|$$

$$\Rightarrow \left| \sqrt{x} - 2 \right| < \varepsilon.$$

$$51. \lim_{x \rightarrow -5} |x - 5| = |(-5) - 5| = |-10| = 10$$

$$\text{Given } \varepsilon > 0: \left| |x - 5| - 10 \right| < \varepsilon$$

$$|-(x - 5) - 10| < \varepsilon \quad (x - 5 < 0)$$

$$|-x - 5| < \varepsilon$$

$$|x - (-5)| < \varepsilon$$

So, let $\delta = \varepsilon$.

So for $|x - (-5)| < \delta = \varepsilon$, you have

$$|-(x + 5)| < \varepsilon$$

$$|-(x - 5) - 10| < \varepsilon$$

$$\left| |x - 5| - 10 \right| < \varepsilon \quad (\text{because } x - 5 < 0)$$

$$|f(x) - L| < \varepsilon.$$

52. $\lim_{x \rightarrow 6} |x - 6| = |6 - 6| = 0$

Given $\varepsilon > 0$: $||x - 6| - 0| < \varepsilon$
 $|x - 6| < \varepsilon$

So, let $\delta = \varepsilon$.

So for $|x - 6| < \delta = \varepsilon$, you have

$$|x - 6| < \varepsilon$$

$$||x - 6| - 0| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

53. $\lim_{x \rightarrow 1} (x^2 + 1) = 2$

Given $\varepsilon > 0$:

$$|(x^2 + 1) - 2| < \varepsilon$$

$$|x^2 - 1| < \varepsilon$$

$$|(x + 1)(x - 1)| < \varepsilon$$

$$|x - 1| < \frac{\varepsilon}{|x + 1|}$$

If you assume $0 < x < 2$, then $\delta = \varepsilon/3$.

So for $0 < |x - 1| < \delta = \frac{\varepsilon}{3}$, you have

$$|x - 1| < \frac{1}{3}\varepsilon < \frac{1}{|x + 1|}\varepsilon$$

$$|x^2 - 1| < \varepsilon$$

$$|(x^2 + 1) - 2| < \varepsilon$$

$$|f(x) - 2| < \varepsilon.$$

54. $\lim_{x \rightarrow -3} (x^2 + 3x) = 0$

Given $\varepsilon > 0$:

$$|(x^2 + 3x) - 0| < \varepsilon$$

$$|x(x + 3)| < \varepsilon$$

$$|x + 3| < \frac{\varepsilon}{|x|}$$

If you assume $-4 < x < -2$, then $\delta = \varepsilon/4$.

So for $0 < |x - (-3)| < \delta = \frac{\varepsilon}{4}$, you have

$$|x + 3| < \frac{1}{4}\varepsilon < \frac{1}{|x|}\varepsilon$$

$$|x(x + 3)| < \varepsilon$$

$$|x^2 + 3x - 0| < \varepsilon$$


$$|f(x) - L| < \varepsilon.$$

55. $\lim_{x \rightarrow \pi} f(x) = \lim_{x \rightarrow \pi} 4 = 4$

56. $\lim_{x \rightarrow \pi} f(x) = \lim_{x \rightarrow \pi} x = \pi$

57. $f(x) = \frac{\sqrt{x + 5} - 3}{x - 4}$


$$\lim_{x \rightarrow 4} f(x) = \frac{1}{6}$$



The domain is $[-5, 4) \cup (4, \infty)$. The graphing utility does not show the hole at $(4, \frac{1}{6})$.

58. $f(x) = \frac{x - 3}{x^2 - 4x + 3}$


$$\lim_{x \rightarrow 3} f(x) = \frac{1}{2}$$



The domain is all $x \neq 1, 3$. The graphing utility does not show the hole at $(3, \frac{1}{2})$.


59. $f(x) = \frac{x - 9}{\sqrt{x} - 3}$

$$\lim_{x \rightarrow 9} f(x) = 6$$



The domain is all $x \geq 0$ except $x = 9$. The graphing utility does not show the hole at $(9, 6)$.

60. $f(x) = \frac{x-3}{x^2-9}$
 $\lim_{x \rightarrow 3} f(x) = \frac{1}{6}$




The domain is all $x \neq \pm 3$. The graphing utility does not show the hole at $(3, \frac{1}{6})$.


61. $\lim_{x \rightarrow 8} f(x) = 25$ means that the values of f approach 25 as x gets closer and closer to 8.

62. In the definition of $\lim_{x \rightarrow c} f(x)$, f must be defined on both sides of c , but does not have to be defined at c itself. The value of f at c has no bearing on the limit as x approaches c .


63. (i) The values of f approach different numbers as x approaches c from different sides of c :



(ii) The values of f increase without bound as x approaches c :



(iii) The values of f oscillate between two fixed numbers as x approaches c :



64. (a) No. The fact that $f(2) = 4$ has no bearing on the existence of the limit of $f(x)$ as x approaches 2.

(b) No. The fact that $\lim_{x \rightarrow 2} f(x) = 4$ has no bearing on the value of f at 2.

65. (a) $C = 2\pi r$

$$r = \frac{C}{2\pi} = \frac{6}{2\pi} = \frac{3}{\pi} \approx 0.9549 \text{ cm}$$

(b) When $C = 5.5$: $r = \frac{5.5}{2\pi} \approx 0.87535 \text{ cm}$

When $C = 6.5$: $r = \frac{6.5}{2\pi} \approx 1.03451 \text{ cm}$

So $0.87535 < r < 1.03451$.

(c) $\lim_{x \rightarrow 3/\pi} (2\pi r) = 6$; $\varepsilon = 0.5$; $\delta \approx 0.0796$

66. $V = \frac{4}{3}\pi r^3, V = 2.48$

(a) $2.48 = \frac{4}{3}\pi r^3$

$$r^3 = \frac{1.86}{\pi}$$

$$r \approx 0.8397 \text{ in.}$$

(b) $2.45 \leq V \leq 2.51$

$$2.45 \leq \frac{4}{3}\pi r^3 \leq 2.51$$


$$0.5849 \leq r^3 \leq 0.5992$$

$$0.8363 \leq r \leq 0.8431$$

(c) For $\varepsilon = 2.51 - 2.48 = 0.03$, $\delta \approx 0.003$

67. $f(x) = (1+x)^{1/x}$

$$\lim_{x \rightarrow 0} (1+x)^{1/x} = e \approx 2.71828$$



x	$f(x)$	x	$f(x)$
-0.1	2.867972	0.1	2.593742
-0.01	2.731999	0.01	2.704814
-0.001	2.719642	0.001	2.716942
-0.0001	2.718418	0.0001	2.718146
-0.00001	2.718295	0.00001	2.718268
-0.000001	2.718283	0.000001	2.718280



68. $f(x) = \frac{|x + 1| - |x - 1|}{x}$

x	-1	-0.5	-0.1	0	0.1	0.5	1.0
$f(x)$	2	2	2	Undef.	2	2	2

$\lim_{x \rightarrow 0} f(x) = 2$


Note that for

$-1 < x < 1, x \neq 0, f(x) = \frac{(x + 1) + (x - 1)}{x} = 2.$



Using the zoom and trace feature, $\delta = 0.001$. So $(2 - \delta, 2 + \delta) = (1.999, 2.001)$.

Note: $\frac{x^2 - 4}{x - 2} = x + 2$ for $x \neq 2$.



From the graph, $\delta = 0.001$. So $(3 - \delta, 3 + \delta) = (2.999, 3.001)$.

Note: $\frac{x^2 - 3x}{x - 3} = x$ for $x \neq 3$.

71. False. The existence or nonexistence of $f(x)$ at $x = c$ has no bearing on the existence of the limit of $f(x)$ as $x \rightarrow c$.

79. If $\lim_{x \rightarrow c} f(x) = L_1$ and $\lim_{x \rightarrow c} f(x) = L_2$, then for every $\varepsilon > 0$, there exists $\delta_1 > 0$ and $\delta_2 > 0$ such that $|x - c| < \delta_1 \Rightarrow |f(x) - L_1| < \varepsilon$ and $|x - c| < \delta_2 \Rightarrow |f(x) - L_2| < \varepsilon$. Let δ equal the smaller of δ_1 and δ_2 . Then for $|x - c| < \delta$, you have $|L_1 - L_2| = |L_1 - f(x) + f(x) - L_2| \leq |L_1 - f(x)| + |f(x) - L_2| < \varepsilon + \varepsilon$. Therefore, $|L_1 - L_2| < 2\varepsilon$. Since $\varepsilon > 0$ is arbitrary, it follows that $L_1 = L_2$.

72. True

73. False. Let

$$f(x) = \begin{cases} x - 4, & x \neq 2 \\ 0, & x = 2 \end{cases}$$

$f(2) = 0$

$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} (x - 4) = 2 \neq 0$

74. False. Let

$$f(x) = \begin{cases} x - 4, & x \neq 2 \\ 0, & x = 2 \end{cases}$$

$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} (x - 4) = 2$ and $f(2) = 0 \neq 2$

75. $f(x) = \sqrt{x}$

$\lim_{x \rightarrow 0.25} \sqrt{x} = 0.5$ is true.

As x approaches $0.25 = \frac{1}{4}$ from either side,

$f(x) = \sqrt{x}$ approaches $\frac{1}{2} = 0.5$.

76. $f(x) = \sqrt{x}$

$\lim_{x \rightarrow 0} \sqrt{x} = 0$ is false.

$f(x) = \sqrt{x}$ is not defined on an open interval containing 0 because the domain of f is $x \geq 0$.

77. Using a graphing utility, you see that

$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

$\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = 2$, etc.

So, $\lim_{x \rightarrow 0} \frac{\sin nx}{x} = n$.

78. Using a graphing utility, you see that

$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$

$\lim_{x \rightarrow 0} \frac{\tan 2x}{x} = 2$, etc.

So, $\lim_{x \rightarrow 0} \frac{\tan(nx)}{x} = n$.

- 80.
- $f(x) = mx + b$
- ,
- $m \neq 0$
- . Let
- $\varepsilon > 0$
- be given. Take

$$\delta = \frac{\varepsilon}{|m|}$$

If $0 < |x - c| < \delta = \frac{\varepsilon}{|m|}$, then

$$|m||x - c| < \varepsilon$$

$$|mx - mc| < \varepsilon$$

$$|(mx + b) - (mc + b)| < \varepsilon$$

which shows that $\lim_{x \rightarrow c} (mx + b) = mc + b$.

- 81.
- $\lim_{x \rightarrow c} [f(x) - L] = 0$
- means that for every
- $\varepsilon > 0$
- there exists
- $\delta > 0$
- such that if

$$0 < |x - c| < \delta,$$

then

$$|(f(x) - L) - 0| < \varepsilon.$$

This means the same as $|f(x) - L| < \varepsilon$ when

$$0 < |x - c| < \delta.$$

So, $\lim_{x \rightarrow c} f(x) = L$.

82. (a)
- $(3x + 1)(3x - 1)x^2 + 0.01 = (9x^2 - 1)x^2 + \frac{1}{100}$
-
- $$= 9x^4 - x^2 + \frac{1}{100}$$
-
- $$= \frac{1}{100}(10x^2 - 1)(90x^2 - 1)$$

So, $(3x + 1)(3x - 1)x^2 + 0.01 > 0$ if

$$10x^2 - 1 < 0 \text{ and } 90x^2 - 1 < 0.$$

$$\text{Let } (a, b) = \left(-\frac{1}{\sqrt{90}}, \frac{1}{\sqrt{90}}\right).$$

For all $x \neq 0$ in (a, b) , the graph is positive. You can verify this with a graphing utility.

- (b) You are given
- $\lim_{x \rightarrow c} g(x) = L > 0$
- . Let

$\varepsilon = \frac{1}{2}L$. There exists $\delta > 0$ such that

$$0 < |x - c| < \delta \text{ implies that}$$

$$|g(x) - L| < \varepsilon = \frac{L}{2}. \text{ That is,}$$

$$-\frac{L}{2} < g(x) - L < \frac{L}{2}$$

$$\frac{L}{2} < g(x) < \frac{3L}{2}$$

For x in the interval $(c - \delta, c + \delta)$, $x \neq c$, you

have $g(x) > \frac{L}{2} > 0$, as desired.

83. Answers will vary.

84. $\lim_{x \rightarrow 4} \frac{x^2 - x - 12}{x - 4} = 7$

n	$4 + [0.1]^n$	$f(4 + [0.1]^n)$
1	4.1	7.1
2	4.01	7.01
3	4.001	7.001
4	4.0001	7.0001

n	$4 - [0.1]^n$	$f(4 - [0.1]^n)$
1	3.9	6.9
2	3.99	6.99
3	3.999	6.999
4	3.9999	6.9999

85. The radius
- OP
- has a length equal to the altitude
- z
- of the triangle plus
- $\frac{h}{2}$
- . So,
- $z = 1 - \frac{h}{2}$
- .

$$\text{Area triangle} = \frac{1}{2}b\left(1 - \frac{h}{2}\right)$$


$$\text{Area rectangle} = bh$$

$$\text{Because these are equal, } \frac{1}{2}b\left(1 - \frac{h}{2}\right) = bh$$

$$1 - \frac{h}{2} = 2h$$

$$\frac{5}{2}h = 1$$

$$h = \frac{2}{5}.$$



86. Consider a cross section of the cone, where EF is a diagonal of the inscribed cube. $AD = 3$, $BC = 2$. Let x be the length of a side of the cube. Then $EF = x\sqrt{2}$.

By similar triangles,

$$\frac{EF}{BC} = \frac{AG}{AD}$$


$$\frac{x\sqrt{2}}{2} = \frac{3-x}{3}$$

Solving for x ,


$$3\sqrt{2}x = 6 - 2x$$

$$(3\sqrt{2} + 2)x = 6$$


$$x = \frac{6}{3\sqrt{2} + 2} = \frac{9\sqrt{2} - 6}{7} \approx 0.96.$$




Section 1.3 Evaluating Limits Analytically




- (a) $\lim_{x \rightarrow 4} h(x) = 0$
- (b) $\lim_{x \rightarrow -1} h(x) = -5$



- $$g(x) = \frac{12(\sqrt{x} - 3)}{x - 9}$$
- (a) $\lim_{x \rightarrow 4} g(x) = 2.4$
 - (b) $\lim_{x \rightarrow 0} g(x) = 4$



- $$f(x) = x \cos x$$
- (a) $\lim_{x \rightarrow 0} f(x) = 0$
 - (b) $\lim_{x \rightarrow \pi/3} f(x) \approx 0.524$
 $\left(= \frac{\pi}{6} \right)$



- $$f(t) = t|t - 4|$$
- (a) $\lim_{t \rightarrow 4} f(t) = 0$
 - (b) $\lim_{t \rightarrow -1} f(t) = -5$

- 5. $\lim_{x \rightarrow 2} x^3 = 2^3 = 8$
- 6. $\lim_{x \rightarrow -2} x^4 = (-2)^4 = 16$
- 7. $\lim_{x \rightarrow 0} (2x - 1) = 2(0) - 1 = -1$
- 8. $\lim_{x \rightarrow -3} (3x + 2) = 3(-3) + 2 = -7$
- 9. $\lim_{x \rightarrow -3} (x^2 + 3x) = (-3)^2 + 3(-3) = 9 - 9 = 0$
- 10. $\lim_{x \rightarrow 1} (-x^2 + 1) = -(1)^2 + 1 = 0$
- 11. $\lim_{x \rightarrow -3} (2x^2 + 4x + 1) = 2(-3)^2 + 4(-3) + 1 = 18 - 12 + 1 = 7$
- 12. $\lim_{x \rightarrow 1} (3x^3 - 2x^2 + 4) = 3(1)^3 - 2(1)^2 + 4 = 5$
- 13. $\lim_{x \rightarrow 3} \sqrt{x + 1} = \sqrt{3 + 1} = 2$
- 14. $\lim_{x \rightarrow 4} \sqrt[3]{x + 4} = \sqrt[3]{4 + 4} = 2$
- 15. $\lim_{x \rightarrow -4} (x + 3)^2 = (-4 + 3)^2 = 1$

16. $\lim_{x \rightarrow 0} (2x - 1)^3 = [2(0) - 1]^3 = -1$

17. $\lim_{x \rightarrow 2} \frac{1}{x} = \frac{1}{2}$

18. $\lim_{x \rightarrow -3} \frac{2}{x + 2} = \frac{2}{-3 + 2} = -2$

19. $\lim_{x \rightarrow 1} \frac{x}{x^2 + 4} = \frac{1}{1^2 + 4} = \frac{1}{5}$

20. $\lim_{x \rightarrow 1} \frac{2x - 3}{x + 5} = \frac{2(1) - 3}{1 + 5} = \frac{-1}{6}$

21. $\lim_{x \rightarrow 7} \frac{3x}{\sqrt{x + 2}} = \frac{3(7)}{\sqrt{7 + 2}} = \frac{21}{3} = 7$

22. $\lim_{x \rightarrow 2} \frac{\sqrt{x + 2}}{x - 4} = \frac{\sqrt{2 + 2}}{2 - 4} = \frac{2}{-2} = -1$

23. (a) $\lim_{x \rightarrow 1} f(x) = 5 - 1 = 4$

(b) $\lim_{x \rightarrow 4} g(x) = 4^3 = 64$

(c) $\lim_{x \rightarrow 1} g(f(x)) = g(f(1)) = g(4) = 64$

24. (a) $\lim_{x \rightarrow -3} f(x) = (-3) + 7 = 4$

(b) $\lim_{x \rightarrow 4} g(x) = 4^2 = 16$

(c) $\lim_{x \rightarrow -3} g(f(x)) = g(4) = 16$

25. (a) $\lim_{x \rightarrow 1} f(x) = 4 - 1 = 3$

(b) $\lim_{x \rightarrow 3} g(x) = \sqrt{3 + 1} = 2$

(c) $\lim_{x \rightarrow 1} g(f(x)) = g(3) = 2$

37. (a) $\lim_{x \rightarrow c} [5g(x)] = 5 \lim_{x \rightarrow c} g(x) = 5(2) = 10$

(b) $\lim_{x \rightarrow c} [f(x) + g(x)] = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = 3 + 2 = 5$

(c) $\lim_{x \rightarrow c} [f(x)g(x)] = \left[\lim_{x \rightarrow c} f(x) \right] \left[\lim_{x \rightarrow c} g(x) \right] = (3)(2) = 6$

(d) $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)} = \frac{3}{2}$

38. (a) $\lim_{x \rightarrow c} [4f(x)] = 4 \lim_{x \rightarrow c} f(x) = 4\left(\frac{3}{2}\right) = 6$

(b) $\lim_{x \rightarrow c} [f(x) + g(x)] = \lim_{x \rightarrow c} f(x) + \lim_{x \rightarrow c} g(x) = \frac{3}{2} + \frac{1}{2} = 2$

(c) $\lim_{x \rightarrow c} [f(x)g(x)] = \left[\lim_{x \rightarrow c} f(x) \right] \left[\lim_{x \rightarrow c} g(x) \right] = \left(\frac{3}{2}\right)\left(\frac{1}{2}\right) = \frac{3}{4}$

(d) $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)} = \frac{3/2}{1/2} = 3$

26. (a) $\lim_{x \rightarrow 4} f(x) = 2(4^2) - 3(4) + 1 = 21$

(b) $\lim_{x \rightarrow 21} g(x) = \sqrt[3]{21 + 6} = 3$

(c) $\lim_{x \rightarrow 4} g(f(x)) = g(21) = 3$

27. $\lim_{x \rightarrow \pi/2} \sin x = \sin \frac{\pi}{2} = 1$

28. $\lim_{x \rightarrow \pi} \tan x = \tan \pi = 0$

29. $\lim_{x \rightarrow 1} \cos \frac{\pi x}{3} = \cos \frac{\pi}{3} = \frac{1}{2}$

30. $\lim_{x \rightarrow 2} \sin \frac{\pi x}{2} = \sin \frac{\pi(2)}{2} = 0$

31. $\lim_{x \rightarrow 0} \sec 2x = \sec 0 = 1$

32. $\lim_{x \rightarrow \pi} \cos 3x = \cos 3\pi = -1$

33. $\lim_{x \rightarrow 5\pi/6} \sin x = \sin \frac{5\pi}{6} = \frac{1}{2}$

34. $\lim_{x \rightarrow 5\pi/3} \cos x = \cos \frac{5\pi}{3} = \frac{1}{2}$

35. $\lim_{x \rightarrow 3} \tan\left(\frac{\pi x}{4}\right) = \tan \frac{3\pi}{4} = -1$

36. $\lim_{x \rightarrow 7} \sec\left(\frac{\pi x}{6}\right) = \sec \frac{7\pi}{6} = \frac{-2\sqrt{3}}{3}$

39. (a) $\lim_{x \rightarrow c} [f(x)]^3 = \left[\lim_{x \rightarrow c} f(x) \right]^3 = (4)^3 = 64$

(b) $\lim_{x \rightarrow c} \sqrt{f(x)} = \sqrt{\lim_{x \rightarrow c} f(x)} = \sqrt{4} = 2$

(c) $\lim_{x \rightarrow c} [3f(x)] = 3 \lim_{x \rightarrow c} f(x) = 3(4) = 12$

(d) $\lim_{x \rightarrow c} [f(x)]^{3/2} = \left[\lim_{x \rightarrow c} f(x) \right]^{3/2} = (4)^{3/2} = 8$

40. (a) $\lim_{x \rightarrow c} \sqrt[3]{f(x)} = \sqrt[3]{\lim_{x \rightarrow c} f(x)} = \sqrt[3]{27} = 3$

(b) $\lim_{x \rightarrow c} \frac{f(x)}{18} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} 18} = \frac{27}{18} = \frac{3}{2}$

(c) $\lim_{x \rightarrow c} [f(x)]^2 = \left[\lim_{x \rightarrow c} f(x) \right]^2 = (27)^2 = 729$

(d) $\lim_{x \rightarrow c} [f(x)]^{2/3} = \left[\lim_{x \rightarrow c} f(x) \right]^{2/3} = (27)^{2/3} = 9$

41. $f(x) = x - 1$ and $g(x) = \frac{x^2 - x}{x}$ agree except at $x = 0$.

(a) $\lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} f(x) = 0 - 1 = -1$

(b) $\lim_{x \rightarrow -1} g(x) = \lim_{x \rightarrow -1} f(x) = -1 - 1 = -2$

42. $f(x) = -x + 3$ and $h(x) = \frac{-x^2 + 3x}{x}$ agree except at $x = 0$.

(a) $\lim_{x \rightarrow 2} h(x) = \lim_{x \rightarrow 2} f(x) = -2 + 3 = 1$

(b) $\lim_{x \rightarrow 0} h(x) = \lim_{x \rightarrow 0} f(x) = -0 + 3 = 3$

43. $f(x) = x(x + 1)$ and $g(x) = \frac{x^3 - x}{x - 1}$ agree except at $x = 1$.

(a) $\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} f(x) = 2$

(b) $\lim_{x \rightarrow -1} g(x) = \lim_{x \rightarrow -1} f(x) = 0$


44. $g(x) = \frac{1}{x - 1}$ and $f(x) = \frac{x}{x^2 - x}$ agree except at $x = 0$.

(a) $\lim_{x \rightarrow 1} f(x)$ does not exist.

(b) $\lim_{x \rightarrow 0} f(x) = -1$


45. $f(x) = \frac{x^2 - 1}{x + 1}$ and $g(x) = x - 1$ agree except at $x = -1$.

$\lim_{x \rightarrow -1} f(x) = \lim_{x \rightarrow -1} g(x) = -2$




46. $f(x) = \frac{2x^2 - x - 3}{x + 1}$ and $g(x) = 2x - 3$ agree except at $x = -1$.

$\lim_{x \rightarrow -1} f(x) = \lim_{x \rightarrow -1} g(x) = -5$




47. $f(x) = \frac{x^3 - 8}{x - 2}$ and $g(x) = x^2 + 2x + 4$ agree except at $x = 2$.

$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} g(x) = 12$



48. $f(x) = \frac{x^3 + 1}{x + 1}$ and $g(x) = x^2 - x + 1$ agree except at $x = -1$.

$\lim_{x \rightarrow -1} f(x) = \lim_{x \rightarrow -1} g(x) = 3$



49. $\lim_{x \rightarrow 0} \frac{x}{x^2 - x} = \lim_{x \rightarrow 0} \frac{x}{x(x - 1)} = \lim_{x \rightarrow 0} \frac{1}{x - 1} = -1$

50. $\lim_{x \rightarrow 0} \frac{3x}{x^2 + 2x} = \lim_{x \rightarrow 0} \frac{3x}{x(x + 2)} = \lim_{x \rightarrow 0} \frac{3}{x + 2} = \frac{3}{2}$

$$\begin{aligned} 51. \lim_{x \rightarrow 4} \frac{x-4}{x^2-16} &= \lim_{x \rightarrow 4} \frac{x-4}{(x+4)(x-4)} \\ &= \lim_{x \rightarrow 4} \frac{1}{x+4} = \frac{1}{8} \end{aligned}$$

$$\begin{aligned} 53. \lim_{x \rightarrow -3} \frac{x^2+x-6}{x^2-9} &= \lim_{x \rightarrow -3} \frac{(x+3)(x-2)}{(x+3)(x-3)} \\ &= \lim_{x \rightarrow -3} \frac{x-2}{x-3} = \frac{-5}{-6} = \frac{5}{6} \end{aligned}$$

$$52. \lim_{x \rightarrow 3} \frac{3-x}{x^2-9} = \lim_{x \rightarrow 3} \frac{3-x}{(x-3)(x+3)} = \lim_{x \rightarrow 3} \frac{-1}{x+3} = -\frac{1}{6}$$

$$\begin{aligned} 54. \lim_{x \rightarrow 4} \frac{x^2-5x+4}{x^2-2x-8} &= \lim_{x \rightarrow 4} \frac{(x-4)(x-1)}{(x-4)(x+2)} \\ &= \lim_{x \rightarrow 4} \frac{(x-1)}{(x+2)} = \frac{3}{6} = \frac{1}{2} \end{aligned}$$

$$\begin{aligned} 55. \lim_{x \rightarrow 4} \frac{\sqrt{x+5}-3}{x-4} &= \lim_{x \rightarrow 4} \frac{\sqrt{x+5}-3}{x-4} \cdot \frac{\sqrt{x+5}+3}{\sqrt{x+5}+3} \\ &= \lim_{x \rightarrow 4} \frac{(x+5)-9}{(x-4)(\sqrt{x+5}+3)} = \lim_{x \rightarrow 4} \frac{1}{\sqrt{x+5}+3} = \frac{1}{\sqrt{9}+3} = \frac{1}{6} \end{aligned}$$

$$56. \lim_{x \rightarrow 3} \frac{\sqrt{x+1}-2}{x-3} = \lim_{x \rightarrow 3} \frac{\sqrt{x+1}-2}{x-3} \cdot \frac{\sqrt{x+1}+2}{\sqrt{x+1}+2} = \lim_{x \rightarrow 3} \frac{x-3}{(x-3)[\sqrt{x+1}+2]} = \lim_{x \rightarrow 3} \frac{1}{\sqrt{x+1}+2} = \frac{1}{4}$$

$$\begin{aligned} 57. \lim_{x \rightarrow 0} \frac{\sqrt{x+5}-\sqrt{5}}{x} &= \lim_{x \rightarrow 0} \frac{\sqrt{x+5}-\sqrt{5}}{x} \cdot \frac{\sqrt{x+5}+\sqrt{5}}{\sqrt{x+5}+\sqrt{5}} \\ &= \lim_{x \rightarrow 0} \frac{(x+5)-5}{x(\sqrt{x+5}+\sqrt{5})} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{x+5}+\sqrt{5}} = \frac{1}{2\sqrt{5}} = \frac{\sqrt{5}}{10} \end{aligned}$$

$$\begin{aligned} 58. \lim_{x \rightarrow 0} \frac{\sqrt{2+x}-\sqrt{2}}{x} &= \lim_{x \rightarrow 0} \frac{\sqrt{2+x}-\sqrt{2}}{x} \cdot \frac{\sqrt{2+x}+\sqrt{2}}{\sqrt{2+x}+\sqrt{2}} \\ &= \lim_{x \rightarrow 0} \frac{2+x-2}{(\sqrt{2+x}+\sqrt{2})x} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{2+x}+\sqrt{2}} = \frac{1}{2\sqrt{2}} = \frac{\sqrt{2}}{4} \end{aligned}$$

$$59. \lim_{x \rightarrow 0} \frac{\frac{1}{3+x} - \frac{1}{3}}{x} = \lim_{x \rightarrow 0} \frac{3-(3+x)}{(3+x)3(x)} = \lim_{x \rightarrow 0} \frac{-x}{(3+x)3(x)} = \lim_{x \rightarrow 0} \frac{-1}{(3+x)3} = -\frac{1}{9}$$

$$\begin{aligned} 60. \lim_{x \rightarrow 0} \frac{\frac{1}{x+4} - \frac{1}{4}}{x} &= \lim_{x \rightarrow 0} \frac{\frac{4-(x+4)}{4(x+4)}}{x} \\ &= \lim_{x \rightarrow 0} \frac{-1}{4(x+4)} = -\frac{1}{16} \end{aligned}$$

$$\begin{aligned} 61. \lim_{\Delta x \rightarrow 0} \frac{2(x+\Delta x)-2x}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{2x+2\Delta x-2x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} 2 = 2 \end{aligned}$$

$$62. \lim_{\Delta x \rightarrow 0} \frac{(x+\Delta x)^2 - x^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x\Delta x + (\Delta x)^2 - x^2}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\Delta x(2x + \Delta x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} (2x + \Delta x) = 2x$$

$$\begin{aligned} 63. \lim_{\Delta x \rightarrow 0} \frac{(x+\Delta x)^2 - 2(x+\Delta x) + 1 - (x^2 - 2x + 1)}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x\Delta x + (\Delta x)^2 - 2x - 2\Delta x + 1 - x^2 + 2x - 1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} (2x + \Delta x - 2) = 2x - 2 \end{aligned}$$

$$64. \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^3 - x^3}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{x^3 + 3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3 - x^3}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(3x^2 + 3x\Delta x + (\Delta x)^2)}{\Delta x} = \lim_{\Delta x \rightarrow 0} (3x^2 + 3x\Delta x + (\Delta x)^2) = 3x^2$$

$$65. \lim_{x \rightarrow 0} \frac{\sin x}{5x} = \lim_{x \rightarrow 0} \left[\left(\frac{\sin x}{x} \right) \left(\frac{1}{5} \right) \right] = (1) \left(\frac{1}{5} \right) = \frac{1}{5}$$

$$66. \lim_{x \rightarrow 0} \frac{3(1 - \cos x)}{x} = \lim_{x \rightarrow 0} \left[3 \left(\frac{1 - \cos x}{x} \right) \right] = (3)(0) = 0$$

$$67. \lim_{x \rightarrow 0} \frac{\sin x(1 - \cos x)}{x^2} = \lim_{x \rightarrow 0} \left[\frac{\sin x}{x} \cdot \frac{1 - \cos x}{x} \right]$$

$$= (1)(0) = 0$$

$$68. \lim_{\theta \rightarrow 0} \frac{\cos \theta \tan \theta}{\theta} = \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

$$69. \lim_{x \rightarrow 0} \frac{\sin^2 x}{x} = \lim_{x \rightarrow 0} \left[\frac{\sin x}{x} \sin x \right] = (1) \sin 0 = 0$$

$$70. \lim_{x \rightarrow 0} \frac{\tan^2 x}{x} = \lim_{x \rightarrow 0} \frac{\sin^2 x}{x \cos^2 x} = \lim_{x \rightarrow 0} \left[\frac{\sin x}{x} \cdot \frac{\sin x}{\cos^2 x} \right]$$

$$= (1)(0) = 0$$


$$71. \lim_{h \rightarrow 0} \frac{(1 - \cos h)^2}{h} = \lim_{h \rightarrow 0} \left[\frac{1 - \cos h}{h} (1 - \cos h) \right]$$

$$= (0)(0) = 0$$

$$77. f(x) = \frac{\sqrt{x+2} - \sqrt{2}}{x}$$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.358	0.354	0.354	?	0.354	0.353	0.349

It appears that the limit is 0.354.



The graph has a hole at $x = 0$.

$$\text{Analytically, } \lim_{x \rightarrow 0} \frac{\sqrt{x+2} - \sqrt{2}}{x} = \lim_{x \rightarrow 0} \frac{\sqrt{x+2} - \sqrt{2}}{x} \cdot \frac{\sqrt{x+2} + \sqrt{2}}{\sqrt{x+2} + \sqrt{2}}$$

$$= \lim_{x \rightarrow 0} \frac{x+2-2}{x(\sqrt{x+2} + \sqrt{2})} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{x+2} + \sqrt{2}} = \frac{1}{2\sqrt{2}} = \frac{\sqrt{2}}{4} \approx 0.354.$$

$$72. \lim_{\phi \rightarrow \pi} \phi \sec \phi = \pi(-1) = -\pi$$

$$73. \lim_{x \rightarrow \pi/2} \frac{\cos x}{\cot x} = \lim_{x \rightarrow \pi/2} \sin x = 1$$

$$74. \lim_{x \rightarrow \pi/4} \frac{1 - \tan x}{\sin x - \cos x} = \lim_{x \rightarrow \pi/4} \frac{\cos x - \sin x}{\sin x \cos x - \cos^2 x}$$

$$= \lim_{x \rightarrow \pi/4} \frac{-(\sin x - \cos x)}{\cos x(\sin x - \cos x)}$$

$$= \lim_{x \rightarrow \pi/4} \frac{-1}{\cos x}$$

$$= \lim_{x \rightarrow \pi/4} (-\sec x)$$

$$= -\sqrt{2}$$

$$75. \lim_{t \rightarrow 0} \frac{\sin 3t}{2t} = \lim_{t \rightarrow 0} \left(\frac{\sin 3t}{3t} \right) \left(\frac{3}{2} \right) = (1) \left(\frac{3}{2} \right) = \frac{3}{2}$$


$$76. \lim_{x \rightarrow 0} \frac{\sin 2x}{\sin 3x} = \lim_{x \rightarrow 0} \left[2 \left(\frac{\sin 2x}{2x} \right) \left(\frac{1}{3} \right) \left(\frac{3x}{\sin 3x} \right) \right]$$

$$= 2(1) \left(\frac{1}{3} \right) (1) = \frac{2}{3}$$

78. $f(x) = \frac{4 - \sqrt{x}}{x - 16}$

x	15.9	15.99	15.999	16	16.001	16.01	16.1
$f(x)$	-0.1252	-0.125	-0.125	?	-0.125	-0.125	-0.1248

It appears that the limit is -0.125.




The graph has a hole at $x = 16$.

Analytically, $\lim_{x \rightarrow 16} \frac{4 - \sqrt{x}}{x - 16} = \lim_{x \rightarrow 16} \frac{(4 - \sqrt{x})}{(\sqrt{x} + 4)(\sqrt{x} - 4)} = \lim_{x \rightarrow 16} \frac{-1}{\sqrt{x} + 4} = -\frac{1}{8}$.

79. $f(x) = \frac{\frac{1}{2+x} - \frac{1}{2}}{x}$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	-0.263	-0.251	-0.250	?	-0.250	-0.249	-0.238

It appears that the limit is -0.250.




The graph has a hole at $x = 0$.

Analytically, $\lim_{x \rightarrow 0} \frac{\frac{1}{2+x} - \frac{1}{2}}{x} = \lim_{x \rightarrow 0} \frac{2 - (2+x)}{2(2+x)} \cdot \frac{1}{x} = \lim_{x \rightarrow 0} \frac{-x}{2(2+x)} \cdot \frac{1}{x} = \lim_{x \rightarrow 0} \frac{-1}{2(2+x)} = -\frac{1}{4}$.

80. $f(x) = \frac{x^5 - 32}{x - 2}$

x	1.9	1.99	1.999	1.9999	2.0	2.0001	2.001	2.01	2.1
$f(x)$	72.39	79.20	79.92	79.99	?	80.01	80.08	80.80	88.41

It appears that the limit is 80.



The graph has a hole at $x = 2$.


Analytically, $\lim_{x \rightarrow 2} \frac{x^5 - 32}{x - 2} = \lim_{x \rightarrow 2} \frac{(x - 2)(x^4 + 2x^3 + 4x^2 + 8x + 16)}{x - 2} = \lim_{x \rightarrow 2} (x^4 + 2x^3 + 4x^2 + 8x + 16) = 80$.

(Hint: Use long division to factor $x^5 - 32$.)

81. $f(t) = \frac{\sin 3t}{t}$

t	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(t)$	2.96	2.9996	3	?	3	2.9996	2.96

It appears that the limit is 3.




The graph has a hole at $t = 0$.

Analytically, $\lim_{t \rightarrow 0} \frac{\sin 3t}{t} = \lim_{t \rightarrow 0} 3 \left(\frac{\sin 3t}{3t} \right) = 3(1) = 3$.

82. $f(x) = \frac{\cos x - 1}{2x^2}$

x	-1	-0.1	-0.01	0.01	0.1	1
$f(x)$	-0.2298	-0.2498	-0.25	-0.25	-0.2498	-0.2298

It appears that the limit is -0.25.



The graph has a hole at $x = 0$.


$$\begin{aligned} \text{Analytically, } \frac{\cos x - 1}{2x^2} \cdot \frac{\cos x + 1}{\cos x + 1} &= \frac{\cos^2 x - 1}{2x^2(\cos x + 1)} \\ &= \frac{-\sin^2 x}{2x^2(\cos x + 1)} \\ &= \frac{\sin^2 x}{x^2} \cdot \frac{-1}{2(\cos x + 1)} \end{aligned}$$

$$\lim_{x \rightarrow 0} \left[\frac{\sin^2 x}{x^2} \cdot \frac{-1}{2(\cos x + 1)} \right] = 1 \left(\frac{-1}{4} \right) = -\frac{1}{4} = -0.25$$

83. $f(x) = \frac{\sin x^2}{x}$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	-0.099998	-0.01	-0.001	?	0.001	0.01	0.099998

It appears that the limit is 0.




The graph has a hole at $x = 0$.

Analytically, $\lim_{x \rightarrow 0} \frac{\sin x^2}{x} = \lim_{x \rightarrow 0} x \left(\frac{\sin x^2}{x^2} \right) = 0(1) = 0$.

84. $f(x) = \frac{\sin x}{\sqrt[3]{x}}$

x	-0.1	-0.01	-0.001	0	0.001	0.01	0.1
$f(x)$	0.215	0.0464	0.01	?	0.01	0.0464	0.215

It appears that the limit is 0.



The graph has a hole at $x = 0$.

Analytically, $\lim_{x \rightarrow 0} \frac{\sin x}{\sqrt[3]{x}} = \lim_{x \rightarrow 0} \sqrt[3]{x^2} \left(\frac{\sin x}{x} \right) = (0)(1) = 0$.

85. $\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{3(x + \Delta x) - 2 - (3x - 2)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{3x + 3\Delta x - 2 - 3x + 2}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{3\Delta x}{\Delta x} = 3$

86. $\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x} - \sqrt{x}}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x + \Delta x} - \sqrt{x}}{\Delta x} \cdot \frac{\sqrt{x + \Delta x} + \sqrt{x}}{\sqrt{x + \Delta x} + \sqrt{x}}$
 $= \lim_{\Delta x \rightarrow 0} \frac{x + \Delta x - x}{\Delta x(\sqrt{x + \Delta x} + \sqrt{x})} = \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x + \Delta x} + \sqrt{x}} = \frac{1}{2\sqrt{x}}$

87. $\lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{\frac{1}{x + \Delta x + 3} - \frac{1}{x + 3}}{\Delta x}$
 $= \lim_{\Delta x \rightarrow 0} \frac{x + 3 - (x + \Delta x + 3)}{(x + \Delta x + 3)(x + 3)} \cdot \frac{1}{\Delta x}$
 $= \lim_{\Delta x \rightarrow 0} \frac{-\Delta x}{(x + \Delta x + 3)(x + 3)\Delta x}$
 $= \lim_{\Delta x \rightarrow 0} \frac{-1}{(x + \Delta x + 3)(x + 3)} = \frac{-1}{(x + 3)^2}$

$$88. \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 - 4(x + \Delta x) - (x^2 - 4x)}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x\Delta x + \Delta x^2 - 4x - 4\Delta x - x^2 + 4x}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{\Delta x(2x + \Delta x - 4)}{\Delta x} = \lim_{\Delta x \rightarrow 0} (2x + \Delta x - 4) = 2x - 4$$

$$89. \lim_{x \rightarrow 0} (4 - x^2) \leq \lim_{x \rightarrow 0} f(x) \leq \lim_{x \rightarrow 0} (4 + x^2)$$

$$4 \leq \lim_{x \rightarrow 0} f(x) \leq 4$$


Therefore, $\lim_{x \rightarrow 0} f(x) = 4$.

$$90. \lim_{x \rightarrow a} [b - |x - a|] \leq \lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} [b + |x - a|]$$

$$b \leq \lim_{x \rightarrow a} f(x) \leq b$$


Therefore, $\lim_{x \rightarrow a} f(x) = b$.

91. $f(x) = x \cos x$




$$\lim_{x \rightarrow 0} (x \cos x) = 0$$

92. $f(x) = |x \sin x|$




$$\lim_{x \rightarrow 0} |x \sin x| = 0$$

93. $f(x) = |x| \sin x$




$$\lim_{x \rightarrow 0} |x| \sin x = 0$$

94. $f(x) = |x| \cos x$




$$\lim_{x \rightarrow 0} |x| \cos x = 0$$

95. $f(x) = x \sin \frac{1}{x}$



$$\lim_{x \rightarrow 0} \left(x \sin \frac{1}{x} \right) = 0$$

96. $h(x) = x \cos \frac{1}{x}$



$$\lim_{x \rightarrow 0} \left(x \cos \frac{1}{x} \right) = 0$$

97. You say that two functions f and g agree at all but one point (on an open interval) if $f(x) = g(x)$ for all x in the interval except for $x = c$, where c is in the interval.

98. $f(x) = \frac{x^2 - 1}{x - 1}$ and $g(x) = x + 1$ agree at all points except $x = 1$.


99. An indeterminate form is obtained when evaluating a limit using direct substitution produces a meaningless fractional expression such as $0/0$. That is,

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$$

for which $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} g(x) = 0$


100. If a function f is squeezed between two functions h and g , $h(x) \leq f(x) \leq g(x)$, and h and g have the same limit L as $x \rightarrow c$, then $\lim_{x \rightarrow c} f(x)$ exists and equals L .

101. $f(x) = x$, $g(x) = \sin x$, $h(x) = \frac{\sin x}{x}$



When the x -values are "close to" 0 the magnitude of f is approximately equal to the magnitude of g . So, $|g|/|f| \approx 1$ when x is "close to" 0.

102. $f(x) = x$, $g(x) = \sin^2 x$, $h(x) = \frac{\sin^2 x}{x}$



When the x -values are "close to" 0 the magnitude of g is "smaller" than the magnitude of f and the magnitude of g is approaching zero "faster" than the magnitude of f . So, $|g|/|f| \approx 0$ when x is "close to" 0.

104. $s(t) = -16t^2 + 500 = 0$ when $t = \sqrt{\frac{500}{16}} = \frac{5\sqrt{5}}{2}$ sec. The velocity at time $a = \frac{5\sqrt{5}}{2}$ is

$$\begin{aligned} \lim_{t \rightarrow \left(\frac{5\sqrt{5}}{2}\right)} \frac{s\left(\frac{5\sqrt{5}}{2}\right) - s(t)}{\frac{5\sqrt{5}}{2} - t} &= \lim_{t \rightarrow \left(\frac{5\sqrt{5}}{2}\right)} \frac{0 - (-16t^2 + 500)}{\frac{5\sqrt{5}}{2} - t} \\ &= \lim_{t \rightarrow \left(\frac{5\sqrt{5}}{2}\right)} \frac{16\left(t^2 - \frac{125}{4}\right)}{\frac{5\sqrt{5}}{2} - t} \\ &= \lim_{t \rightarrow \left(\frac{5\sqrt{5}}{2}\right)} \frac{16\left(t + \frac{5\sqrt{5}}{2}\right)\left(t - \frac{5\sqrt{5}}{2}\right)}{\frac{5\sqrt{5}}{2} - t} \\ &= \lim_{t \rightarrow \frac{5\sqrt{5}}{2}} \left[-16\left(t + \frac{5\sqrt{5}}{2}\right)\right] = -80\sqrt{5} \text{ ft/sec} \\ &\approx -178.9 \text{ ft/sec.} \end{aligned}$$

The velocity of the wrench when it hits the ground is about 178.9 ft/sec.

103. $s(t) = -16t^2 + 500$

$$\begin{aligned} \lim_{t \rightarrow 2} \frac{s(2) - s(t)}{2 - t} &= \lim_{t \rightarrow 2} \frac{-16(2)^2 + 500 - (-16t^2 + 500)}{2 - t} \\ &= \lim_{t \rightarrow 2} \frac{436 + 16t^2 - 500}{2 - t} \\ &= \lim_{t \rightarrow 2} \frac{16(t^2 - 4)}{2 - t} \\ &= \lim_{t \rightarrow 2} \frac{16(t - 2)(t + 2)}{2 - t} \\ &= \lim_{t \rightarrow 2} -16(t + 2) = -64 \text{ ft/sec} \end{aligned}$$

The wrench is falling at about 64 feet/second.

105. $s(t) = -4.9t^2 + 200$

$$\begin{aligned} \lim_{t \rightarrow 3} \frac{s(3) - s(t)}{3 - t} &= \lim_{t \rightarrow 3} \frac{-4.9(3)^2 + 200 - (-4.9t^2 + 200)}{3 - t} \\ &= \lim_{t \rightarrow 3} \frac{4.9(t^2 - 9)}{3 - t} \\ &= \lim_{t \rightarrow 3} \frac{4.9(t - 3)(t + 3)}{3 - t} \\ &= \lim_{t \rightarrow 3} [-4.9(t + 3)] \\ &= -29.4 \text{ m/sec} \end{aligned}$$

The object is falling about 29.4 m/sec.

106. $-4.9t^2 + 200 = 0$ when $t = \sqrt{\frac{200}{4.9}} = \frac{20\sqrt{5}}{7}$ sec. The velocity at time $a = \frac{20\sqrt{5}}{7}$ is

$$\begin{aligned} \lim_{t \rightarrow a} \frac{s(a) - s(t)}{a - t} &= \lim_{t \rightarrow a} \frac{0 - [-4.9t^2 + 200]}{a - t} \\ &= \lim_{t \rightarrow a} \frac{4.9(t + a)(t - a)}{a - t} \\ &= \lim_{t \rightarrow \frac{20\sqrt{5}}{7}} \left[-4.9 \left(t + \frac{20\sqrt{5}}{7} \right) \right] = -28\sqrt{5} \text{ m/sec} \\ &\approx -62.6 \text{ m/sec.} \end{aligned}$$

The velocity of the object when it hits the ground is about 62.6 m/sec.

107. Let $f(x) = 1/x$ and $g(x) = -1/x$. $\lim_{x \rightarrow 0} f(x)$ and $\lim_{x \rightarrow 0} g(x)$ do not exist. However,

$$\lim_{x \rightarrow 0} [f(x) + g(x)] = \lim_{x \rightarrow 0} \left[\frac{1}{x} + \left(-\frac{1}{x} \right) \right] = \lim_{x \rightarrow 0} [0] = 0$$

and therefore does not exist.

108. Suppose, on the contrary, that $\lim_{x \rightarrow c} g(x)$ exists. Then, because $\lim_{x \rightarrow c} f(x)$ exists, so would $\lim_{x \rightarrow c} [f(x) + g(x)]$, which is a contradiction. So, $\lim_{x \rightarrow c} g(x)$ does not exist.

109. Given $f(x) = b$, show that for every $\varepsilon > 0$ there exists a $\delta > 0$ such that $|f(x) - b| < \varepsilon$ whenever $|x - c| < \delta$. Because $|f(x) - b| = |b - b| = 0 < \varepsilon$ for every $\varepsilon > 0$, any value of $\delta > 0$ will work.

110. Given $f(x) = x^n$, n is a positive integer, then

$$\begin{aligned} \lim_{x \rightarrow c} x^n &= \lim_{x \rightarrow c} (cx^{n-1}) \\ &= \left[\lim_{x \rightarrow c} x \right] \left[\lim_{x \rightarrow c} x^{n-1} \right] = c \left[\lim_{x \rightarrow c} (cx^{n-2}) \right] \\ &= c \left[\lim_{x \rightarrow c} x \right] \left[\lim_{x \rightarrow c} x^{n-2} \right] = c(c) \lim_{x \rightarrow c} (cx^{n-3}) \\ &= \dots = c^n. \end{aligned}$$

111. If $b = 0$, the property is true because both sides are equal to 0. If $b \neq 0$, let $\varepsilon > 0$ be given. Because $\lim_{x \rightarrow c} f(x) = L$, there exists $\delta > 0$ such that

$$|f(x) - L| < \varepsilon/|b| \text{ whenever } 0 < |x - c| < \delta. \text{ So, whenever } 0 < |x - c| < \delta, \text{ we have}$$

$$|b||f(x) - L| < \varepsilon \text{ or } |bf(x) - bL| < \varepsilon$$

which implies that $\lim_{x \rightarrow c} [bf(x)] = bL$.

112. Given $\lim_{x \rightarrow c} f(x) = 0$:

For every $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - 0| < \varepsilon \text{ whenever } 0 < |x - c| < \delta.$$

Now $|f(x) - 0| = |f(x)| = ||f(x)| - 0| < \varepsilon$ for

$$|x - c| < \delta. \text{ Therefore, } \lim_{x \rightarrow c} |f(x)| = 0.$$

113.
$$\begin{aligned} -M|f(x)| &\leq f(x)g(x) \leq M|f(x)| \\ \lim_{x \rightarrow c} (-M|f(x)|) &\leq \lim_{x \rightarrow c} f(x)g(x) \leq \lim_{x \rightarrow c} (M|f(x)|) \\ -M(0) &\leq \lim_{x \rightarrow c} f(x)g(x) \leq M(0) \\ 0 &\leq \lim_{x \rightarrow c} f(x)g(x) \leq 0 \end{aligned}$$

Therefore, $\lim_{x \rightarrow c} f(x)g(x) = 0$.

114. (a) If $\lim_{x \rightarrow c} |f(x)| = 0$, then $\lim_{x \rightarrow c} [-|f(x)|] = 0$.

$$-|f(x)| \leq f(x) \leq |f(x)|$$

$$\lim_{x \rightarrow c} [-|f(x)|] \leq \lim_{x \rightarrow c} f(x) \leq \lim_{x \rightarrow c} |f(x)|$$

$$0 \leq \lim_{x \rightarrow c} f(x) \leq 0$$

Therefore, $\lim_{x \rightarrow c} f(x) = 0$.

(b) Given $\lim_{x \rightarrow c} f(x) = L$:

For every $\varepsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - L| < \varepsilon \text{ whenever } 0 < |x - c| < \delta. \text{ Since}$$

$$||f(x)| - |L|| \leq |f(x) - L| < \varepsilon \text{ for}$$

$$|x - c| < \delta, \text{ then } \lim_{x \rightarrow c} |f(x)| = |L|.$$

115. Let

$$f(x) = \begin{cases} 4, & \text{if } x \geq 0 \\ -4, & \text{if } x < 0 \end{cases}$$

$$\lim_{x \rightarrow 0^+} |f(x)| = \lim_{x \rightarrow 0^+} 4 = 4.$$


$\lim_{x \rightarrow 0} f(x)$ does not exist because for

$$x < 0, f(x) = -4 \text{ and for } x \geq 0, f(x) = 4.$$

116. $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2} f(x) = 3$

The value of f at $x = 2$ is irrelevant.

117. The limit does not exist because the function approaches 1 from the right side of 0 and approaches -1 from the left side of 0.



118. False. $\lim_{x \rightarrow \pi} \frac{\sin x}{x} = \frac{0}{\pi} = 0$


119. True.

120. False. Let

$$f(x) = \begin{cases} x & x \neq 1 \\ 3 & x = 1 \end{cases}, \quad c = 1.$$

Then $\lim_{x \rightarrow 1} f(x) = 1$ but $f(1) \neq 1$.

121. False. The limit does not exist because $f(x)$ approaches 3 from the left side of 2 and approaches 0 from the right side of 2.



122. False. Let

$$f(x) = \frac{1}{2}x^2 \text{ and } g(x) = x^2.$$

Then $f(x) < g(x)$ for all $x \neq 0$. But

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} g(x) = 0.$$

$$\begin{aligned} 123. \lim_{x \rightarrow 0} \frac{1 - \cos x}{x} &= \lim_{x \rightarrow 0} \frac{1 - \cos x}{x} \cdot \frac{1 + \cos x}{1 + \cos x} \\ &= \lim_{x \rightarrow 0} \frac{1 - \cos^2 x}{x(1 + \cos x)} = \lim_{x \rightarrow 0} \frac{\sin^2 x}{x(1 + \cos x)} \\ &= \lim_{x \rightarrow 0} \frac{\sin x}{x} \cdot \frac{\sin x}{1 + \cos x} \\ &= \left[\lim_{x \rightarrow 0} \frac{\sin x}{x} \right] \left[\lim_{x \rightarrow 0} \frac{\sin x}{1 + \cos x} \right] \\ &= (1)(0) = 0 \end{aligned}$$

124. $f(x) = \begin{cases} 0, & \text{if } x \text{ is rational} \\ 1, & \text{if } x \text{ is irrational} \end{cases}$

$g(x) = \begin{cases} 0, & \text{if } x \text{ is rational} \\ x, & \text{if } x \text{ is irrational} \end{cases}$

$\lim_{x \rightarrow 0} f(x)$ does not exist.

No matter how "close to" 0 x is, there are still an infinite number of rational and irrational numbers so that


$\lim_{x \rightarrow 0} f(x)$ does not exist.

$$\lim_{x \rightarrow 0} g(x) = 0$$

when x is "close to" 0, both parts of the function are "close to" 0.

125. $f(x) = \frac{\sec x - 1}{x^2}$

(a) The domain of f is all $x \neq 0, \pi/2 + n\pi$.



The domain is not obvious. The hole at $x = 0$ is not apparent.

(c) $\lim_{x \rightarrow 0} f(x) = \frac{1}{2}$

$$\begin{aligned} \text{(d)} \quad \frac{\sec x - 1}{x^2} &= \frac{\sec x - 1}{x^2} \cdot \frac{\sec x + 1}{\sec x + 1} = \frac{\sec^2 x - 1}{x^2(\sec x + 1)} \\ &= \frac{\tan^2 x}{x^2(\sec x + 1)} = \frac{1}{\cos^2 x} \left(\frac{\sin^2 x}{x^2} \right) \frac{1}{\sec x + 1} \end{aligned}$$

$$\begin{aligned} \text{So, } \lim_{x \rightarrow 0} \frac{\sec x - 1}{x^2} &= \lim_{x \rightarrow 0} \frac{1}{\cos^2 x} \left(\frac{\sin^2 x}{x^2} \right) \frac{1}{\sec x + 1} \\ &= 1(1) \left(\frac{1}{2} \right) = \frac{1}{2}. \end{aligned}$$

$$\begin{aligned}
 126. (a) \quad \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} &= \lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} \cdot \frac{1 + \cos x}{1 + \cos x} \\
 &= \lim_{x \rightarrow 0} \frac{1 - \cos^2 x}{x^2(1 + \cos x)} \\
 &= \lim_{x \rightarrow 0} \frac{\sin^2 x}{x^2} \cdot \frac{1}{1 + \cos x} \\
 &= (1) \left(\frac{1}{2} \right) = \frac{1}{2}
 \end{aligned}$$

(b) From part (a),

$$\frac{1 - \cos x}{x^2} \approx \frac{1}{2} \Rightarrow 1 - \cos x \approx \frac{1}{2}x^2 \Rightarrow \cos x \approx 1 - \frac{1}{2}x^2 \text{ for } x \approx 0.$$

(c) $\cos(0.1) \approx 1 - \frac{1}{2}(0.1)^2 = 0.995$

(d) $\cos(0.1) \approx 0.9950$, which agrees with part (c).

127. The graphing utility was set in degree mode, instead of *radian* mode.

Section 1.4 Continuity and One-Sided Limits

1. (a) $\lim_{x \rightarrow 4^+} f(x) = 3$

(b) $\lim_{x \rightarrow 4^-} f(x) = 3$

(c) $\lim_{x \rightarrow 4} f(x) = 3$

The function is continuous at $x = 4$ and is continuous on $(-\infty, \infty)$.

2. (a) $\lim_{x \rightarrow -2^+} f(x) = -2$

(b) $\lim_{x \rightarrow -2^-} f(x) = -2$

(c) $\lim_{x \rightarrow -2} f(x) = -2$

The function is continuous at $x = -2$.

3. (a) $\lim_{x \rightarrow 3^+} f(x) = 0$

(b) $\lim_{x \rightarrow 3^-} f(x) = 0$

(c) $\lim_{x \rightarrow 3} f(x) = 0$

The function is NOT continuous at $x = 3$.

4. (a) $\lim_{x \rightarrow -3^+} f(x) = 3$

(b) $\lim_{x \rightarrow -3^-} f(x) = 3$

(c) $\lim_{x \rightarrow -3} f(x) = 3$

The function is NOT continuous at $x = -3$ because $f(-3) = 4 \neq \lim_{x \rightarrow -3} f(x)$.

5. (a) $\lim_{x \rightarrow 2^+} f(x) = -3$

(b) $\lim_{x \rightarrow 2^-} f(x) = 3$

(c) $\lim_{x \rightarrow 2} f(x)$ does not exist

The function is NOT continuous at $x = 2$.

6. (a) $\lim_{x \rightarrow -1^+} f(x) = 0$

(b) $\lim_{x \rightarrow -1^-} f(x) = 2$

(c) $\lim_{x \rightarrow -1} f(x)$ does not exist.

The function is NOT continuous at $x = -1$.

7. $\lim_{x \rightarrow 8^+} \frac{1}{x + 8} = \frac{1}{8 + 8} = \frac{1}{16}$

8. $\lim_{x \rightarrow 5^-} -\frac{3}{x + 5} = -\frac{3}{5 + 5} = -\frac{3}{10}$

9. $\lim_{x \rightarrow 5^+} \frac{x - 5}{x^2 - 25} = \lim_{x \rightarrow 5^+} \frac{1}{x + 5} = \frac{1}{10}$

10. $\lim_{x \rightarrow 2^+} \frac{2 - x}{x^2 - 4} = \lim_{x \rightarrow 2^+} -\frac{1}{x + 2} = -\frac{1}{4}$

11. $\lim_{x \rightarrow -3^-} \frac{x}{\sqrt{x^2 - 9}}$ does not exist because

$\frac{x}{\sqrt{x^2 - 9}}$ decreases without bound as $x \rightarrow -3^-$.

$$\begin{aligned} 12. \lim_{x \rightarrow 9^-} \frac{\sqrt{x} - 3}{x - 9} &= \lim_{x \rightarrow 9^-} \frac{\sqrt{x} - 3}{x - 9} \cdot \frac{\sqrt{x} + 3}{\sqrt{x} + 3} \\ &= \lim_{x \rightarrow 9^-} \frac{x - 9}{(x - 9)(\sqrt{x} + 3)} \\ &= \lim_{x \rightarrow 9^-} \frac{1}{\sqrt{x} + 3} = \frac{1}{6} \end{aligned}$$

$$13. \lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} \frac{-x}{x} = -1$$

$$14. \lim_{x \rightarrow 10^+} \frac{|x - 10|}{x - 10} = \lim_{x \rightarrow 10^+} \frac{x - 10}{x - 10} = 1$$

$$\begin{aligned} 15. \lim_{\Delta x \rightarrow 0^-} \frac{\frac{1}{x + \Delta x} - \frac{1}{x}}{\Delta x} &= \lim_{\Delta x \rightarrow 0^-} \frac{x - (x + \Delta x)}{x(x + \Delta x)} \cdot \frac{1}{\Delta x} = \lim_{\Delta x \rightarrow 0^-} \frac{-\Delta x}{x(x + \Delta x)} \cdot \frac{1}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0^-} \frac{-1}{x(x + \Delta x)} \\ &= \frac{-1}{x(x + 0)} = -\frac{1}{x^2} \end{aligned}$$

$$\begin{aligned} 16. \lim_{\Delta x \rightarrow 0^+} \frac{(x + \Delta x)^2 + (x + \Delta x) - (x^2 + x)}{\Delta x} &= \lim_{\Delta x \rightarrow 0^+} \frac{x^2 + 2x(\Delta x) + (\Delta x)^2 + x + \Delta x - x^2 - x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0^+} \frac{2x(\Delta x) + (\Delta x)^2 + \Delta x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0^+} (2x + \Delta x + 1) \\ &= 2x + 0 + 1 = 2x + 1 \end{aligned}$$

$$17. \lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} \frac{x + 2}{2} = \frac{5}{2}$$

$$24. \lim_{x \rightarrow 2^+} (2x - \lfloor x \rfloor) = 2(2) - 2 = 2$$

$$\begin{aligned} 18. \lim_{x \rightarrow 2^+} f(x) &= \lim_{x \rightarrow 2^+} (-x^2 + 4x - 2) = 2 \\ \lim_{x \rightarrow 2^-} f(x) &= \lim_{x \rightarrow 2^-} (x^2 - 4x + 6) = 2 \\ \lim_{x \rightarrow 2} f(x) &= 2 \end{aligned}$$

$$25. \lim_{x \rightarrow 3} (2 - \lfloor -x \rfloor) \text{ does not exist because}$$

$$\lim_{x \rightarrow 3^-} (2 - \lfloor -x \rfloor) = 2 - (-3) = 5$$

and

$$\lim_{x \rightarrow 3^+} (2 - \lfloor -x \rfloor) = 2 - (-4) = 6.$$

$$19. \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x + 1) = 2$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^3 + 1) = 2$$

$$\lim_{x \rightarrow 1} f(x) = 2$$

$$26. \lim_{x \rightarrow 1} \left(1 - \left\lfloor -\frac{x}{2} \right\rfloor \right) = 1 - (-1) = 2$$

$$27. f(x) = \frac{1}{x^2 - 4}$$

has discontinuities at $x = -2$ and $x = 2$ because $f(-2)$ and $f(2)$ are not defined.

$$20. \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (1 - x) = 0$$

$$21. \lim_{x \rightarrow \pi} \cot x \text{ does not exist because}$$

$$\lim_{x \rightarrow \pi^+} \cot x \text{ and } \lim_{x \rightarrow \pi^-} \cot x \text{ do not exist.}$$

$$28. f(x) = \frac{x^2 - 1}{x + 1}$$

has a discontinuity at $x = -1$ because $f(-1)$ is not defined.

$$22. \lim_{x \rightarrow \pi/2} \sec x \text{ does not exist because}$$

$$\lim_{x \rightarrow (\pi/2)^+} \sec x \text{ and } \lim_{x \rightarrow (\pi/2)^-} \sec x \text{ do not exist.}$$

$$23. \lim_{x \rightarrow 4^-} (5\lfloor x \rfloor - 7) = 5(3) - 7 = 8$$

$$(\lfloor x \rfloor = 3 \text{ for } 3 \leq x < 4)$$

$$29. f(x) = \frac{\lfloor x \rfloor}{2} + x$$

has discontinuities at each integer k because

$$\lim_{x \rightarrow k^-} f(x) \neq \lim_{x \rightarrow k^+} f(x).$$

30. $f(x) = \begin{cases} x, & x < 1 \\ 2, & x = 1 \\ 2x - 1, & x > 1 \end{cases}$ has a discontinuity at $x = 1$ because $f(1) = 2 \neq \lim_{x \rightarrow 1} f(x) = 1$.

31. $g(x) = \sqrt{49 - x^2}$ is continuous on $[-7, 7]$.

32. $f(t) = 3 - \sqrt{9 - t^2}$ is continuous on $[-3, 3]$.

33. $\lim_{x \rightarrow 0^-} f(x) = 3 = \lim_{x \rightarrow 0^+} f(x)$. f is continuous on $[-1, 4]$.

34. $g(2)$ is not defined. g is continuous on $[-1, 2)$.

35. $f(x) = \frac{6}{x}$ has a nonremovable discontinuity at $x = 0$.

36. $f(x) = \frac{3}{x - 2}$ has a nonremovable discontinuity at $x = 2$.

37. $f(x) = x^2 - 9$ is continuous for all real x .

38. $f(x) = x^2 - 2x + 1$ is continuous for all real x .

39. $f(x) = \frac{1}{4 - x^2} = \frac{1}{(2 - x)(2 + x)}$ has nonremovable discontinuities at $x = \pm 2$ because $\lim_{x \rightarrow 2} f(x)$ and $\lim_{x \rightarrow -2} f(x)$ do not exist.

40. $f(x) = \frac{1}{x^2 + 1}$ is continuous for all real x .

41. $f(x) = 3x - \cos x$ is continuous for all real x .

42. $f(x) = \cos \frac{\pi x}{2}$ is continuous for all real x .

43. $f(x) = \frac{x}{x^2 - x}$ is not continuous at $x = 0, 1$. Because $\frac{x}{x^2 - x} = \frac{1}{x - 1}$ for $x \neq 0, x = 0$ is a removable discontinuity, whereas $x = 1$ is a nonremovable discontinuity.

44. $f(x) = \frac{x}{x^2 - 1}$ has nonremovable discontinuities at $x = 1$ and $x = -1$ because $\lim_{x \rightarrow 1} f(x)$ and $\lim_{x \rightarrow -1} f(x)$ do not exist.

45. $f(x) = \frac{x}{x^2 + 1}$ is continuous for all real x .

46. $f(x) = \frac{x - 6}{x^2 - 36}$

has a nonremovable discontinuity at $x = -6$ because $\lim_{x \rightarrow -6} f(x)$ does not exist, and has a removable discontinuity at $x = 6$ because

$$\lim_{x \rightarrow 6} f(x) = \lim_{x \rightarrow 6} \frac{1}{x + 6} = \frac{1}{12}$$

47. $f(x) = \frac{x + 2}{(x + 2)(x - 5)}$

has a nonremovable discontinuity at $x = 5$ because $\lim_{x \rightarrow 5} f(x)$ does not exist, and has a removable discontinuity at $x = -2$ because

$$\lim_{x \rightarrow -2} f(x) = \lim_{x \rightarrow -2} \frac{1}{x - 5} = -\frac{1}{7}$$

48. $f(x) = \frac{x - 1}{(x + 2)(x - 1)}$

has a nonremovable discontinuity at $x = -2$ because $\lim_{x \rightarrow -2} f(x)$ does not exist, and has a removable discontinuity at $x = 1$ because

$$\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} \frac{1}{x + 2} = \frac{1}{3}$$

49. $f(x) = \frac{|x + 7|}{x + 7}$

has a nonremovable discontinuity at $x = -7$ because $\lim_{x \rightarrow -7} f(x)$ does not exist.

50. $f(x) = \frac{|x - 8|}{x - 8}$ has a nonremovable discontinuity at $x = 8$ because $\lim_{x \rightarrow 8} f(x)$ does not exist.

51. $f(x) = \begin{cases} x, & x \leq 1 \\ x^2, & x > 1 \end{cases}$

has a **possible** discontinuity at $x = 1$.

1. $f(1) = 1$

2. $\left. \begin{aligned} \lim_{x \rightarrow 1^-} f(x) &= \lim_{x \rightarrow 1^-} x = 1 \\ \lim_{x \rightarrow 1^+} f(x) &= \lim_{x \rightarrow 1^+} x^2 = 1 \end{aligned} \right\} \lim_{x \rightarrow 1} f(x) = 1$

3. $f(-1) = \lim_{x \rightarrow -1} f(x)$

f is continuous at $x = 1$, therefore, f is continuous for all real x .

$$52. f(x) = \begin{cases} -2x + 3, & x < 1 \\ x^2, & x \geq 1 \end{cases}$$

has a **possible** discontinuity at $x = 1$.

$$1. f(1) = 1^2 = 1$$

$$2. \left. \begin{array}{l} \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (-2x + 3) = 1 \\ \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} x^2 = 1 \end{array} \right\} \lim_{x \rightarrow 1} f(x) = 1$$

$$3. f(1) = \lim_{x \rightarrow 1} f(x)$$

f is continuous at $x = 1$, therefore, f is continuous for all real x .

$$53. f(x) = \begin{cases} \frac{x}{2} + 1, & x \leq 2 \\ 3 - x, & x > 2 \end{cases}$$

has a **possible** discontinuity at $x = 2$.

$$1. f(2) = \frac{2}{2} + 1 = 2$$

$$2. \left. \begin{array}{l} \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} \left(\frac{x}{2} + 1 \right) = 2 \\ \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (3 - x) = 1 \end{array} \right\} \lim_{x \rightarrow 2} f(x) \text{ does not exist.}$$

Therefore, f has a nonremovable discontinuity at $x = 2$.

$$54. f(x) = \begin{cases} -2x, & x \leq 2 \\ x^2 - 4x + 1, & x > 2 \end{cases}$$

has a **possible** discontinuity at $x = 2$.

$$1. f(2) = -2(2) = -4$$

$$2. \left. \begin{array}{l} \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (-2x) = -4 \\ \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x^2 - 4x + 1) = -3 \end{array} \right\} \lim_{x \rightarrow 2} f(x) \text{ does not exist.}$$

Therefore, f has a nonremovable discontinuity at $x = 2$.

$$55. f(x) = \begin{cases} \tan \frac{\pi x}{4}, & |x| < 1 \\ x, & |x| \geq 1 \end{cases}$$

$$= \begin{cases} \tan \frac{\pi x}{4}, & -1 < x < 1 \\ x, & x \leq -1 \text{ or } x \geq 1 \end{cases}$$

has **possible** discontinuities at $x = -1, x = 1$.

$$1. f(-1) = -1 \qquad f(1) = 1$$

$$2. \lim_{x \rightarrow -1} f(x) = -1 \qquad \lim_{x \rightarrow 1} f(x) = 1$$

$$3. f(-1) = \lim_{x \rightarrow -1} f(x) \qquad f(1) = \lim_{x \rightarrow 1} f(x)$$

f is continuous at $x = \pm 1$, therefore, f is continuous for all real x .

$$56. f(x) = \begin{cases} \csc \frac{\pi x}{6}, & |x - 3| \leq 2 \\ 2, & |x - 3| > 2 \end{cases}$$

$$= \begin{cases} \csc \frac{\pi x}{6}, & 1 \leq x \leq 5 \\ 2, & x < 1 \text{ or } x > 5 \end{cases}$$

has **possible** discontinuities at $x = 1, x = 5$.

$$1. f(1) = \csc \frac{\pi}{6} = 2 \qquad f(5) = \csc \frac{5\pi}{6} = 2$$

$$2. \lim_{x \rightarrow 1^-} f(x) = 2 \qquad \lim_{x \rightarrow 5^-} f(x) = 2$$

$$3. f(1) = \lim_{x \rightarrow 1^-} f(x) \qquad f(5) = \lim_{x \rightarrow 5^-} f(x)$$

f is continuous at $x = 1$ and $x = 5$, therefore, f is continuous for all real x .

57. $f(x) = \csc 2x$ has nonremovable discontinuities at integer multiples of $\pi/2$.

58. $f(x) = \tan \frac{\pi x}{2}$ has nonremovable discontinuities at each $2k + 1$, k is an integer.


59. $f(x) = \llbracket x - 8 \rrbracket$ has nonremovable discontinuities at each integer k .

60. $f(x) = 5 - \llbracket x \rrbracket$ has nonremovable discontinuities at each integer k .

$$61. \lim_{x \rightarrow 0^+} f(x) = 0$$

$$\lim_{x \rightarrow 0^-} f(x) = 0$$


f is not continuous at $x = -2$.



$$62. \lim_{x \rightarrow 0^+} f(x) = 0$$

$$\lim_{x \rightarrow 0^-} f(x) = 0$$

f is not continuous at $x = -4$.



$$63. f(1) = 3$$

$$\text{Find } a \text{ so that } \lim_{x \rightarrow 1^-} (ax - 4) = 3$$

$$a(1) - 4 = 3$$

$$a = 7.$$

$$64. f(1) = 3$$

$$\text{Find } a \text{ so that } \lim_{x \rightarrow 1^+} (ax + 5) = 3$$

$$a(1) + 5 = 3$$

$$a = -2.$$

$$65. f(2) = 8$$

$$\text{Find } a \text{ so that } \lim_{x \rightarrow 2^+} ax^2 = 8 \Rightarrow a = \frac{8}{2^2} = 2.$$

$$66. \lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} \frac{4 \sin x}{x} = 4$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (a - 2x) = a$$

Let $a = 4$.

67. Find a and b such that $\lim_{x \rightarrow -1^+} (ax + b) = -a + b = 2$ and $\lim_{x \rightarrow 3^-} (ax + b) = 3a + b = -2$.

$$\begin{aligned} a - b &= -2 \\ (+)3a + b &= -2 \\ \hline 4a &= -4 \\ a &= -1 \\ b &= 2 + (-1) = 1 \end{aligned} \quad f(x) = \begin{cases} 2, & x \leq -1 \\ -x + 1, & -1 < x < 3 \\ -2, & x \geq 3 \end{cases}$$

68. $\lim_{x \rightarrow a} g(x) = \lim_{x \rightarrow a} \frac{x^2 - a^2}{x - a} = \lim_{x \rightarrow a} (x + a) = 2a$

Find a such $2a = 8 \Rightarrow a = 4$.

69. $f(g(x)) = (x - 1)^2$
Continuous for all real x .

70. $f(g(x)) = \frac{1}{\sqrt{x-1}}$

Nonremovable discontinuity at $x = 1$. Continuous for all $x > 1$.


71. $f(g(x)) = \frac{1}{(x^2 + 5) - 6} = \frac{1}{x^2 - 1}$

Nonremovable discontinuities at $x = \pm 1$

72. $f(g(x)) = \sin x^2$
Continuous for all real x


73. $y = \llbracket x \rrbracket - x$

Nonremovable discontinuity at each integer




74. $h(x) = \frac{1}{(x+1)(x-2)}$

Nonremovable discontinuities at $x = -1$ and $x = 2$.



75. $g(x) = \begin{cases} x^2 - 3x, & x > 4 \\ 2x - 5, & x \leq 4 \end{cases}$

There is a nonremovable discontinuity at $x = 4$.



76. $f(x) = \begin{cases} \frac{\cos x - 1}{x}, & x < 0 \\ 5x, & x \geq 0 \end{cases}$


$$f(0) = 5(0) = 0$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \frac{\cos x - 1}{x} = 0$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (5x) = 0$$

Therefore, $\lim_{x \rightarrow 0} f(x) = 0 = f(0)$ and f is continuous on the entire real line.

($x = 0$ was the only possible discontinuity.)



77. $f(x) = \frac{x}{x^2 + x + 2}$

Continuous on $(-\infty, \infty)$

78. $f(x) = x\sqrt{x+3}$

Continuous on $[-3, \infty)$

79. $f(x) = \sec \frac{\pi x}{4}$


Continuous on:

..., $(-6, -2), (-2, 2), (2, 6), (6, 10), \dots$

80. $f(x) = \frac{x+1}{\sqrt{x}}$


Continuous on $(0, \infty)$

81. $f(x) = \frac{\sin x}{x}$



The graph **appears** to be continuous on the interval $[-4, 4]$. Because $f(0)$ is not defined, you know that f has a discontinuity at $x = 0$. This discontinuity is removable so it does not show up on the graph.

82. $f(x) = \frac{x^3 - 8}{x - 2}$



The graph **appears** to be continuous on the interval $[-4, 4]$. Because $f(2)$ is not defined, you know that f has a discontinuity at $x = 2$. This discontinuity is removable so it does not show up on the graph.

83. $f(x) = \frac{1}{12}x^4 - x^3 + 4$ is continuous on the interval $[1, 2]$. $f(1) = \frac{37}{12}$ and $f(2) = -\frac{8}{3}$. By the Intermediate Value Theorem, there exists a number c in $[1, 2]$ such that $f(c) = 0$.

84. $f(x) = x^3 + 5x - 3$ is continuous on the interval $[0, 1]$. $f(0) = -3$ and $f(1) = 3$. By the Intermediate Value Theorem, there exists a number c in $[0, 1]$ such that $f(c) = 0$.

85. $f(x) = x^2 - 2 - \cos x$ is continuous on $[0, \pi]$. $f(0) = -3$ and $f(\pi) = \pi^2 - 1 \approx 8.87 > 0$. By the Intermediate Value Theorem, $f(c) = 0$ for at least one value of c between 0 and π .

86. $f(x) = -\frac{5}{x} + \tan\left(\frac{\pi x}{10}\right)$ is continuous on the interval $[1, 4]$.

$$f(1) = -5 + \tan\left(\frac{\pi}{10}\right) \approx -4.7 \text{ and}$$

$$f(4) = -\frac{5}{4} + \tan\left(\frac{2\pi}{5}\right) \approx 1.8. \text{ By the Intermediate}$$

Value Theorem, there exists a number c in $[1, 4]$ such that $f(c) = 0$.

87. $f(x) = x^3 + x - 1$

$f(x)$ is continuous on $[0, 1]$.

$$f(0) = -1 \text{ and } f(1) = 1$$

By the Intermediate Value Theorem, $f(c) = 0$ for at least one value of c between 0 and 1. Using a graphing utility to zoom in on the graph of $f(x)$, you find that $x \approx 0.68$. Using the root feature, you find that $x \approx 0.6823$.

88. $f(x) = x^3 + 5x - 3$

$f(x)$ is continuous on $[0, 1]$.

$$f(0) = -3 \text{ and } f(1) = 3$$

By the Intermediate Value Theorem, $f(c) = 0$ for at least one value of c between 0 and 1. Using a graphing utility to zoom in on the graph of $f(x)$, you find that $x \approx 0.56$. Using the root feature, you find that $x \approx 0.5641$.

89. $g(t) = 2 \cos t - 3t$

g is continuous on $[0, 1]$.

$$g(0) = 2 > 0 \text{ and } g(1) \approx -1.9 < 0.$$

By the Intermediate Value Theorem, $g(c) = 0$ for at least one value of c between 0 and 1. Using a graphing utility to zoom in on the graph of $g(t)$, you find that $t \approx 0.56$. Using the root feature, you find that $t \approx 0.5636$.

90. $h(\theta) = 1 + \theta - 3 \tan \theta$

h is continuous on $[0, 1]$.

$$h(0) = 1 > 0 \text{ and } h(1) \approx -2.67 < 0.$$

By the Intermediate Value Theorem, $h(c) = 0$ for at least one value of c between 0 and 1. Using a graphing utility to zoom in on the graph of $h(\theta)$, you find that $\theta \approx 0.45$. Using the root feature, you find that $\theta \approx 0.4503$.

91. $f(x) = x^2 + x - 1$

 f is continuous on $[0, 5]$.

$f(0) = -1$ and $f(5) = 29$

$-1 < 11 < 29$

The Intermediate Value Theorem applies.

$x^2 + x - 1 = 11$

$x^2 + x - 12 = 0$

$(x + 4)(x - 3) = 0$

$x = -4$ or $x = 3$

$c = 3$ ($x = -4$ is not in the interval.)

So, $f(3) = 11$.

92. $f(x) = x^2 - 6x + 8$

 f is continuous on $[0, 3]$.

$f(0) = 8$ and $f(3) = -1$

$-1 < 0 < 8$

The Intermediate Value Theorem applies.

$x^2 - 6x + 8 = 0$

$(x - 2)(x - 4) = 0$

$x = 2$ or $x = 4$

$c = 2$ ($x = 4$ is not in the interval.)

So, $f(2) = 0$.

93. $f(x) = x^3 - x^2 + x - 2$

 f is continuous on $[0, 3]$.

$f(0) = -2$ and $f(3) = 19$

$-2 < 4 < 19$

The Intermediate Value Theorem applies.

$x^3 - x^2 + x - 2 = 4$

$x^3 - x^2 + x - 6 = 0$

$(x - 2)(x^2 + x + 3) = 0$

$x = 2$

 $(x^2 + x + 3)$ has no real solution.)

$c = 2$

So, $f(2) = 4$.

94. $f(x) = \frac{x^2 + x}{x - 1}$

 f is continuous on $\left[\frac{5}{2}, 4\right]$. The nonremovable discontinuity, $x = 1$, lies outside the interval.

$f\left(\frac{5}{2}\right) = \frac{35}{6}$ and $f(4) = \frac{20}{3}$

$\frac{35}{6} < 6 < \frac{20}{3}$

The Intermediate Value Theorem applies.

$\frac{x^2 + x}{x - 1} = 6$

$x^2 + x = 6x - 6$

$x^2 - 5x + 6 = 0$


$(x - 2)(x - 3) = 0$

$x = 2$ or $x = 3$

$c = 3$ ($x = 2$ is not in the interval.)

So, $f(3) = 6$.

95. (a) The limit does not exist at $x = c$.
 (b) The function is not defined at $x = c$.
 (c) The limit exists at $x = c$, but it is not equal to the value of the function at $x = c$.
 (d) The limit does not exist at $x = c$.

96. Answers will vary. *Sample answer:*The function is not continuous at $x = 3$ because

$$\lim_{x \rightarrow 3^+} f(x) = 1 \neq 0 = \lim_{x \rightarrow 3^-} f(x).$$

97. If f and g are continuous for all real x , then so is $f + g$ (Theorem 1.11, part 2). However, f/g might not be continuous if $g(x) = 0$. For example, let $f(x) = x$ and $g(x) = x^2 - 1$. Then f and g are continuous for all real x , but f/g is not continuous at $x = \pm 1$.


98. A discontinuity at c is removable if the function f can be made continuous at c by appropriately defining (or redefining) $f(c)$. Otherwise, the discontinuity is nonremovable.

(a) $f(x) = \frac{|x - 4|}{x - 4}$

(b) $f(x) = \frac{\sin(x + 4)}{x + 4}$

(c) $f(x) = \begin{cases} 1, & x \geq 4 \\ 0, & -4 < x < 4 \\ 1, & x = -4 \\ 0, & x < -4 \end{cases}$

$x = 4$ is nonremovable, $x = -4$ is removable



99. True

1. $f(c) = L$ is defined.
2. $\lim_{x \rightarrow c} f(x) = L$ exists.
3. $f(c) = \lim_{x \rightarrow c} f(x)$

All of the conditions for continuity are met.

100. True. If $f(x) = g(x)$, $x \neq c$, then

$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} g(x)$ (if they exist) and at least one of these limits then does not equal the corresponding function value at $x = c$.

101. False. A rational function can be written as $P(x)/Q(x)$ where P and Q are polynomials of degree m and n , respectively. It can have, at most, n discontinuities.

102. False. $f(1)$ is not defined and $\lim_{x \rightarrow 1} f(x)$ does not exist.

103. $\lim_{t \rightarrow 4^-} f(t) \approx 28$
 $\lim_{t \rightarrow 4^+} f(t) \approx 56$

At the end of day 3, the amount of chlorine in the pool has decreased to about 28 oz. At the beginning of day 4, more chlorine was added, and the amount is now about 56 oz.

104. The functions agree for integer values of x :

$$\left. \begin{aligned} g(x) &= 3 - \lfloor -x \rfloor = 3 - (-x) = 3 + x \\ f(x) &= 3 + \lceil x \rceil = 3 + x \end{aligned} \right\} \text{for } x \text{ an integer}$$


However, for non-integer values of x , the functions differ by 1.

$$f(x) = 3 + \lceil x \rceil = g(x) - 1 = 2 - \lfloor -x \rfloor.$$

For example,

$$f\left(\frac{1}{2}\right) = 3 + 0 = 3, g\left(\frac{1}{2}\right) = 3 - (-1) = 4.$$

105. $C(t) = \begin{cases} 0.40, & 0 < t \leq 10 \\ 0.40 + 0.05\lceil t - 9 \rceil, & t > 10, t \text{ not an integer} \\ 0.40 + 0.05(t - 10), & t > 10, t \text{ an integer} \end{cases}$



There is a nonremovable discontinuity at each integer greater than or equal to 10.


Note: You could also express C as

$$C(t) = \begin{cases} 0.40, & 0 < t \leq 10 \\ 0.40 - 0.05\lfloor 10 - t \rfloor, & t > 10 \end{cases}$$

106. $N(t) = 25\left(2\left\lceil\frac{t+2}{2}\right\rceil - t\right)$

t	0	1	1.8	2	3	3.8
$N(t)$	50	25	5	50	25	5

Discontinuous at every positive even integer. The company replenishes its inventory every two months.



107. Let $s(t)$ be the position function for the run up to the campsite. $s(0) = 0$ ($t = 0$ corresponds to 8:00 A.M., $s(20) = k$ (distance to campsite)). Let $r(t)$ be the position function for the run back down the mountain: $r(0) = k, r(10) = 0$. Let $f(t) = s(t) - r(t)$.

When $t = 0$ (8:00 A.M.),
 $f(0) = s(0) - r(0) = 0 - k < 0$.

When $t = 10$ (8:00 A.M.), $f(10) = s(10) - r(10) > 0$.

Because $f(0) < 0$ and $f(10) > 0$, then there must be a value t in the interval $[0, 10]$ such that $f(t) = 0$. If

$f(t) = 0$, then $s(t) - r(t) = 0$, which gives us

$s(t) = r(t)$. Therefore, at some time t , where

$0 \leq t \leq 10$, the position functions for the run up and the run down are equal.

108. Let $V = \frac{4}{3}\pi r^3$ be the volume of a sphere with radius r .

V is continuous on $[5, 8]$. $V(5) = \frac{500\pi}{3} \approx 523.6$ and

$V(8) = \frac{2048\pi}{3} \approx 2144.7$. Because

$523.6 < 1500 < 2144.7$, the Intermediate Value Theorem guarantees that there is at least one value r between 5 and 8 such that $V(r) = 1500$. (In fact, $r \approx 7.1012$.)

109. Suppose there exists x_1 in $[a, b]$ such that

$f(x_1) > 0$ and there exists x_2 in $[a, b]$ such that

$f(x_2) < 0$. Then by the Intermediate Value Theorem,

$f(x)$ must equal zero for some value of x in

$[x_1, x_2]$ (or $[x_2, x_1]$ if $x_2 < x_1$). So, f would have a zero in $[a, b]$, which is a contradiction. Therefore, $f(x) > 0$ for all x in $[a, b]$ or $f(x) < 0$ for all x in $[a, b]$.

110. Let c be any real number. Then $\lim_{x \rightarrow c} f(x)$ does not exist because there are both rational and irrational numbers arbitrarily close to c . Therefore, f is not continuous at c .

111. If $x = 0$, then $f(0) = 0$ and $\lim_{x \rightarrow 0} f(x) = 0$. So, f is continuous at $x = 0$.

If $x \neq 0$, then $\lim_{t \rightarrow x} f(t) = 0$ for x rational, whereas


$\lim_{t \rightarrow x} f(t) = \lim_{t \rightarrow x} kt = kx \neq 0$ for x irrational. So, f is not continuous for all $x \neq 0$.

$$112. \operatorname{sgn}(x) = \begin{cases} -1, & \text{if } x < 0 \\ 0, & \text{if } x = 0 \\ 1, & \text{if } x > 0 \end{cases}$$


(a) $\lim_{x \rightarrow 0^-} \operatorname{sgn}(x) = -1$

(b) $\lim_{x \rightarrow 0^+} \operatorname{sgn}(x) = 1$

(c) $\lim_{x \rightarrow 0} \operatorname{sgn}(x)$ does not exist.




113. (a)




(b) There appears to be a limiting speed and a possible cause is air resistance.

114. (a) $f(x) = \begin{cases} 0, & 0 \leq x < b \\ b, & b < x \leq 2b \end{cases}$



NOT continuous at $x = b$.

(b) $g(x) = \begin{cases} \frac{x}{2}, & 0 \leq x \leq b \\ b - \frac{x}{2}, & b < x \leq 2b \end{cases}$



Continuous on $[0, 2b]$.

115. $f(x) = \begin{cases} 1 - x^2, & x \leq c \\ x, & x > c \end{cases}$

f is continuous for $x < c$ and for $x > c$. At $x = c$, you need $1 - c^2 = c$. Solving $c^2 + c - 1$, you obtain

$$c = \frac{-1 \pm \sqrt{1+4}}{2} = \frac{-1 \pm \sqrt{5}}{2}.$$

117. $f(x) = \frac{\sqrt{x+c^2} - c}{x}, c > 0$

Domain: $x + c^2 \geq 0 \Rightarrow x \geq -c^2$ and $x \neq 0, [-c^2, 0) \cup (0, \infty)$

$$\lim_{x \rightarrow 0} \frac{\sqrt{x+c^2} - c}{x} = \lim_{x \rightarrow 0} \frac{\sqrt{x+c^2} - c}{x} \cdot \frac{\sqrt{x+c^2} + c}{\sqrt{x+c^2} + c} = \lim_{x \rightarrow 0} \frac{(x+c^2) - c^2}{x[\sqrt{x+c^2} + c]} = \lim_{x \rightarrow 0} \frac{1}{\sqrt{x+c^2} + c} = \frac{1}{2c}$$

Define $f(0) = 1/(2c)$ to make f continuous at $x = 0$.

118. 1. $f(c)$ is defined.


2. $\lim_{x \rightarrow c} f(x) = \lim_{\Delta x \rightarrow 0} f(c + \Delta x) = f(c)$ exists.

[Let $x = c + \Delta x$. As $x \rightarrow c$, $\Delta x \rightarrow 0$]

3. $\lim_{x \rightarrow c} f(x) = f(c)$.

Therefore, f is continuous at $x = c$.

119. $h(x) = x \llbracket x \rrbracket$



h has nonremovable discontinuities at $x = \pm 1, \pm 2, \pm 3, \dots$

120. (a) Define $f(x) = f_2(x) - f_1(x)$. Because f_1 and f_2 are continuous on $[a, b]$, so is f .

$$f(a) = f_2(a) - f_1(a) > 0 \text{ and}$$

$$f(b) = f_2(b) - f_1(b) < 0$$

By the Intermediate Value Theorem, there exists c in $[a, b]$ such that $f(c) = 0$.

$$f(c) = f_2(c) - f_1(c) = 0 \Rightarrow f_1(c) = f_2(c)$$

(b) Let $f_1(x) = x$ and $f_2(x) = \cos x$, continuous on $[0, \pi/2]$, $f_1(0) < f_2(0)$ and $f_1(\pi/2) > f_2(\pi/2)$.

So by part (a), there exists c in $[0, \pi/2]$ such that $c = \cos(c)$.

Using a graphing utility, $c \approx 0.739$.

116. Let y be a real number. If $y = 0$, then $x = 0$. If $y > 0$, then let $0 < x_0 < \pi/2$ such that

$M = \tan x_0 > y$ (this is possible since the tangent function increases without bound on $[0, \pi/2)$). By the Intermediate Value Theorem, $f(x) = \tan x$ is continuous on $[0, x_0]$ and $0 < y < M$, which implies that there exists x between 0 and x_0 such that $\tan x = y$. The argument is similar if $y < 0$.

121. The statement is true.

If $y \geq 0$ and $y \leq 1$, then $y(y-1) \leq 0 \leq x^2$, as desired. So assume $y > 1$. There are now two cases.

Case 1: If $x \leq y - \frac{1}{2}$, then $2x + 1 \leq 2y$ and

$$\begin{aligned} y(y-1) &= y(y+1) - 2y \\ &\leq (x+1)^2 - 2y \\ &= x^2 + 2x + 1 - 2y \\ &\leq x^2 + 2y - 2y \\ &= x^2 \end{aligned}$$

Case 2: If $x \geq y - \frac{1}{2}$

$$\begin{aligned} x^2 &\geq \left(y - \frac{1}{2}\right)^2 \\ &= y^2 - y + \frac{1}{4} \\ &> y^2 - y \\ &= y(y-1) \end{aligned}$$

In both cases, $y(y-1) \leq x^2$.

122. $P(1) = P(0^2 + 1) = P(0)^2 + 1 = 1$

$$P(2) = P(1^2 + 1) = P(1)^2 + 1 = 2$$

$$P(5) = P(2^2 + 1) = P(2)^2 + 1 = 5$$

Continuing this pattern, you see that $P(x) = x$ for infinitely many values of x . So, the finite degree polynomial must be constant: $P(x) = x$ for all x .

Section 1.5 Infinite Limits

1. $f(x) = \frac{1}{x - 4}$

As x approaches 4 from the left, $x - 4$ is a small negative number. So,

$$\lim_{x \rightarrow 4^-} f(x) = -\infty$$

As x approaches 4 from the right, $x - 4$ is a small positive number. So,

$$\lim_{x \rightarrow 4^+} f(x) = \infty$$

2. $f(x) = \frac{-1}{x - 4}$

As x approaches 4 from the left, $x - 4$ is a small negative number. So,

$$\lim_{x \rightarrow 4^-} f(x) = \infty$$

As x approaches 4 from the right, $x - 4$ is a small positive number. So,

$$\lim_{x \rightarrow 4^+} f(x) = -\infty$$

3. $f(x) = \frac{1}{(x - 4)^2}$

As x approaches 4 from the left or right, $(x - 4)^2$ is a small positive number. So,


$$\lim_{x \rightarrow 4^+} f(x) = \lim_{x \rightarrow 4^-} f(x) = \infty$$

9. $f(x) = \frac{1}{x^2 - 9}$

x	-3.5	-3.1	-3.01	-3.001	-2.999	-2.99	-2.9	-2.5
$f(x)$	0.308	1.639	16.64	166.6	-166.7	-16.69	-1.695	-0.364

$$\lim_{x \rightarrow -3^-} f(x) = \infty$$

$$\lim_{x \rightarrow -3^+} f(x) = -\infty$$




10. $f(x) = \frac{x}{x^2 - 9}$

x	-3.5	-3.1	-3.01	-3.001	-2.999	-2.99	-2.9	-2.5
$f(x)$	-1.077	-5.082	-50.08	-500.1	499.9	49.92	4.915	0.9091

$$\lim_{x \rightarrow -3^-} f(x) = -\infty$$

$$\lim_{x \rightarrow -3^+} f(x) = \infty$$



4. $f(x) = \frac{-1}{(x - 4)^2}$

As x approaches 4 from the left or right, $(x - 4)^2$ is a small positive number. So,

$$\lim_{x \rightarrow 4^-} f(x) = \lim_{x \rightarrow 4^+} f(x) = -\infty$$

5. $\lim_{x \rightarrow -2^+} 2 \left| \frac{x}{x^2 - 4} \right| = \infty$

$$\lim_{x \rightarrow -2^-} 2 \left| \frac{x}{x^2 - 4} \right| = \infty$$

6. $\lim_{x \rightarrow -2^+} \frac{1}{x + 2} = \infty$

$$\lim_{x \rightarrow -2^-} \frac{1}{x + 2} = -\infty$$

7. $\lim_{x \rightarrow -2^+} \tan \frac{\pi x}{4} = -\infty$

$$\lim_{x \rightarrow -2^-} \tan \frac{\pi x}{4} = \infty$$

8. $\lim_{x \rightarrow -2^+} \sec \frac{\pi x}{4} = \infty$


$$\lim_{x \rightarrow -2^-} \sec \frac{\pi x}{4} = -\infty$$

11. $f(x) = \frac{x^2}{x^2 - 9}$

x	-3.5	-3.1	-3.01	-3.001	-2.999	-2.99	-2.9	-2.5
$f(x)$	3.769	15.75	150.8	1501	-1499	-149.3	-14.25	-2.273

$\lim_{x \rightarrow -3^-} f(x) = \infty$

$\lim_{x \rightarrow -3^+} f(x) = -\infty$




12. $f(x) = \sec \frac{\pi x}{6}$

x	-3.5	-3.1	-3.01	-3.001	-2.999	-2.99	-2.9	-2.5
$f(x)$	-3.864	-19.11	-191.0	-1910	1910	191.0	19.11	3.864

$\lim_{x \rightarrow -3^-} f(x) = -\infty$

$\lim_{x \rightarrow -3^+} f(x) = \infty$



13. $\lim_{x \rightarrow 0^+} \frac{1}{x^2} = \infty = \lim_{x \rightarrow 0^-} \frac{1}{x^2}$

Therefore, $x = 0$ is a vertical asymptote.

14. $\lim_{x \rightarrow 2^+} \frac{4}{(x-2)^3} = \infty$

$\lim_{x \rightarrow 2^-} \frac{4}{(x-2)^3} = -\infty$

Therefore, $x = 2$ is a vertical asymptote.

15. $\lim_{x \rightarrow -2^-} \frac{x^2}{x^2 - 4} = \infty$ and $\lim_{x \rightarrow -2^+} \frac{x^2}{x^2 - 4} = -\infty$

Therefore, $x = -2$ is a vertical asymptote.

$\lim_{x \rightarrow 2^-} \frac{x^2}{x^2 - 4} = -\infty$ and $\lim_{x \rightarrow 2^+} \frac{x^2}{x^2 - 4} = \infty$

Therefore, $x = 2$ is a vertical asymptote.

16. No vertical asymptote because the denominator is never zero.

17. No vertical asymptote because the denominator is never zero.

18. $\lim_{s \rightarrow -5^-} h(s) = -\infty$ and $\lim_{s \rightarrow -5^+} h(s) = \infty$.

Therefore, $s = -5$ is a vertical asymptote.

$\lim_{s \rightarrow 5^-} h(s) = -\infty$ and $\lim_{s \rightarrow 5^+} h(s) = \infty$.

Therefore, $s = 5$ is a vertical asymptote.

19. $\lim_{x \rightarrow 2^+} \frac{x^2 - 2}{(x-2)(x+1)} = \infty$

$\lim_{x \rightarrow 2^-} \frac{x^2 - 2}{(x-2)(x+1)} = -\infty$

Therefore, $x = 2$ is a vertical asymptote.

$\lim_{x \rightarrow -1^+} \frac{x^2 - 2}{(x-2)(x+1)} = \infty$

$\lim_{x \rightarrow -1^-} \frac{x^2 - 2}{(x-2)(x+1)} = -\infty$

Therefore, $x = -1$ is a vertical asymptote.

$$20. \lim_{x \rightarrow 0^-} \frac{2+x}{x^2(1-x)} = \lim_{x \rightarrow 0^+} \frac{2+x}{x^2(1-x)} = \infty$$

Therefore, $x = 0$ is a vertical asymptote.

$$\lim_{x \rightarrow 1^-} \frac{2+x}{x^2(1-x)} = \infty$$

$$\lim_{x \rightarrow 1^+} \frac{2+x}{x^2(1-x)} = -\infty$$

Therefore, $x = 1$ is a vertical asymptote.

$$21. \lim_{t \rightarrow 0^+} \left(1 - \frac{4}{t^2}\right) = -\infty = \lim_{t \rightarrow 0^-} \left(1 - \frac{4}{t^2}\right)$$

Therefore, $t = 0$ is a vertical asymptote.

$$22. g(x) = \frac{(1/2)x^3 - x^2 - 4x}{3x^2 - 6x - 24} = \frac{1}{6} \frac{x(x^2 - 2x - 8)}{x^2 - 2x - 8}$$

$$= \frac{1}{6}x, \quad x \neq -2, 4$$

No vertical asymptote. The graph has holes at $x = -2$ and $x = 4$.

$$23. f(x) = \frac{3}{x^2 + x - 2} = \frac{3}{(x+2)(x-1)}$$

Vertical asymptotes at $x = -2$ and $x = 1$.

$$24. f(x) = \frac{4(x^2 + x - 6)}{x(x^3 - 2x^2 - 9x + 18)}$$

$$= \frac{4(x+3)(x-2)}{x(x-2)(x^2-9)}$$

$$= \frac{4}{x(x-3)}, \quad x \neq -3, 2$$

Vertical asymptotes at $x = 0$ and $x = 3$. The graph has holes at $x = -3$ and $x = 2$.

$$25. f(x) = \frac{x^3 + 1}{x + 1} = \frac{(x+1)(x^2 - x + 1)}{x + 1}$$

has no vertical asymptote because

$$\lim_{x \rightarrow -1} f(x) = \lim_{x \rightarrow -1} (x^2 - x + 1) = 3.$$

The graph has a hole at $x = -1$.

$$26. h(x) = \frac{x^2 - 4}{x^3 + 2x^2 + x + 2} = \frac{(x+2)(x-2)}{(x+2)(x^2+1)}$$

has no vertical asymptote because

$$\lim_{x \rightarrow -2} h(x) = \lim_{x \rightarrow -2} \frac{x-2}{x^2+1} = -\frac{4}{5}.$$

The graph has a hole at $x = -2$.

$$27. f(x) = \frac{(x-5)(x+3)}{(x-5)(x^2+1)} = \frac{x+3}{x^2+1}, \quad x \neq 5$$

No vertical asymptote. The graph has a hole at $x = 5$.

$$28. h(t) = \frac{t(t-2)}{(t-2)(t+2)(t^2+4)}$$

$$= \frac{t}{(t+2)(t^2+4)}, \quad t \neq 2$$

Vertical asymptote at $t = -2$. The graph has a hole at $t = 2$.

$$29. f(x) = \tan \pi x = \frac{\sin \pi x}{\cos \pi x} \text{ has vertical asymptotes at}$$

$$x = \frac{2n+1}{2}, \quad n \text{ any integer.}$$

$$30. f(x) = \sec \pi x = \frac{1}{\cos \pi x} \text{ has vertical asymptotes at}$$

$$x = \frac{2n+1}{2}, \quad n \text{ any integer.}$$

$$31. s(t) = \frac{t}{\sin t} \text{ has vertical asymptotes at } t = n\pi, \quad n \text{ a}$$

nonzero integer. There is no vertical asymptote at $t = 0$ since

$$\lim_{t \rightarrow 0} \frac{t}{\sin t} = 1.$$


$$32. g(\theta) = \frac{\tan \theta}{\theta} = \frac{\sin \theta}{\theta \cos \theta} \text{ has vertical asymptotes at}$$

$$\theta = \frac{(2n+1)\pi}{2} = \frac{\pi}{2} + n\pi, \quad n \text{ any integer.}$$

There is no vertical asymptote at $\theta = 0$ because


$$\lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1.$$

$$33. \lim_{x \rightarrow -1} \frac{x^2 - 1}{x + 1} = \lim_{x \rightarrow -1} (x - 1) = -2$$



Removable discontinuity at $x = -1$

34. $\lim_{x \rightarrow -1} \frac{x^2 - 6x - 7}{x + 1} = \lim_{x \rightarrow -1} (x - 7) = -8$




Removable discontinuity at $x = -1$

35. $\lim_{x \rightarrow -1^+} \frac{x^2 + 1}{x + 1} = \infty$


$\lim_{x \rightarrow -1^-} \frac{x^2 + 1}{x + 1} = -\infty$

Vertical asymptote at $x = -1$



36. $\lim_{x \rightarrow -1} \frac{\sin(x + 1)}{x + 1} = 1$

Removable discontinuity at $x = -1$



37. $\lim_{x \rightarrow -1^+} \frac{1}{x + 1} = \infty$

38. $\lim_{x \rightarrow 1^-} \frac{-1}{(x - 1)^2} = -\infty$

39. $\lim_{x \rightarrow 2^+} \frac{x}{x - 2} = \infty$

40. $\lim_{x \rightarrow 1^+} \frac{2 + x}{1 - x} = -\infty$

41. $\lim_{x \rightarrow 1^+} \frac{x^2}{(x - 1)^2} = \infty$

42. $\lim_{x \rightarrow 4^-} \frac{x^2}{x^2 + 16} = \frac{1}{2}$

43. $\lim_{x \rightarrow 3^-} \frac{x + 3}{(x^2 + x - 6)} = \lim_{x \rightarrow 3^-} \frac{x + 3}{(x + 3)(x - 2)}$
 $= \lim_{x \rightarrow 3^-} \frac{1}{x - 2} = -\frac{1}{5}$

44. $\lim_{x \rightarrow -(1/2)^+} \frac{6x^2 + x - 1}{4x^2 - 4x - 3} = \lim_{x \rightarrow -(1/2)^+} \frac{(3x - 1)(2x + 1)}{(2x - 3)(2x + 1)}$
 $= \lim_{x \rightarrow -(1/2)^+} \frac{3x - 1}{2x - 3} = \frac{5}{8}$

45. $\lim_{x \rightarrow 1} \frac{x - 1}{(x^2 + 1)(x - 1)} = \lim_{x \rightarrow 1} \frac{1}{x^2 + 1} = \frac{1}{2}$

46. $\lim_{x \rightarrow 3} \frac{x - 2}{x^2} = \frac{1}{9}$

47. $\lim_{x \rightarrow 0^-} \left(1 + \frac{1}{x}\right) = -\infty$

48. $\lim_{x \rightarrow 0^-} \left(x^2 - \frac{1}{x}\right) = \infty$

49. $\lim_{x \rightarrow 0^+} \frac{2}{\sin x} = \infty$

50. $\lim_{x \rightarrow (\pi/2)^+} \frac{-2}{\cos x} = \infty$

51. $\lim_{x \rightarrow \pi} \frac{\sqrt{x}}{\csc x} = \lim_{x \rightarrow \pi} (\sqrt{x} \sin x) = 0$


52. $\lim_{x \rightarrow 0} \frac{(x + 2)}{\cot x} = \lim_{x \rightarrow 0} [(x + 2)\tan x] = 0$

53. $\lim_{x \rightarrow (1/2)^-} x \sec(\pi x) = \infty$ and $\lim_{x \rightarrow (1/2)^+} x \sec(\pi x) = -\infty$.
 Therefore, $\lim_{x \rightarrow (1/2)} x \sec(\pi x)$ does not exist.


54. $\lim_{x \rightarrow (1/2)^-} x^2 \tan \pi x = \infty$ and
 $\lim_{x \rightarrow (1/2)^+} x^2 \tan \pi x = -\infty$. Therefore,
 $\lim_{x \rightarrow (1/2)} x^2 \tan \pi x$ does not exist.

55. $f(x) = \frac{x^2 + x + 1}{x^3 - 1} = \frac{x^2 + x + 1}{(x - 1)(x^2 + x + 1)}$


$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} \frac{1}{x - 1} = \infty$




56. $f(x) = \frac{x^3 - 1}{x^2 + x + 1} = \frac{(x - 1)(x^2 + x + 1)}{x^2 + x + 1}$
 $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x - 1) = 0$



57. $f(x) = \frac{1}{x^2 - 25}$
 $\lim_{x \rightarrow 5^-} f(x) = -\infty$



58. $f(x) = \sec \frac{\pi x}{8}$
 $\lim_{x \rightarrow 4^+} f(x) = -\infty$



59. A limit in which $f(x)$ increases or decreases without bound as x approaches c is called an infinite limit. ∞ is not a number. Rather, the symbol

$$\lim_{x \rightarrow c} f(x) = \infty$$


says how the limit fails to exist.

60. The line $x = c$ is a vertical asymptote if the graph of f approaches $\pm\infty$ as x approaches c .

61. One answer is

$$f(x) = \frac{x - 3}{(x - 6)(x + 2)} = \frac{x - 3}{x^2 - 4x - 12}$$

62. No. For example, $f(x) = \frac{1}{x^2 + 1}$ has no vertical asymptote.



64. No, it is not true. Consider $p(x) = x^2 - 1$. The function

$$f(x) = \frac{x^2 - 1}{x - 1} = \frac{p(x)}{x - 1}$$

has a hole at $(1, 2)$, not a vertical asymptote.

65. $m = \frac{m_0}{\sqrt{1 - (v^2/c^2)}}$

$$\lim_{v \rightarrow c^-} m = \lim_{v \rightarrow c^-} \frac{m_0}{\sqrt{1 - (v^2/c^2)}} = \infty$$

66. $P = \frac{k}{V}$

$$\lim_{V \rightarrow 0^+} \frac{k}{V} = k(\infty) = \infty$$

(In this case you know that $k > 0$.)

67. (a) $r = 50\pi \sec^2 \frac{\pi}{6} = \frac{200\pi}{3}$ ft/sec

(b) $r = 50\pi \sec^2 \frac{\pi}{3} = 200\pi$ ft/sec

(c) $\lim_{\theta \rightarrow (\pi/2)^-} [50\pi \sec^2 \theta] = \infty$

68. (a) $r = \frac{2(7)}{\sqrt{625 - 49}} = \frac{7}{12}$ ft/sec

(b) $r = \frac{2(15)}{\sqrt{625 - 225}} = \frac{3}{2}$ ft/sec

(c) $\lim_{x \rightarrow 25^-} \frac{2x}{\sqrt{625 - x^2}} = \infty$

69. (a) Average speed = $\frac{\text{Total distance}}{\text{Total time}}$

$$50 = \frac{2d}{(d/x) + (d/y)}$$

$$50 = \frac{2xy}{y + x}$$

$$50y + 50x = 2xy$$

$$50x = 2xy - 50y$$

$$50x = 2y(x - 25)$$

$$\frac{25x}{x - 25} = y$$

Domain: $x > 25$

(b)


x	30	40	50	60
y	150	66.667	50	42.857

(c) $\lim_{x \rightarrow 25^+} \frac{25x}{\sqrt{x - 25}} = \infty$

As x gets close to 25 mi/h, y becomes larger and larger.

70. (a)


x	1	0.5	0.2	0.1	0.01	0.001	0.0001
$f(x)$	0.1585	0.0411	0.0067	0.0017	≈ 0	≈ 0	≈ 0



$$\lim_{x \rightarrow 0^+} \frac{x - \sin x}{x} = 0$$

(b)


x	1	0.5	0.2	0.1	0.01	0.001	0.0001
$f(x)$	0.1585	0.0823	0.0333	0.0167	0.0017	≈ 0	≈ 0



$$\lim_{x \rightarrow 0^+} \frac{x - \sin x}{x^2} = 0$$

(c)


x	1	0.5	0.2	0.1	0.01	0.001	0.0001
$f(x)$	0.1585	0.1646	0.1663	0.1666	0.1667	0.1667	0.1667



$$\lim_{x \rightarrow 0^+} \frac{x - \sin x}{x^3} = 0.1667 \text{ (1/6)}$$

(d)

x	1	0.5	0.2	0.1	0.01	0.001	0.0001
$f(x)$	0.1585	0.3292	0.8317	1.6658	16.67	166.7	1667.0



$$\lim_{x \rightarrow 0^+} \frac{x - \sin x}{x^4} = \infty$$


$$\text{For } n > 3, \lim_{x \rightarrow 0^+} \frac{x - \sin x}{x^n} = \infty.$$

71. (a) $A = \frac{1}{2}bh - \frac{1}{2}r^2\theta = \frac{1}{2}(10)(10 \tan \theta) - \frac{1}{2}(10)^2\theta = 50 \tan \theta - 50 \theta$

Domain: $\left(0, \frac{\pi}{2}\right)$

(b)

θ	0.3	0.6	0.9	1.2	1.5
$f(\theta)$	0.47	4.21	18.0	68.6	630.1



(c) $\lim_{\theta \rightarrow \pi/2^-} A = \infty$

72. (a) Because the circumference of the motor is half that of the saw arbor, the saw makes $1700/2 = 850$ revolutions per minute.

(b) The direction of rotation is reversed.

(c) $2(20 \cot \phi) + 2(10 \cot \phi)$: straight sections. The angle subtended in each circle is $2\pi - \left(2\left(\frac{\pi}{2} - \phi\right)\right) = \pi + 2\phi$.


So, the length of the belt around the pulleys is $20(\pi + 2\phi) + 10(\pi + 2\phi) = 30(\pi + 2\phi)$.

Total length = $60 \cot \phi + 30(\pi + 2\phi)$

Domain: $\left(0, \frac{\pi}{2}\right)$

(d)

ϕ	0.3	0.6	0.9	1.2	1.5
L	306.2	217.9	195.9	189.6	188.5



(f) $\lim_{\phi \rightarrow (\pi/2)^-} L = 60\pi \approx 188.5$

(All the belts are around pulleys.)

(g) $\lim_{\phi \rightarrow 0^+} L = \infty$

73. False. For instance, let

$$f(x) = \frac{x^2 - 1}{x - 1} \text{ or}$$

$$g(x) = \frac{x}{x^2 + 1}.$$

74. True

75. False. The graphs of

$$y = \tan x, y = \cot x, y = \sec x \text{ and } y = \csc x \text{ have}$$

vertical asymptotes.

76. False. Let

$$f(x) = \begin{cases} \frac{1}{x}, & x \neq 0 \\ 3, & x = 0. \end{cases}$$

The graph of f has a vertical asymptote at $x = 0$, but $f(0) = 3$.

77. Let $f(x) = \frac{1}{x^2}$ and $g(x) = \frac{1}{x^4}$, and $c = 0$.

$$\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty \text{ and } \lim_{x \rightarrow 0} \frac{1}{x^4} = \infty, \text{ but}$$

$$\lim_{x \rightarrow 0} \left(\frac{1}{x^2} - \frac{1}{x^4} \right) = \lim_{x \rightarrow 0} \left(\frac{x^2 - 1}{x^4} \right) = -\infty \neq 0.$$

78. Given $\lim_{x \rightarrow c} f(x) = \infty$ and $\lim_{x \rightarrow c} g(x) = L$:

(2) Product:

If $L > 0$, then for $\varepsilon = L/2 > 0$ there exists

$\delta_1 > 0$ such that $|g(x) - L| < L/2$ whenever

$0 < |x - c| < \delta_1$. So, $L/2 < g(x) < 3L/2$. Because

$\lim_{x \rightarrow c} f(x) = \infty$ then for $M > 0$, there exists

$\delta_2 > 0$ such that $f(x) > M(2/L)$ whenever

$|x - c| < \delta_2$. Let δ be the smaller of δ_1 and

δ_2 . Then for $0 < |x - c| < \delta$, you have

$f(x)g(x) > M(2/L)(L/2) = M$. Therefore

$\lim_{x \rightarrow c} f(x)g(x) = \infty$. The proof is similar for $L < 0$.

(3) Quotient: Let $\varepsilon > 0$ be given.

There exists $\delta_1 > 0$ such that

$f(x) > 3L/2\varepsilon$ whenever $0 < |x - c| < \delta_1$ and

there exists $\delta_2 > 0$ such that

$|g(x) - L| < L/2$ whenever $0 < |x - c| < \delta_2$. This

inequality gives us $L/2 < g(x) < 3L/2$. Let δ be

the smaller of δ_1 and δ_2 . Then for

$0 < |x - c| < \delta$, you have

$$\left| \frac{g(x)}{f(x)} \right| < \frac{3L/2}{3L/2\varepsilon} = \varepsilon.$$

Therefore, $\lim_{x \rightarrow c} \frac{g(x)}{f(x)} = 0$.

82. $f(x) = \frac{1}{x-5}$ is defined for all $x < 5$. Let $N < 0$ be given. You need $\delta > 0$ such that $f(x) = \frac{1}{x-5} < N$ whenever

$5 - \delta < x < 5$. Equivalently, $x - 5 > \frac{1}{N}$ whenever $|x - 5| < \delta, x < 5$. Equivalently, $\frac{1}{|x-5|} < -\frac{1}{N}$ whenever

$|x - 5| < \delta, x < 5$. So take $\delta = -\frac{1}{N}$. Note that $\delta > 0$ because $N < 0$. For $|x - 5| < \delta$ and

$x < 5, \frac{1}{|x-5|} > \frac{1}{\delta} = -N$, and $\frac{1}{x-5} = -\frac{1}{|x-5|} < N$.

79. Given $\lim_{x \rightarrow c} f(x) = \infty$, let $g(x) = 1$. Then

$$\lim_{x \rightarrow c} \frac{g(x)}{f(x)} = 0 \text{ by Theorem 1.15.}$$

80. Given $\lim_{x \rightarrow c} \frac{1}{f(x)} = 0$. Suppose $\lim_{x \rightarrow c} f(x)$ exists and equals L .

$$\text{Then, } \lim_{x \rightarrow c} \frac{1}{f(x)} = \frac{\lim_{x \rightarrow c} 1}{\lim_{x \rightarrow c} f(x)} = \frac{1}{L} = 0.$$

This is not possible. So, $\lim_{x \rightarrow c} f(x)$ does not exist.

81. $f(x) = \frac{1}{x-3}$ is defined for all $x > 3$. Let $M > 0$ be given. You need $\delta > 0$ such that

$$f(x) = \frac{1}{x-3} > M \text{ whenever } 3 < x < 3 + \delta.$$

Equivalently, $x - 3 < \frac{1}{M}$ whenever


$$|x - 3| < \delta, x > 3.$$

So take $\delta = \frac{1}{M}$. Then for $x > 3$ and

$$|x - 3| < \delta, \frac{1}{x-3} > \frac{1}{\delta} = M \text{ and so } f(x) > M.$$

Review Exercises for Chapter 1

1. Calculus required. Using a graphing utility, you can estimate the length to be 8.3. Or, the length is slightly longer than the distance between the two points, approximately 8.25.




2. Precalculus. $L = \sqrt{(9 - 1)^2 + (3 - 1)^2} \approx 8.25$

3. $f(x) = \frac{4}{x + 2} - 2$

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	-1.0526	-1.0050	-1.0005	-0.9995	-0.9950	-0.9524


$\lim_{x \rightarrow 0} f(x) \approx -1.0$



4.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	1.432	1.416	1.414	1.414	1.413	1.397

$\lim_{x \rightarrow 0} f(x) \approx 1.414$



5. $\lim_{x \rightarrow 1} (x + 4) = 1 + 4 = 5$

Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon$. Then for $0 < |x - 1| < \delta = \varepsilon$, you have

$$|x - 1| < \varepsilon$$

$$|(x + 4) - 5| < \varepsilon$$

$$|f(x) - L| < \varepsilon.$$

6. $\lim_{x \rightarrow 9} \sqrt{x} = \sqrt{9} = 3$

Let $\varepsilon > 0$ be given. You need

$$|\sqrt{x} - 3| < \varepsilon \Rightarrow |\sqrt{x} + 3| |\sqrt{x} - 3| < \varepsilon |\sqrt{x} + 3| \Rightarrow |x - 9| < \varepsilon |\sqrt{x} + 3|.$$

Assuming $4 < x < 16$, you can choose $\delta = 5\varepsilon$.

So, for $0 < |x - 9| < \delta = 5\varepsilon$, you have

$$\begin{aligned} |x - 9| < 5\varepsilon &< |\sqrt{x} + 3| \varepsilon \\ |\sqrt{x} - 3| &< \varepsilon \\ |f(x) - L| &< \varepsilon. \end{aligned}$$

7. $\lim_{x \rightarrow 2} (1 - x^2) = 1 - 2^2 = -3$

Let $\varepsilon > 0$ be given. You need

$$|1 - x^2 - (-3)| < \varepsilon \Rightarrow |x^2 - 4| = |x - 2| |x + 2| < \varepsilon \Rightarrow |x - 2| < \frac{1}{|x + 2|} \varepsilon$$

Assuming $1 < x < 3$, you can choose $\delta = \frac{\varepsilon}{5}$.

So, for $0 < |x - 2| < \delta = \frac{\varepsilon}{5}$, you have

$$\begin{aligned} |x - 2| < \frac{\varepsilon}{5} &< \frac{\varepsilon}{|x + 2|} \\ |x - 2| |x + 2| &< \varepsilon \\ |x^2 - 4| &< \varepsilon \\ |4 - x^2| &< \varepsilon \\ |(1 - x^2) - (-3)| &< \varepsilon \\ |f(x) - L| &< \varepsilon. \end{aligned}$$

8. $\lim_{x \rightarrow 5} 9 = 9$. Let $\varepsilon > 0$ be given. δ can be any positive number. So, for $0 < |x - 5| < \delta$, you have

$$\begin{aligned} |9 - 9| &< \varepsilon \\ |f(x) - L| &< \varepsilon. \end{aligned}$$

9. $h(x) = \frac{4x - x^2}{x} = \frac{x(4 - x)}{x} = 4 - x, x \neq 0$

(a) $\lim_{x \rightarrow 0} h(x) = 4 - 0 = 4$

(b) $\lim_{x \rightarrow -1} h(x) = 4 - (-1) = 5$

10. $g(x) = \frac{-2x}{x - 3}$

(a) $\lim_{x \rightarrow 3} g(x)$ does not exist

(b) $\lim_{x \rightarrow 0} g(x) = \frac{-2(0)}{0 - 3} = 0$

11. $\lim_{x \rightarrow 6} (x - 2)^2 = (6 - 2)^2 = 16$

12. $\lim_{x \rightarrow 7} (10 - x)^4 = (10 - 7)^4 = 3^4 = 81$

13. $\lim_{t \rightarrow 4} \sqrt{t + 2} = \sqrt{4 + 2} = \sqrt{6} = 2.45$

14. $\lim_{y \rightarrow 4} 3|y - 1| = 3|4 - 1| = 9$

15. $\lim_{t \rightarrow -2} \frac{t + 2}{t^2 - 4} = \lim_{t \rightarrow -2} \frac{1}{t - 2} = -\frac{1}{4}$

16. $\lim_{t \rightarrow 3} \frac{t^2 - 9}{t - 3} = \lim_{t \rightarrow 3} (t + 3) = 6$

$$\begin{aligned}
 17. \lim_{x \rightarrow 4} \frac{\sqrt{x-3} - 1}{x-4} &= \lim_{x \rightarrow 4} \frac{\sqrt{x-3} - 1}{x-4} \cdot \frac{\sqrt{x-3} + 1}{\sqrt{x-3} + 1} \\
 &= \lim_{x \rightarrow 4} \frac{(x-3) - 1}{(x-4)(\sqrt{x-3} + 1)} \\
 &= \lim_{x \rightarrow 4} \frac{1}{\sqrt{x-3} + 1} = \frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 19. \lim_{x \rightarrow 0} \frac{[1/(x+1)] - 1}{x} &= \lim_{x \rightarrow 0} \frac{1 - (x+1)}{x(x+1)} \\
 &= \lim_{x \rightarrow 0} \frac{-1}{x+1} = -1
 \end{aligned}$$

$$\begin{aligned}
 18. \lim_{x \rightarrow 0} \frac{\sqrt{4+x} - 2}{x} &= \lim_{x \rightarrow 0} \frac{\sqrt{4+x} - 2}{x} \cdot \frac{\sqrt{4+x} + 2}{\sqrt{4+x} + 2} \\
 &= \lim_{x \rightarrow 0} \frac{1}{\sqrt{4+x} + 2} = \frac{1}{4}
 \end{aligned}$$

$$\begin{aligned}
 20. \lim_{s \rightarrow 0} \frac{(1/\sqrt{1+s}) - 1}{s} &= \lim_{s \rightarrow 0} \left[\frac{(1/\sqrt{1+s}) - 1}{s} \cdot \frac{(1/\sqrt{1+s}) + 1}{(1/\sqrt{1+s}) + 1} \right] \\
 &= \lim_{s \rightarrow 0} \frac{[1/(1+s)] - 1}{s[(1/\sqrt{1+s}) + 1]} = \lim_{s \rightarrow 0} \frac{-1}{(1+s)[(1/\sqrt{1+s}) + 1]} = -\frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
 21. \lim_{x \rightarrow -5} \frac{x^3 + 125}{x + 5} &= \lim_{x \rightarrow -5} \frac{(x+5)(x^2 - 5x + 25)}{x + 5} \\
 &= \lim_{x \rightarrow -5} (x^2 - 5x + 25) = 75
 \end{aligned}$$

$$\begin{aligned}
 22. \lim_{x \rightarrow -2} \frac{x^2 - 4}{x^3 + 8} &= \lim_{x \rightarrow -2} \frac{(x+2)(x-2)}{(x+2)(x^2 - 2x + 4)} \\
 &= \lim_{x \rightarrow -2} \frac{x-2}{x^2 - 2x + 4} = -\frac{4}{12} = -\frac{1}{3}
 \end{aligned}$$

$$23. \lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x} = \lim_{x \rightarrow 0} \left(\frac{x}{\sin x} \right) \left(\frac{1 - \cos x}{x} \right) = (1)(0) = 0$$

$$24. \lim_{x \rightarrow (\pi/4)} \frac{4x}{\tan x} = \frac{4(\pi/4)}{1} = \pi$$

$$\begin{aligned}
 25. \lim_{\Delta x \rightarrow 0} \frac{\sin[(\pi/6) + \Delta x] - (1/2)}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{\sin(\pi/6)\cos \Delta x + \cos(\pi/6)\sin \Delta x - (1/2)}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{1}{2} \cdot \frac{(\cos \Delta x - 1)}{\Delta x} + \lim_{\Delta x \rightarrow 0} \frac{\sqrt{3}}{2} \cdot \frac{\sin \Delta x}{\Delta x} = 0 + \frac{\sqrt{3}}{2}(1) = \frac{\sqrt{3}}{2}
 \end{aligned}$$

$$\begin{aligned}
 26. \lim_{\Delta x \rightarrow 0} \frac{\cos(\pi + \Delta x) + 1}{\Delta x} &= \lim_{\Delta x \rightarrow 0} \frac{\cos \pi \cos \Delta x - \sin \pi \sin \Delta x + 1}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \left[-\frac{(\cos \Delta x - 1)}{\Delta x} \right] - \lim_{\Delta x \rightarrow 0} \left[\sin \pi \frac{\sin \Delta x}{\Delta x} \right] \\
 &= -0 - (0)(1) = 0
 \end{aligned}$$

$$27. \lim_{x \rightarrow c} [f(x) \cdot g(x)] = \left(-\frac{3}{4}\right)\left(\frac{2}{3}\right) = -\frac{1}{2}$$

$$29. \lim_{x \rightarrow c} [f(x) + 2g(x)] = -\frac{3}{4} + 2\left(\frac{2}{3}\right) = \frac{7}{12}$$

$$28. \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow c} f(x)}{\lim_{x \rightarrow c} g(x)} = \frac{-3/4}{2/3} = -\frac{9}{8}$$

$$30. \lim_{x \rightarrow c} [f(x)]^2 = \left[\lim_{x \rightarrow c} f(x) \right]^2 = \left(-\frac{3}{4}\right)^2 = \frac{9}{16}$$

31. $f(x) = \frac{\sqrt{2x+1} - \sqrt{3}}{x-1}$

(a)

x	1.1	1.01	1.001	1.0001
$f(x)$	0.5680	0.5764	0.5773	0.5773

$$\lim_{x \rightarrow 1^+} \frac{\sqrt{2x+1} - \sqrt{3}}{x-1} \approx 0.577 \quad (\text{Actual limit is } \sqrt{3}/3.)$$

(b)  The graph has a hole at $x = 1$.

$$\lim_{x \rightarrow 1^+} f(x) \approx 0.5774.$$

(c)

$$\begin{aligned} \lim_{x \rightarrow 1^+} \frac{\sqrt{2x+1} - \sqrt{3}}{x-1} &= \lim_{x \rightarrow 1^+} \frac{\sqrt{2x+1} - \sqrt{3}}{x-1} \cdot \frac{\sqrt{2x+1} + \sqrt{3}}{\sqrt{2x+1} + \sqrt{3}} \\ &= \lim_{x \rightarrow 1^+} \frac{(2x+1) - 3}{(x-1)(\sqrt{2x+1} + \sqrt{3})} \\ &= \lim_{x \rightarrow 1^+} \frac{2}{\sqrt{2x+1} + \sqrt{3}} \\ &= \frac{2}{2\sqrt{3}} = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3} \end{aligned}$$

32. $f(x) = \frac{1 - \sqrt[3]{x}}{x-1}$

(a)

x	1.1	1.01	1.001	1.0001
$f(x)$	-0.3228	-0.3322	-0.3332	-0.3333

$$\lim_{x \rightarrow 1^+} \frac{1 - \sqrt[3]{x}}{x-1} \approx -0.333 \quad (\text{Actual limit is } -\frac{1}{3}.)$$

(b)  The graph has a hole at $x = 1$.

$$\lim_{x \rightarrow 1^+} f(x) \approx 0.333.$$

(c)

$$\begin{aligned} \lim_{x \rightarrow 1^+} \frac{1 - \sqrt[3]{x}}{x-1} &= \lim_{x \rightarrow 1^+} \frac{1 - \sqrt[3]{x}}{x-1} \cdot \frac{1 + \sqrt[3]{x} + (\sqrt[3]{x})^2}{1 + \sqrt[3]{x} + (\sqrt[3]{x})^2} \\ &= \lim_{x \rightarrow 1^+} \frac{1-x}{(x-1)[1 + \sqrt[3]{x} + (\sqrt[3]{x})^2]} \\ &= \lim_{x \rightarrow 1^+} \frac{-1}{1 + \sqrt[3]{x} + (\sqrt[3]{x})^2} = -\frac{1}{3} \end{aligned}$$

$$\begin{aligned}
 33. \quad v &= \lim_{t \rightarrow 4} \frac{s(4) - s(t)}{4 - t} \\
 &= \lim_{t \rightarrow 4} \frac{[-4.9(16) + 250] - [-4.9t^2 + 250]}{4 - t} \\
 &= \lim_{t \rightarrow 4} \frac{4.9(t^2 - 16)}{4 - t} \\
 &= \lim_{t \rightarrow 4} \frac{4.9(t - 4)(t + 4)}{4 - t} \\
 &= \lim_{t \rightarrow 4} [-4.9(t + 4)] = -39.2 \text{ m/sec}
 \end{aligned}$$

The object is falling at about 39.2 m/sec.

$$34. \quad -4.9t^2 + 250 = 0 \Rightarrow t = \frac{50}{7} \text{ sec}$$

When $a = \frac{50}{7}$, the velocity is

$$\begin{aligned}
 \lim_{t \rightarrow a} \frac{s(a) - s(t)}{a - t} &= \lim_{t \rightarrow a} \frac{[-4.9a^2 + 250] - [-4.9t^2 + 250]}{a - t} \\
 &= \lim_{t \rightarrow a} \frac{4.9(t^2 - a^2)}{a - t} \\
 &= \lim_{t \rightarrow a} \frac{4.9(t - a)(t + a)}{a - t} \\
 &= \lim_{t \rightarrow a} [-4.9(t + a)] \\
 &= -4.9(2a) \quad \left(a = \frac{50}{7} \right) \\
 &= -70 \text{ m/sec.}
 \end{aligned}$$

The velocity of the object when it hits the ground is about 70 m/sec.

$$35. \quad \lim_{x \rightarrow 3^-} \frac{|x - 3|}{x - 3} = \lim_{x \rightarrow 3^-} \frac{-(x - 3)}{x - 3} = -1$$

36. $\lim_{x \rightarrow 4} \llbracket x - 1 \rrbracket$ does not exist. There is a break in the graph at $x = 4$.

$$37. \quad \lim_{x \rightarrow 2} f(x) = 0$$

$$38. \quad \lim_{x \rightarrow 1^+} g(x) = 1 + 1 = 2$$

39. $\lim_{t \rightarrow 1} h(t)$ does not exist because $\lim_{t \rightarrow 1^-} h(t) = 1 + 1 = 2$

$$\text{and } \lim_{t \rightarrow 1^+} h(t) = \frac{1}{2}(1 + 1) = 1.$$

$$40. \quad \lim_{s \rightarrow -2} f(s) = 2$$

$$41. \quad f(x) = -3x^2 + 7$$

Continuous on $(-\infty, \infty)$

$$42. \quad f(x) = x^2 - \frac{2}{x}$$

Continuous on $(-\infty, 0) \cup (0, \infty)$

$$43. \quad f(x) = \llbracket x + 3 \rrbracket$$

$$\lim_{x \rightarrow k^+} \llbracket x + 3 \rrbracket = k + 3 \text{ where } k \text{ is an integer.}$$

$$\lim_{x \rightarrow k^-} \llbracket x + 3 \rrbracket = k + 2 \text{ where } k \text{ is an integer.}$$

Nonremovable discontinuity at each integer k

Continuous on $(k, k + 1)$ for all integers k

$$44. \quad f(x) = \frac{3x^2 - x - 2}{x - 1} = \frac{(3x + 2)(x - 1)}{x - 1}$$

Continuous on $(-\infty, 1) \cup (1, \infty)$

$$45. \quad f(x) = \frac{3x^2 - x - 2}{x - 1} = \frac{(3x + 2)(x - 1)}{x - 1}$$

$$\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} (3x + 2) = 5$$

Removable discontinuity at $x = 1$

Continuous on $(-\infty, 1) \cup (1, \infty)$

46. $f(x) = \begin{cases} 5 - x, & x \leq 2 \\ 2x - 3, & x > 2 \end{cases}$

$\lim_{x \rightarrow 2^-} (5 - x) = 3$

$\lim_{x \rightarrow 2^+} (2x - 3) = 1$

Nonremovable discontinuity at $x = 2$

Continuous on $(-\infty, 2) \cup (2, \infty)$

47. $f(x) = \frac{1}{(x - 2)^2}$

$\lim_{x \rightarrow 2} \frac{1}{(x - 2)^2} = \infty$

Nonremovable discontinuity at $x = 2$

Continuous on $(-\infty, 2) \cup (2, \infty)$

48. $f(x) = \sqrt{\frac{x+1}{x}} = \sqrt{1 + \frac{1}{x}}$

$\lim_{x \rightarrow 0^+} \sqrt{1 + \frac{1}{x}} = \infty$

Domain: $(-\infty, -1], (0, \infty)$

Nonremovable discontinuity at $x = 0$

Continuous on $(-\infty, -1] \cup (0, \infty)$

49. $f(x) = \frac{3}{x+1}$

$\lim_{x \rightarrow 1^-} f(x) = -\infty$

$\lim_{x \rightarrow 1^+} f(x) = \infty$

Nonremovable discontinuity at $x = -1$

Continuous on $(-\infty, -1) \cup (-1, \infty)$

54. $\lim_{x \rightarrow 1^+} (x + 1) = 2$

$\lim_{x \rightarrow 3^-} (x + 1) = 4$

Find b and c so that $\lim_{x \rightarrow 1^-} (x^2 + bx + c) = 2$ and $\lim_{x \rightarrow 3^+} (x^2 + bx + c) = 4$.

Consequently you get $1 + b + c = 2$ and $9 + 3b + c = 4$.

Solving simultaneously, $b = -3$ and $c = 4$.

55. f is continuous on $[1, 2]$. $f(1) = -1 < 0$ and

$f(2) = 13 > 0$. Therefore by the Intermediate Value

Theorem, there is at least one value c in $(1, 2)$ such that

$2c^3 - 3 = 0$.

50. $f(x) = \frac{x+1}{2x+2}$

$\lim_{x \rightarrow -1} \frac{x+1}{2(x+1)} = \frac{1}{2}$

Removable discontinuity at $x = -1$

Continuous on $(-\infty, -1) \cup (-1, \infty)$

51. $f(x) = \csc \frac{\pi x}{2}$

Nonremovable discontinuities at each even integer.

Continuous on

$(2k, 2k + 2)$

for all integers k .

52. $f(x) = \tan 2x$

Nonremovable discontinuities when

$x = \frac{(2n+1)\pi}{4}$

Continuous on

$\left(\frac{(2n-1)\pi}{4}, \frac{(2n+1)\pi}{4}\right)$

for all integers n .

53. $f(2) = 5$

Find c so that $\lim_{x \rightarrow 2^+} (cx + 6) = 5$.


$c(2) + 6 = 5$

$2c = -1$

$c = -\frac{1}{2}$

56. $C(x) = 12.80 + 2.50[-\lceil -x \rceil - 1], \quad x > 0$

$= 12.80 - 2.50[\lceil -x \rceil + 1], \quad x > 0$



C has a nonremovable discontinuity at each integer

$1, 2, 3, \dots$

$$57. f(x) = \frac{x^2 - 4}{|x - 2|} = (x + 2) \left[\frac{x - 2}{|x - 2|} \right]$$

- (a) $\lim_{x \rightarrow 2^-} f(x) = -4$
 (b) $\lim_{x \rightarrow 2^+} f(x) = 4$
 (c) $\lim_{x \rightarrow 2} f(x)$ does not exist.

$$58. f(x) = \sqrt{(x-1)x}$$

- (a) Domain: $(-\infty, 0] \cup [1, \infty)$
 (b) $\lim_{x \rightarrow 0^-} f(x) = 0$
 (c) $\lim_{x \rightarrow 1^+} f(x) = 0$

$$59. g(x) = 1 + \frac{2}{x}$$

Vertical asymptote at $x = 0$

$$60. h(x) = \frac{4x}{4 - x^2}$$

Vertical asymptotes at $x = 2$ and $x = -2$

$$61. f(x) = \frac{8}{(x-10)^2}$$

Vertical asymptote at $x = 10$

$$62. f(x) = \csc \pi x$$

Vertical asymptote at every integer k

$$63. \lim_{x \rightarrow -2^-} \frac{2x^2 + x + 1}{x + 2} = -\infty$$

$$64. \lim_{x \rightarrow (1/2)^+} \frac{x}{2x - 1} = \infty$$

$$65. \lim_{x \rightarrow -1^+} \frac{x+1}{x^3+1} = \lim_{x \rightarrow -1^+} \frac{1}{x^2-x+1} = \frac{1}{3}$$

$$76. f(x) = \frac{\tan 2x}{x}$$

(a)

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
$f(x)$	2.0271	2.0003	2.0000	2.0000	2.0003	2.0271

$$\lim_{x \rightarrow 0} \frac{\tan 2x}{x} = 2$$

(b) Yes, define $f(x) = \begin{cases} \frac{\tan 2x}{x}, & x \neq 0 \\ 2, & x = 0 \end{cases}$.

Now $f(x)$ is continuous at $x = 0$.

$$66. \lim_{x \rightarrow -1^-} \frac{x+1}{x^4-1} = \lim_{x \rightarrow -1^-} \frac{1}{(x^2+1)(x-1)} = -\frac{1}{4}$$

$$67. \lim_{x \rightarrow 1^-} \frac{x^2 + 2x + 1}{x - 1} = -\infty$$

$$68. \lim_{x \rightarrow -1^+} \frac{x^2 - 2x + 1}{x + 1} = \infty$$

$$69. \lim_{x \rightarrow 0^+} \left(x - \frac{1}{x^3} \right) = -\infty$$

$$70. \lim_{x \rightarrow 2^-} \frac{1}{\sqrt[3]{x^2 - 4}} = -\infty$$

$$71. \lim_{x \rightarrow 0^+} \frac{\sin 4x}{5x} = \lim_{x \rightarrow 0^+} \left[\frac{4}{5} \left(\frac{\sin 4x}{4x} \right) \right] = \frac{4}{5}$$

$$72. \lim_{x \rightarrow 0^+} \frac{\sec x}{x} = \infty$$

$$73. \lim_{x \rightarrow 0^+} \frac{\csc 2x}{x} = \lim_{x \rightarrow 0^+} \frac{1}{x \sin 2x} = \infty$$

$$74. \lim_{x \rightarrow 0^-} \frac{\cos^2 x}{x} = -\infty$$

$$75. C = \frac{80,000p}{100 - p}, 0 \leq p < 100$$

(a) $C(15) \approx \$14,117.65$

(b) $C(50) = \$80,000$

(c) $C(90) = \$720,000$

(d) $\lim_{p \rightarrow 100^-} \frac{80,000p}{100 - p} = \infty$

Problem Solving for Chapter 1

1. (a) Perimeter $\Delta PAO = \sqrt{x^2 + (y-1)^2} + \sqrt{x^2 + y^2} + 1$
 $= \sqrt{x^2 + (x^2 - 1)^2} + \sqrt{x^2 + x^4} + 1$
 Perimeter $\Delta PBO = \sqrt{(x-1)^2 + y^2} + \sqrt{x^2 + y^2} + 1$
 $= \sqrt{(x-1)^2 + x^4} + \sqrt{x^2 + x^4} + 1$

(b) $r(x) = \frac{\sqrt{x^2 + (x^2 - 1)^2} + \sqrt{x^2 + x^4} + 1}{\sqrt{(x-1)^2 + x^4} + \sqrt{x^2 + x^4} + 1}$

x	4	2	1	0.1	0.01
Perimeter ΔPAO	33.02	9.08	3.41	2.10	2.01
Perimeter ΔPBO	33.77	9.60	3.41	2.00	2.00
$r(x)$	0.98	0.95	1	1.05	1.005

(c) $\lim_{x \rightarrow 0^+} r(x) = \frac{1 + 0 + 1}{1 + 0 + 1} = \frac{2}{2} = 1$

2. (a) Area $\Delta PAO = \frac{1}{2}bh = \frac{1}{2}(1)(x) = \frac{x}{2}$
 Area $\Delta PBO = \frac{1}{2}bh = \frac{1}{2}(1)(y) = \frac{y}{2} = \frac{x^2}{2}$



(b) $a(x) = \frac{\text{Area } \Delta PBO}{\text{Area } \Delta PAO} = \frac{x^2/2}{x/2} = x$

x	4	2	1	0.1	0.01
Area ΔPAO	2	1	1/2	1/20	1/200
Area ΔPBO	8	2	1/2	1/200	1/20,000
$a(x)$	4	2	1	1/10	1/100

(c) $\lim_{x \rightarrow 0^+} a(x) = \lim_{x \rightarrow 0^+} x = 0$

3. (a) There are 6 triangles, each with a central angle of $60^\circ = \pi/3$. So,

$$\text{Area hexagon} = 6 \left[\frac{1}{2}bh \right] = 6 \left[\frac{1}{2}(1) \sin \frac{\pi}{3} \right] = \frac{3\sqrt{3}}{2} \approx 2.598.$$



$$\text{Error} = \text{Area (Circle)} - \text{Area (Hexagon)} = \pi - \frac{3\sqrt{3}}{2} \approx 0.5435$$

(b) There are n triangles, each with central angle of $\theta = 2\pi/n$. So,

$$A_n = n \left[\frac{1}{2}bh \right] = n \left[\frac{1}{2}(1) \sin \frac{2\pi}{n} \right] = \frac{n \sin(2\pi/n)}{2}.$$

(c)

n	6	12	24	48	96
A_n	2.598	3	3.106	3.133	3.139

(d) As n gets larger and larger, $2\pi/n$ approaches 0. Letting $x = 2\pi/n$, $A_n = \frac{\sin(2\pi/n)}{2/n} = \frac{\sin(2\pi/n)}{(2\pi/n)}\pi = \frac{\sin x}{x}\pi$ which approaches $(1)\pi = \pi$.

4. (a) Slope = $\frac{4 - 0}{3 - 0} = \frac{4}{3}$

(b) Slope = $-\frac{3}{4}$ Tangent line: $y - 4 = -\frac{3}{4}(x - 3)$
 $y = -\frac{3}{4}x + \frac{25}{4}$

(c) Let $Q = (x, y) = (x, \sqrt{25 - x^2})$

$$m_x = \frac{\sqrt{25 - x^2} - 4}{x - 3}$$

(d) $\lim_{x \rightarrow 3} m_x = \lim_{x \rightarrow 3} \frac{\sqrt{25 - x^2} - 4}{x - 3} \cdot \frac{\sqrt{25 - x^2} + 4}{\sqrt{25 - x^2} + 4}$
 $= \lim_{x \rightarrow 3} \frac{25 - x^2 - 16}{(x - 3)(\sqrt{25 - x^2} + 4)}$
 $= \lim_{x \rightarrow 3} \frac{(3 - x)(3 + x)}{(x - 3)(\sqrt{25 - x^2} + 4)}$
 $= \lim_{x \rightarrow 3} \frac{-(3 + x)}{\sqrt{25 - x^2} + 4} = \frac{-6}{4 + 4} = -\frac{3}{4}$

This is the slope of the tangent line at P .

5. (a) Slope = $-\frac{12}{5}$

(b) Slope of tangent line is $\frac{5}{12}$.

$$y + 12 = \frac{5}{12}(x - 5)$$

$$y = \frac{5}{12}x - \frac{169}{12} \text{ Tangent line}$$

(c) $Q = (x, y) = (x, -\sqrt{169 - x^2})$

$$m_x = \frac{-\sqrt{169 - x^2} + 12}{x - 5}$$

(d) $\lim_{x \rightarrow 5} m_x = \lim_{x \rightarrow 5} \frac{12 - \sqrt{169 - x^2}}{x - 5} \cdot \frac{12 + \sqrt{169 - x^2}}{12 + \sqrt{169 - x^2}}$
 $= \lim_{x \rightarrow 5} \frac{144 - (169 - x^2)}{(x - 5)(12 + \sqrt{169 - x^2})}$
 $= \lim_{x \rightarrow 5} \frac{x^2 - 25}{(x - 5)(12 + \sqrt{169 - x^2})}$
 $= \lim_{x \rightarrow 5} \frac{(x + 5)}{12 + \sqrt{169 - x^2}} = \frac{10}{12 + 12} = \frac{5}{12}$

This is the same slope as part (b).

6. $\frac{\sqrt{a + bx} - \sqrt{3}}{x} = \frac{\sqrt{a + bx} - \sqrt{3}}{x} \cdot \frac{\sqrt{a + bx} + \sqrt{3}}{\sqrt{a + bx} + \sqrt{3}}$
 $= \frac{(a + bx) - 3}{x(\sqrt{a + bx} + \sqrt{3})}$

Letting $a = 3$ simplifies the numerator.

So,

$$\lim_{x \rightarrow 0} \frac{\sqrt{3 + bx} - \sqrt{3}}{x} = \lim_{x \rightarrow 0} \frac{bx}{x(\sqrt{3 + bx} + \sqrt{3})}$$

$$= \lim_{x \rightarrow 0} \frac{b}{\sqrt{3 + bx} + \sqrt{3}}$$

Setting $\frac{b}{\sqrt{3} + \sqrt{3}} = \sqrt{3}$, you obtain $b = 6$. So,

$a = 3$ and $b = 6$.


7. (a) $3 + x^{1/3} \geq 0$

$$x^{1/3} \geq -3$$

$$x \geq -27$$

Domain: $x \geq -27, x \neq 1$ or $[-27, 1) \cup (1, \infty)$

(b)



$$(c) \lim_{x \rightarrow -27^+} f(x) = \frac{\sqrt{3 + (-27)^{1/3}} - 2}{-27 - 1} = \frac{-2}{-28} = \frac{1}{14} \approx 0.0714$$

$$\begin{aligned} (d) \lim_{x \rightarrow 1} f(x) &= \lim_{x \rightarrow 1} \frac{\sqrt{3 + x^{1/3}} - 2}{x - 1} \cdot \frac{\sqrt{3 + x^{1/3}} + 2}{\sqrt{3 + x^{1/3}} + 2} \\ &= \lim_{x \rightarrow 1} \frac{3 + x^{1/3} - 4}{(x - 1)(\sqrt{3 + x^{1/3}} + 2)} \\ &= \lim_{x \rightarrow 1} \frac{x^{1/3} - 1}{(x^{1/3} - 1)(x^{2/3} + x^{1/3} + 1)(\sqrt{3 + x^{1/3}} + 2)} \\ &= \lim_{x \rightarrow 1} \frac{1}{(x^{2/3} + x^{1/3} + 1)(\sqrt{3 + x^{1/3}} + 2)} \\ &= \frac{1}{(1 + 1 + 1)(2 + 2)} = \frac{1}{12} \end{aligned}$$

8. $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (a^2 - 2) = a^2 - 2$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{ax}{\tan x} = a \left(\text{because } \lim_{x \rightarrow 0} \frac{\tan x}{x} = 1 \right)$$

Thus,

$$a^2 - 2 = a$$

$$a^2 - a - 2 = 0$$

$$(a - 2)(a + 1) = 0$$


$$a = -1, 2$$

9. (a) $\lim_{x \rightarrow 2} f(x) = 3$; g_1, g_4

(b) f continuous at 2: g_1

(c) $\lim_{x \rightarrow 2^-} f(x) = 3$; g_1, g_3, g_4

10.



(a) $f\left(\frac{1}{4}\right) = \llbracket 4 \rrbracket = 4$

$$f(3) = \left\lfloor \frac{1}{3} \right\rfloor = 0$$

$$f(1) = \llbracket 1 \rrbracket = 1$$

(b) $\lim_{x \rightarrow 1^-} f(x) = 1$

$$\lim_{x \rightarrow 1^+} f(x) = 0$$


$$\lim_{x \rightarrow 0^-} f(x) = -\infty$$

$$\lim_{x \rightarrow 0^+} f(x) = \infty$$

(c) f is continuous for all real numbers except

$$x = 0, \pm 1, \pm \frac{1}{2}, \pm \frac{1}{3}, \dots$$

11.



- (a) $f(1) = \llbracket 1 \rrbracket + \llbracket -1 \rrbracket = 1 + (-1) = 0$
 $f(0) = 0$
 $f\left(\frac{1}{2}\right) = 0 + (-1) = -1$
 $f(-2.7) = -3 + 2 = -1$
- (b) $\lim_{x \rightarrow 1^-} f(x) = -1$
 $\lim_{x \rightarrow 1^+} f(x) = -1$
 $\lim_{x \rightarrow 1/2} f(x) = -1$
- (c) f is continuous for all real numbers except
 $x = 0, \pm 1, \pm 2, \pm 3, \dots$

12. (a) $v^2 = \frac{192,000}{r} + v_0^2 - 48$

$$\frac{192,000}{r} = v^2 - v_0^2 + 48$$

$$r = \frac{192,000}{v^2 - v_0^2 + 48}$$

$$\lim_{v \rightarrow 0} r = \frac{192,000}{48 - v_0^2}$$

Let $v_0 = \sqrt{48} = 4\sqrt{3}$ mi/sec.

(b) $v^2 = \frac{1920}{r} + v_0^2 - 2.17$

$$\frac{1920}{r} = v^2 - v_0^2 + 2.17$$

$$r = \frac{1920}{v^2 - v_0^2 + 2.17}$$

$$\lim_{v \rightarrow 0} r = \frac{1920}{2.17 - v_0^2}$$

Let $v_0 = \sqrt{2.17}$ mi/sec (≈ 1.47 mi/sec).


(c) $r = \frac{10,600}{v^2 - v_0^2 + 6.99}$

$$\lim_{v \rightarrow 0} r = \frac{10,600}{6.99 - v_0^2}$$

Let $v_0 = \sqrt{6.99} \approx 2.64$ mi/sec.

Because this is smaller than the escape velocity for Earth, the mass is less.

13. (a)



- (b) (i) $\lim_{x \rightarrow a^+} P_{a,b}(x) = 1$
 (ii) $\lim_{x \rightarrow a^-} P_{a,b}(x) = 0$
 (iii) $\lim_{x \rightarrow b^+} P_{a,b}(x) = 0$
 (iv) $\lim_{x \rightarrow b^-} P_{a,b}(x) = 1$
- (c) $P_{a,b}$ is continuous for all positive real numbers except $x = a, b$.
- (d) The area under the graph of U , and above the x -axis, is 1.

14. Let $a \neq 0$ and let $\varepsilon > 0$ be given. There exists $\delta_1 > 0$ such that if $0 < |x - 0| < \delta_1$ then $|f(x) - L| < \varepsilon$. Let $\delta = \delta_1/|a|$. Then for $0 < |x - 0| < \delta = \delta_1/|a|$, you have

$$|x| < \frac{\delta_1}{|a|}$$

$$|ax| < \delta_1$$

$$|f(ax) - L| < \varepsilon.$$

As a counterexample, let

$$a = 0 \text{ and } f(x) = \begin{cases} 1, & x \neq 0 \\ 2, & x = 0 \end{cases}$$

Then $\lim_{x \rightarrow 0} f(x) = 1 = L$, but

$$\lim_{x \rightarrow 0} f(ax) = \lim_{x \rightarrow 0} f(0) = \lim_{x \rightarrow 0} 2 = 2.$$