

Solutions for

EXERCISES
for

Weather & Climate

NINTH EDITION

Greg Carbone

University of South Carolina – Columbia

PEARSON

Copyright © 2016, 2013, 2010, 2004, 2001 Pearson Education, Inc. All rights reserved. Printed in the United States of America. This publication is protected by copyright, and permission should be obtained from the publisher prior to any prohibited reproduction, storage in a retrieval system, or transmission in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise. For information regarding permissions, request forms and the appropriate contacts within the Pearson Education Global Rights & Permissions department, please visit <http://www.pearsoned.com/permissions/>.

Unless otherwise indicated herein, any third-party trademarks that may appear in this work are the property of their respective owners and any references to third-party trademarks, logos or other trade dress are for demonstrative or descriptive purposes only. Such references are not intended to imply any sponsorship, endorsement, authorization, or promotion of Pearson's products by the owners of such marks, or any relationship between the owner and Pearson Education, Inc. or its affiliates, authors, licensees or distributors.

PEARSON

www.pearsonhighered.com

Contents

1	Vertical Structure of the Atmosphere	1
2	Earth–Sun Geometry	5
3	The Surface Energy Budget	9
4	The Global Energy Budget	12
5	Atmospheric Moisture	14
6	Saturation and Atmospheric Stability	18
7	Cloud Droplets and Raindrops	21
8	Atmospheric Motion	23
9	Weather Map Analysis	30
10	Mid-Latitude Cyclones	35
11	Weather Forecasting	40
12	Thunderstorms and Tornadoes	45
13	Hurricanes	48
14	Climate Controls	52
15	Climate Classification	55
16	Climatic Variability and Change	57
17	Simulating Climate Change	60
<i>Appendix A</i>	Dimensions and Units	62
<i>Appendix B</i>	Earth Measures	64


1

Vertical Structure of the Atmosphere

1. Height (km) % of atmosphere above

22.4	6.25
16.8	12.5
11.2	25
5.6	50

2. & 3.



4. 25% 250 mb
58.4% 584 mb


5. 210 mb

6. 123 mb 58.4%
69 mb 33%

7. Ozone absorbs solar radiation (particularly in the ultraviolet portion of the electromagnetic spectrum). This absorption leads to warming in the stratosphere.

8.	2000	4000	6000	8000	10,000
	2.0°C	-11°C	-24°C	-37°C	-50°C

9. & 11.



10. a. Key West b. Key West c. Fairbanks

11. See 9 above.

12. Key West tropopause: ~16,000 m, ~ -75°C; Fairbanks tropopause: ~10,000 m, ~ -53°C;

13. The greater the average temperature, the higher the tropopause. Our example suggests that vertical mixing is greater when temperature is warmer.

14. 170 mb

15. 92 mb

16. Because of greater air density in the lower layer, the pressure drop between 2 and 4 km is nearly double that between 8 and 10 km.


17. 182 mb

18. Air pressure decreases with height because there is less atmosphere to exert downward force. The pressure drop will be greatest when air density is highest because the mass of the atmosphere above decreases at a faster rate.


19. California desert: 1003.9 mb; Michigan UP: 1018.6 mb; New Brunswick: 1003.7 mb.

20. The Michigan and New Orleans stations have the same pressure (1018.6 mb), but a 30°F temperature difference. The New Brunswick and southern California stations have similar low pressures (1003.7 mb and 1003.9 mb), but a 30°F temperature difference.
21. The ideal gas law shows that pressure is proportional to the product of density times temperature. Therefore, to have a similar pressure, but be 30°F warmer, New Orleans must have a lower density.
22. The Michigan and New Brunswick stations have higher air density than the other two.
23. The 1.4 g/kg mixing ratio in Fairbanks is considerably smaller than that at the surface of Key West (16 g/kg) on the same day. In fact the surface moisture value at Fairbanks is as low as the mixing ratio value 10 km above Key West.
24. The atomic weight of hydrogen equals 1; the atomic weight of oxygen equals 16. Therefore, the atomic weight of H₂O is 1 g/mol + 1 g/mol + 16 g/mol = 18 g/mol.
25. The solution to the previous question shows that water vapor, at 18 g/mol, is lighter than dry air (28.6 g/mol).

26.



27.




28. Minimum atmospheric CO₂ concentrations occur during the northern hemisphere's late fall at which time CO₂ has been taken up by vegetation for the previous 6–8 months. Atmospheric CO₂ concentrations occur in April/May after many months of leaf decay and just before spring green-up in the Northern Hemisphere.

Review Questions


1. Air pressure and density decrease exponentially with height above Earth's surface. This is because gas molecules are concentrated near the surface and a given height increase at these lower levels means passing through more molecules than the same height increase at higher elevations. Temperature also decreases with height in the troposphere. This rate of decrease varies, but is typically linear compared to pressure or density.
2. The thickness of the troposphere is a function of temperature. Warmer temperatures in tropical regions create mixing to greater depths, pushing the tropopause higher.
3. Pressure changes much faster vertically than it does horizontally. It drops 100 mb in the lowest kilometer of the atmosphere.

2 Earth–Sun Geometry


1. March 21




March 21 profile view



June 21



June 21 profile view



2. 63.5°; December 21

3. 26.5°

4. a. 0° (equator)
 b. 23.5° N
 c. 0° (equator)
 d. 23.5° S
 e. [variable]

5.

	New Orleans	Helsinki
a.	60°	30°
b.	83.5°	53.5°
c.	60°	30°
d.	36.5°	6.5°
e.	[variable]	[variable]

6. [variable]

7. Answer is date dependent. Example for 34° N latitude on February 1, a two-meter pole casts a shadow measuring 2.52 meters.


$$\tan \Theta = \frac{\text{length of pole}}{\text{length of shadow}}$$

$$\Theta = \tan^{-1}(0.7937)$$


$$\Theta = 38.44^\circ$$

8. [variable]


9. 60° N December 22



30° N June 21



60° N June 21



10. Summer temperature is highest because solar radiation is more concentrated. During the winter, it's cooler as the solar beam is spread over a greater surface area.


11. There is a much greater seasonal range in daylight hours in polar regions than in tropical regions.

	30° N	60° N
June solstice	14	18
Equinoxes	12	12
December solstice	10	6


13. 60° N

14. The steepness of the curves suggests that daylight hours increase and decrease faster from one day to the next near the equinoxes, and change slower from one day to the next near the solstices.

15.



16. On the December solstice, the sun rises a bit south of east and sets a bit south of west. On the June solstice, the sun rises in the northeast and sets in the northwest. At 60° N, this shift is even more dramatic.



18. The seasonal difference in solar intensity (beam spreading) and daylight hours is greater at 60° N than at 30° N.
19. The difference in beam spreading between 60° N and 30° N is very large in winter and not very different in summer. Furthermore, 60° N has a shorter daylight period than 30° N in winter, while in summer the daylight hours are actually greater at 60° N.
20. Most direct rays: 1 unit beam = 1.000 surface units; Date: March 21, September 22
 Least direct rays: 1 unit beam = 1.090 surface units; Date: June 21, December 21
21. 9%
22. [variable]
23. [variable]
24. The higher the latitude, the greater the seasonal range in solar intensity. This results in a larger annual temperature range at high latitudes than in the tropics.
- 25.
- | | December Solstice | June Solstice |
|-------|-------------------|---------------|
| 60° N | 8.834 | 1.244 |
| 50° N | 3.521 | 1.117 |
| 40° N | 2.241 | 1.043 |
| 30° N | 1.681 | 1.006 |
| 20° N | 1.379 | 1.002 |
26. The solar intensity gradient across the mid-latitudes is much greater in winter and contributes to a greater temperature gradient.

Review Questions

1. A given change at low sun angles is much more effective than the same change at higher sun angles. Therefore, the seasonal shift of sun angle from 36.5° to 83.5° at New Orleans results in less change in solar intensity than the shift from 6.5° to 53.5° at Helsinki.
2. A greater range in solar intensity and daylight hours will result in a greater range in solar radiation received and temperature.