

# Complete Solutions Manual to Accompany

---

## Probability and Statistics for Engineering and the Sciences

**NINTH EDITION**

**Jay Devore**

California Polytechnic State University,  
San Luis Obispo, CA

Prepared by

**Matthew A. Carlton**

California Polytechnic State University, San Luis Obispo, CA



---

Australia • Brazil • Mexico • Singapore • United Kingdom • United States

© 2016 Cengage Learning

ALL RIGHTS RESERVED. No part of this work covered by the copyright herein may be reproduced, transmitted, stored, or used in any form or by any means graphic, electronic, or mechanical, including but not limited to photocopying, recording, scanning, digitizing, taping, Web distribution, information networks, or information storage and retrieval systems, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without the prior written permission of the publisher except as may be permitted by the license terms below.

For product information and technology assistance, contact us at  
**Cengage Learning Customer & Sales Support,**  
**1-800-354-9706.**

For permission to use material from this text or product, submit  
all requests online at [www.cengage.com/permissions](http://www.cengage.com/permissions)  
Further permissions questions can be emailed to  
[permissionrequest@cengage.com](mailto:permissionrequest@cengage.com).

ISBN-13: 978-1-305-26061-0  
ISBN-10: 1-305-26061-9

**Cengage Learning**  
20 Channel Center Street  
Fourth Floor  
Boston, MA 02210  
USA

Cengage Learning is a leading provider of customized learning solutions with office locations around the globe, including Singapore, the United Kingdom, Australia, Mexico, Brazil, and Japan. Locate your local office at:  
[www.cengage.com/global](http://www.cengage.com/global).

Cengage Learning products are represented in Canada by Nelson Education, Ltd.

To learn more about Cengage Learning Solutions, visit [www.cengage.com](http://www.cengage.com).

Purchase any of our products at your local college store or at our preferred online store  
[www.cengagebrain.com](http://www.cengagebrain.com).

**NOTE: UNDER NO CIRCUMSTANCES MAY THIS MATERIAL OR ANY PORTION THEREOF BE SOLD, LICENSED, AUCTIONED, OR OTHERWISE REDISTRIBUTED EXCEPT AS MAY BE PERMITTED BY THE LICENSE TERMS HEREIN.**

#### READ IMPORTANT LICENSE INFORMATION

Dear Professor or Other Supplement Recipient:

Cengage Learning has provided you with this product (the "Supplement") for your review and, to the extent that you adopt the associated textbook for use in connection with your course (the "Course"), you and your students who purchase the textbook may use the Supplement as described below. Cengage Learning has established these use limitations in response to concerns raised by authors, professors, and other users regarding the pedagogical problems stemming from unlimited distribution of Supplements.

Cengage Learning hereby grants you a nontransferable license to use the Supplement in connection with the Course, subject to the following conditions. The Supplement is for your personal, noncommercial use only and may not be reproduced, posted electronically or distributed, except that portions of the Supplement may be provided to your students IN PRINT FORM ONLY in connection with your instruction of the Course, so long as such students are advised that they

may not copy or distribute any portion of the Supplement to any third party. You may not sell, license, auction, or otherwise redistribute the Supplement in any form. We ask that you take reasonable steps to protect the Supplement from unauthorized use, reproduction, or distribution. Your use of the Supplement indicates your acceptance of the conditions set forth in this Agreement. If you do not accept these conditions, you must return the Supplement unused within 30 days of receipt.

All rights (including without limitation, copyrights, patents, and trade secrets) in the Supplement are and will remain the sole and exclusive property of Cengage Learning and/or its licensors. The Supplement is furnished by Cengage Learning on an "as is" basis without any warranties, express or implied. This Agreement will be governed by and construed pursuant to the laws of the State of New York, without regard to such State's conflict of law rules.

Thank you for your assistance in helping to safeguard the integrity of the content contained in this Supplement. We trust you find the Supplement a useful teaching tool.

# CONTENTS

Chapter 1	Overview and Descriptive Statistics	1
Chapter 2	Probability	48
Chapter 3	Discrete Random Variables and Probability Distributions	90
Chapter 4	Continuous Random Variables and Probability Distributions	126
Chapter 5	Joint Probability Distributions and Random Samples	177
Chapter 6	Point Estimation	206
Chapter 7	Statistical Intervals Based on a Single Sample	217
Chapter 8	Tests of Hypotheses Based on a Single Sample	234
Chapter 9	Inferences Based on Two Samples	255
Chapter 10	The Analysis of Variance	285
Chapter 11	Multifactor Analysis of Variance	299
Chapter 12	Simple Linear Regression and Correlation	330
Chapter 13	Nonlinear and Multiple Regression	368
Chapter 14	Goodness-of-Fit Tests and Categorical Data Analysis	406
Chapter 15	Distribution-Free Procedures	424
Chapter 16	Quality Control Methods	434

# CHAPTER 1

## Section 1.1

1.
  - a. *Los Angeles Times, Oberlin Tribune, Gainesville Sun, Washington Post*
  - b. Duke Energy, Clorox, Seagate, Neiman Marcus
  - c. Vince Correa, Catherine Miller, Michael Cutler, Ken Lee
  - d. 2.97, 3.56, 2.20, 2.97
  
2.
  - a. 29.1 yd, 28.3 yd, 24.7 yd, 31.0 yd
  - b. 432 pp, 196 pp, 184 pp, 321 pp
  - c. 2.1, 4.0, 3.2, 6.3
  - d. 0.07 g, 1.58 g, 7.1 g, 27.2 g
  
3.
  - a. How likely is it that more than half of the sampled computers will need or have needed warranty service? What is the expected number among the 100 that need warranty service? How likely is it that the number needing warranty service will exceed the expected number by more than 10?
  - b. Suppose that 15 of the 100 sampled needed warranty service. How confident can we be that the proportion of *all* such computers needing warranty service is between .08 and .22? Does the sample provide compelling evidence for concluding that more than 10% of all such computers need warranty service?

## Chapter 1: Overview and Descriptive Statistics

- 4.
- a. Concrete populations: all living U.S. Citizens, all mutual funds marketed in the U.S., all books published in 1980  
Hypothetical populations: all grade point averages for University of California undergraduates during the next academic year, page lengths for all books published during the next calendar year, batting averages for all major league players during the next baseball season
  - b. (Concrete) Probability: In a sample of 5 mutual funds, what is the chance that all 5 have rates of return which exceeded 10% last year?  
Statistics: If previous year rates-of-return for 5 mutual funds were 9.6, 14.5, 8.3, 9.9 and 10.2, can we conclude that the average rate for all funds was below 10%?  
(Hypothetical) Probability: In a sample of 10 books to be published next year, how likely is it that the average number of pages for the 10 is between 200 and 250?  
Statistics: If the sample average number of pages for 10 books is 227, can we be highly confident that the average for all books is between 200 and 245?
- 5.
- a. No. All students taking a large statistics course who participate in an SI program of this sort.
  - b. The advantage to randomly allocating students to the two groups is that the two groups should then be fairly comparable before the study. If the two groups perform differently in the class, we might attribute this to the treatments (SI and control). If it were left to students to choose, stronger or more dedicated students might gravitate toward SI, confounding the results.
  - c. If all students were put in the treatment group, there would be no firm basis for assessing the effectiveness of SI (nothing to which the SI scores could reasonably be compared).
- 6.
- One could take a simple random sample of students from all students in the California State University system and ask each student in the sample to report the distance from their hometown to campus. Alternatively, the sample could be generated by taking a stratified random sample by taking a simple random sample from each of the 23 campuses and again asking each student in the sample to report the distance from their hometown to campus. Certain problems might arise with self reporting of distances, such as recording error or poor recall. This study is enumerative because there exists a finite, identifiable population of objects from which to sample.
- 7.
- One could generate a simple random sample of all single-family homes in the city, or a stratified random sample by taking a simple random sample from each of the 10 district neighborhoods. From each of the selected homes, values of all desired variables would be determined. This would be an enumerative study because there exists a finite, identifiable population of objects from which to sample.

## Chapter 1: Overview and Descriptive Statistics

- 8.
- a. Number observations equal  $2 \times 2 \times 2 = 8$
  - b. This could be called an analytic study because the data would be collected on an existing process. There is no sampling frame.
- 9.
- a. There could be several explanations for the variability of the measurements. Among them could be measurement error (due to mechanical or technical changes across measurements), recording error, differences in weather conditions at time of measurements, etc.
  - b. No, because there is no sampling frame.

### Section 1.2

- 10.
- a.

5	9	
6	33588	
7	00234677889	
8	127	
9	077	stem: ones
10	7	leaf: tenths
11	368	

A representative strength for these beams is around 7.8 MPa, but there is a reasonably large amount of variation around that representative value.

(What constitutes large or small variation usually depends on context, but variation is usually considered large when the range of the data – the difference between the largest and smallest value – is comparable to a representative value. Here, the range is  $11.8 - 5.9 = 5.9$  MPa, which is similar in size to the representative value of 7.8 MPa. So, most researchers would call this a large amount of variation.)

- b. The data display is not perfectly symmetric around some middle/representative value. There is some positive skewness in this data.
- c. Outliers are data points that appear to be *very* different from the pack. Looking at the stem-and-leaf display in part (a), there appear to be no outliers in this data. (A later section gives a more precise definition of what constitutes an outlier.)
- d. From the stem-and-leaf display in part (a), there are 4 values greater than 10. Therefore, the proportion of data values that exceed 10 is  $4/27 = .148$ , or, about 15%.

## Chapter 1: Overview and Descriptive Statistics

11.


3L	1	
3H	56678	
4L	000112222234	
4H	5667888	stem: tenths
5L	144	leaf : hundredths
5H	58	
6L	2	
6H	6678	
7L		
7H	5	

The stem-and-leaf display shows that .45 is a good representative value for the data. In addition, the display is not symmetric and appears to be positively skewed. The range of the data is  $.75 - .31 = .44$ , which is comparable to the typical value of .45. This constitutes a reasonably large amount of variation in the data. The data value .75 is a possible outlier.

12.

The sample size for this data set is  $n = 5 + 15 + 27 + 34 + 22 + 14 + 7 + 2 + 4 + 1 = 131$ .

- a. The first four intervals correspond to observations less than 5, so the proportion of values less than 5 is  $(5 + 15 + 27 + 34)/131 = 81/131 = .618$ .
- b. The last four intervals correspond to observations at least 6, so the proportion of values at least 6 is  $(7 + 2 + 4 + 1)/131 = 14/131 = .107$ .
- c. & d. The relative (percent) frequency and density histograms appear below. The distribution of  $\text{CeO}_2$  sizes is not symmetric, but rather positively skewed. Notice that the relative frequency and density histograms are essentially identical, other than the vertical axis labeling, because the bin widths are all the same.







## Chapter 1: Overview and Descriptive Statistics

14.

a.

2	23	stem: 1.0
3	2344567789	leaf: .10
4	01356889	
5	00001114455666789	
6	0000122223344456667789999	
7	00012233455555668	
8	02233448	
9	012233335666788	
10	2344455688	
11	2335999	
12	37	
13	8	
14	36	
15	0035	
16		
17		
18	9	

- b. A representative is around 7.0.
- c. The data exhibit a moderate amount of variation (this is subjective).
- d. No, the data is skewed to the right, or positively skewed.
- e. The value 18.9 appears to be an outlier, being more than two stem units from the previous value.

15.

American		French
	8	1
755543211000	9	00234566
9432	10	2356
6630	11	1369
850	12	223558
8	13	7
	14	
	15	8
2	16	

American movie times are unimodal strongly positively skewed, while French movie times appear to be bimodal. A typical American movie runs about 95 minutes, while French movies are typically either around 95 minutes or around 125 minutes. American movies are generally shorter than French movies and are less variable in length. Finally, both American and French movies occasionally run very long (outliers at 162 minutes and 158 minutes, respectively, in the samples).

## Chapter 1: Overview and Descriptive Statistics

16.


a.

Beams		Cylinders	
9	5	8	
88533	6	16	
98877643200	7	012488	
721	8	13359	stem: ones
770	9	278	leaf: tenths
7	10		
863	11	2	
	12	6	
	13		
	14	1	

The data appears to be slightly skewed to the right, or positively skewed. The value of 14.1 MPa appears to be an outlier. Three out of the twenty, or 15%, of the observations exceed 10 MPa.

b. The majority of observations are between 5 and 9 MPa for both beams and cylinders, with the modal class being 7.0-7.9 MPa. The observations for cylinders are more variable, or spread out, and the maximum value of the cylinder observations is higher.


c.



17.

The sample size for this data set is  $n = 7 + 20 + 26 + \dots + 3 + 2 = 108$ .


- a. “At most five bidders” means 2, 3, 4, or 5 bidders. The proportion of contracts that involved at most 5 bidders is  $(7 + 20 + 26 + 16)/108 = 69/108 = .639$ . Similarly, the proportion of contracts that involved at least 5 bidders (5 through 11) is equal to  $(16 + 11 + 9 + 6 + 8 + 3 + 2)/108 = 55/108 = .509$ .
- b. The number of contracts with between 5 and 10 bidders, inclusive, is  $16 + 11 + 9 + 6 + 8 + 3 = 53$ , so the proportion is  $53/108 = .491$ . “Strictly” between 5 and 10 means 6, 7, 8, or 9 bidders, for a proportion equal to  $(11 + 9 + 6 + 8)/108 = 34/108 = .315$ .
- c. The distribution of number of bidders is positively skewed, ranging from 2 to 11 bidders, with a typical value of around 4-5 bidders.



## Chapter 1: Overview and Descriptive Statistics

18.

- a. The most interesting feature of the histogram is the heavy presence of three very large outliers (21, 24, and 32 directors). Absent these three corporations, the distribution of number of directors would be roughly symmetric with a typical value of around 9.



Note: One way to have Minitab automatically construct a histogram from grouped data such as this is to use Minitab's ability to enter multiple copies of the same number by typing, for example, 42(9) to enter 42 copies of the number 9. The frequency data in this exercise was entered using the following Minitab commands:

```
MTB > set c1
DATA> 3(4) 12(5) 13(6) 25(7) 24(8) 42(9) 23(10) 19(11) 16(12)
11(13) 5(14) 4(15) 1(16) 3(17) 1(21) 1(24) 1(32)
DATA> end
```

- b. The accompanying frequency distribution is nearly identical to the one in the textbook, except that the three largest values are compacted into the “ $\geq 18$ ” category. If this were the originally-presented information, we could not create a histogram, because we would not know the upper boundary for the rectangle corresponding to the “ $\geq 18$ ” category.


No. dir.	4	5	6	7	8	9	10	11
Freq.	3	12	13	25	24	42	23	19
No dir.	12	13	14	15	16	17	$\geq 18$	
Freq.	16	11	5	4	1	3	3	

- c. The sample size is  $3 + 12 + \dots + 3 + 1 + 1 + 1 = 204$ . So, the proportion of these corporations that have at most 10 directors is  $(3 + 12 + 13 + 25 + 24 + 42 + 23)/204 = 142/204 = .696$ .
- d. Similarly, the proportion of these corporations with more than 15 directors is  $(1 + 3 + 1 + 1 + 1)/204 = 7/204 = .034$ .

## Chapter 1: Overview and Descriptive Statistics

19.

- a. From this frequency distribution, the proportion of wafers that contained at least one particle is  $(100-1)/100 = .99$ , or 99%. Note that it is much easier to subtract 1 (which is the number of wafers that contain 0 particles) from 100 than it would be to add all the frequencies for 1, 2, 3, ... particles. In a similar fashion, the proportion containing at least 5 particles is  $(100 - 1-2-3-12-11)/100 = 71/100 = .71$ , or, 71%.
- b. The proportion containing between 5 and 10 particles is  $(15+18+10+12+4+5)/100 = 64/100 = .64$ , or 64%. The proportion that contain strictly between 5 and 10 (meaning strictly *more* than 5 and strictly *less* than 10) is  $(18+10+12+4)/100 = 44/100 = .44$ , or 44%.
- c. The following histogram was constructed using Minitab. The histogram is *almost* symmetric and unimodal; however, the distribution has a few smaller modes and has a very slight positive skew.



20.

- a. The following stem-and-leaf display was constructed:


```

0|123334555599
1|00122234688      stem: thousands
2|1112344477      leaf: hundreds
3|0113338
4|37
5|23778
    
```

A typical data value is somewhere in the low 2000's. The display is bimodal (the stem at 5 would be considered a mode, the stem at 0 another) and has a positive skew.


## Chapter 1: Overview and Descriptive Statistics

- b. A histogram of this data, using classes boundaries of 0, 1000, 2000, ..., 6000 is shown below. The proportion of subdivisions with total length less than 2000 is  $(12+11)/47 = .489$ , or 48.9%. Between 2000 and 4000, the proportion is  $(10+7)/47 = .362$ , or 36.2%. The histogram shows the same general shape as depicted by the stem-and-leaf in part (a).




21.

- a. A histogram of the  $y$  data appears below. From this histogram, the number of subdivisions having no cul-de-sacs (i.e.,  $y = 0$ ) is  $17/47 = .362$ , or 36.2%. The proportion having at least one cul-de-sac ( $y \geq 1$ ) is  $(47 - 17)/47 = 30/47 = .638$ , or 63.8%. Note that subtracting the number of cul-de-sacs with  $y = 0$  from the total, 47, is an easy way to find the number of subdivisions with  $y \geq 1$ .




## Chapter 1: Overview and Descriptive Statistics

- b. A histogram of the  $z$  data appears below. From this histogram, the number of subdivisions with at most 5 intersections (i.e.,  $z \leq 5$ ) is  $42/47 = .894$ , or 89.4%. The proportion having fewer than 5 intersections (i.e.,  $z < 5$ ) is  $39/47 = .830$ , or 83.0%.




22. A very large percentage of the data values are greater than 0, which indicates that most, but not all, runners do slow down at the end of the race. The histogram is also positively skewed, which means that some runners slow down a *lot* compared to the others. A typical value for this data would be in the neighborhood of 200 seconds. The proportion of the runners who ran the last 5 km faster than they did the first 5 km is very small, about 1% or so.
23. Note: since the class intervals have unequal length, we must use a *density scale*.



The distribution of tantrum durations is unimodal and heavily positively skewed. Most tantrums last between 0 and 11 minutes, but a few last more than half an hour! With such heavy skewness, it's difficult to give a representative value.


## Chapter 1: Overview and Descriptive Statistics

24. The distribution of shear strengths is roughly symmetric and bell-shaped, centered at about 5000 lbs and ranging from about 4000 to 6000 lbs.




25. The transformation creates a much more symmetric, mound-shaped histogram.

Histogram of original data:



## Chapter 1: Overview and Descriptive Statistics

Histogram of transformed data:



26.

- Yes: the proportion of sampled angles smaller than  $15^\circ$  is  $.177 + .166 + .175 = .518$ .
- The proportion of sampled angles at least  $30^\circ$  is  $.078 + .044 + .030 = .152$ .
- The proportion of angles between  $10^\circ$  and  $25^\circ$  is roughly  $.175 + .136 + (.194)/2 = .408$ .
- The distribution of misorientation angles is heavily positively skewed. Though angles can range from  $0^\circ$  to  $90^\circ$ , nearly 85% of all angles are less than  $30^\circ$ . Without more precise information, we cannot tell if the data contain outliers.






## Chapter 1: Overview and Descriptive Statistics

27.

- a. The endpoints of the class intervals overlap. For example, the value 50 falls in both of the intervals 0–50 and 50–100.
- b. The lifetime distribution is positively skewed. A representative value is around 100. There is a great deal of variability in lifetimes and several possible candidates for outliers.


<b>Class Interval</b>	<b>Frequency</b>	<b>Relative Frequency</b>
0–< 50	9	0.18
50–<100	19	0.38
100–<150	11	0.22
150–<200	4	0.08
200–<250	2	0.04
250–<300	2	0.04
300–<350	1	0.02
350–<400	1	0.02
400–<450	0	0.00
450–<500	0	0.00
500–<550	1	0.02
	<hr/> 50	<hr/> 1.00



## Chapter 1: Overview and Descriptive Statistics

- c. There is much more symmetry in the distribution of the transformed values than in the values themselves, and less variability. There are no longer gaps or obvious outliers.

<u>Class Interval</u>	<u>Frequency</u>	<u>Relative Frequency</u>
2.25–<2.75	2	0.04
2.75–<3.25	2	0.04
3.25–<3.75	3	0.06
3.75–<4.25	8	0.16
4.25–<4.75	18	0.36
4.75–<5.25	10	0.20
5.25–<5.75	4	0.08
5.75–<6.25	3	0.06




- d. The proportion of lifetime observations in this sample that are less than 100 is  $.18 + .38 = .56$ , and the proportion that is at least 200 is  $.04 + .04 + .02 + .02 + .02 = .14$ .

28. The sample size for this data set is  $n = 804$ .
- $(5 + 11 + 13 + 30 + 46)/804 = 105/804 = .131$ .
  - $(73 + 38 + 19 + 11)/804 = 141/804 = .175$ .
  - The number of trials resulting in deposited energy of 3.6 mJ or more is  $126 + 92 + 73 + 38 + 19 + 11 = 359$ . Additionally, 141 trials resulted in deposited energy within the interval 3.4–<3.6. If we assume that roughly half of these were in the interval 3.5–<3.6 (since 3.5 is the midpoint), then our estimated frequency is  $359 + (141)/2 = 429.5$ , for a rough proportion equal to  $429.5/804 = .534$ .

## Chapter 1: Overview and Descriptive Statistics


- d. The deposited energy distribution is roughly symmetric or perhaps slightly negatively skewed (there is a somewhat long left tail). Notice that the histogram must be made on a density scale, since the interval widths are not all the same.




# Chapter 1: Overview and Descriptive Statistics

29.

Physical Activity	Frequency	Relative Frequency
A	28	.28
B	19	.19
C	18	.18
D	17	.17
E	9	.09
F	9	.09
	100	1.00



30.



Chapter 1: Overview and Descriptive Statistics


31.

Class	Frequency	Cum. Freq.	Cum. Rel. Freq.
0.0–<4.0	2	2	0.050
4.0–<8.0	14	16	0.400
8.0–<12.0	11	27	0.675
12.0–<16.0	8	35	0.875
16.0–<20.0	4	39	0.975
20.0–<24.0	0	39	0.975
24.0–<28.0	1	40	1.000

32.

- a. Cumulative percents must be restored to relative frequencies. Then the histogram may be constructed (see below). The relative frequency distribution is almost unimodal and exhibits a large positive skew. The typical middle value is somewhere between 400 and 450, although the skewness makes it difficult to pinpoint more exactly than this.

Class	Rel. Freq.	Class	Rel. Freq.
0–< 150	.193	900–<1050	.019
150–< 300	.183	1050–<1200	.029
300–< 450	.251	1200–<1350	.005
450–< 600	.148	1350–<1500	.004
600–< 750	.097	1500–<1650	.001
750–< 900	.066	1650–<1800	.002
		1800–<1950	.002



- b. The proportion of the fire loads less than 600 is  $.193 + .183 + .251 + .148 = .775$ . The proportion of loads that are at least 1200 is  $.005 + .004 + .001 + .002 + .002 = .014$ .
- c. The proportion of loads between 600 and 1200 is  $1 - .775 - .014 = .211$ .

**Section 1.3**

33.

- a. Using software,  $\bar{x} = 640.5$  (\$640,500) and  $\tilde{x} = 582.5$  (\$582,500). The average sale price for a home in this sample was \$640,500. Half the sales were for less than \$582,500, while half were for more than \$582,500.
- b. Changing that one value lowers the sample mean to 610.5 (\$610,500) but has no effect on the sample median.
- c. After removing the two largest and two smallest values,  $\bar{x}_{tr(20)} = 591.2$  (\$591,200).
- d. A 10% trimmed mean from removing just the highest and lowest values is  $\bar{x}_{tr(10)} = 596.3$ . To form a 15% trimmed mean, take the average of the 10% and 20% trimmed means to get  $\bar{x}_{tr(15)} = (591.2 + 596.3)/2 = 593.75$  (\$593,750).

34.

- a. For urban homes,  $\bar{x} = 21.55$  EU/mg; for farm homes,  $\bar{x} = 8.56$  EU/mg. The average endotoxin concentration in urban homes is more than double the average endotoxin concentration in farm homes.
- b. For urban homes,  $\tilde{x} = 17.00$  EU/mg; for farm homes,  $\tilde{x} = 8.90$  EU/mg. The median endotoxin concentration in urban homes is nearly double the median endotoxin concentration in farm homes. The mean and median endotoxin concentration for urban homes are so different because the few large values, especially the extreme value of 80.0, raise the mean but not the median.
- c. For urban homes, deleting the smallest ( $x = 4.0$ ) and largest ( $x = 80.0$ ) values gives a trimmed mean of  $\bar{x}_{tr} = 153/9 = 17$  EU/mg. The corresponding trimming percentage is  $100(1/11) \approx 9.1\%$ . The trimmed mean is less than the mean of the entire sample, since the sample was positively skewed. Coincidentally, the median and trimmed mean are equal.

For farm homes, deleting the smallest ( $x = 0.3$ ) and largest ( $x = 21.0$ ) values gives a trimmed mean of  $\bar{x}_{tr} = 107.1/13 = 8.24$  EU/mg. The corresponding trimming percentage is  $100(1/15) \approx 6.7\%$ . The trimmed mean is below, though not far from, the mean and median of the entire sample.

35.

- The sample size is  $n = 15$ .
- a. The sample mean is  $\bar{x} = 18.55/15 = 1.237$   $\mu\text{g/g}$  and the sample median is  $\tilde{x}$  = the 8<sup>th</sup> ordered value = .56  $\mu\text{g/g}$ . These values are very different due to the heavy positive skewness in the data.
- b. A 1/15 trimmed mean is obtained by removing the largest and smallest values and averaging the remaining 13 numbers:  $(.22 + \dots + 3.07)/13 = 1.162$ . Similarly, a 2/15 trimmed mean is the average of the middle 11 values:  $(.25 + \dots + 2.25)/11 = 1.074$ . Since the average of 1/15 and 2/15 is .1 (10%), a 10% trimmed mean is given by the midpoint of these two trimmed means:  $(1.162 + 1.074)/2 = 1.118$   $\mu\text{g/g}$ .

## Chapter 1: Overview and Descriptive Statistics

- c. The median of the data set will remain .56 so long as that's the 8<sup>th</sup> ordered observation. Hence, the value .20 could be increased to as high as .56 without changing the fact that the 8<sup>th</sup> ordered observation is .56. Equivalently, .20 could be increased by as much as .36 without affecting the value of the sample median.

36.

- a. A stem-and leaf display of this data appears below:

32	55	stem: ones
33	49	leaf: tenths
34		
35	6699	
36	34469	
37	03345	
38	9	
39	2347	
40	23	
41		
42	4	

The display is reasonably symmetric, so the mean and median will be close.

- b. The sample mean is  $\bar{x} = 9638/26 = 370.7$  sec, while the sample median is  $\tilde{x} = (369+370)/2 = 369.50$  sec.
- c. The largest value (currently 424) could be increased by any amount. Doing so will not change the fact that the middle two observations are 369 and 370, and hence, the median will not change. However, the value  $x = 424$  cannot be changed to a number less than 370 (a change of  $424 - 370 = 54$ ) since that will change the middle two values.
- d. Expressed in minutes, the mean is  $(370.7 \text{ sec})/(60 \text{ sec}) = 6.18$  min, while the median is 6.16 min.

37.  $\bar{x} = 12.01$ ,  $\tilde{x} = 11.35$ ,  $\bar{x}_{tr(10)} = 11.46$ . The median or the trimmed mean would be better choices than the mean because of the outlier 21.9.

## Chapter 1: Overview and Descriptive Statistics

38.

- a. The reported values are (in increasing order) 110, 115, 120, 120, 125, 130, 130, 135, and 140. Thus the median of the reported values is 125.
- b. 127.6 is reported as 130, so the median is now 130, a very substantial change. When there is rounding or grouping, the median can be highly sensitive to small change.

39.

- a.  $\Sigma x_i = 16.475$  so  $\bar{x} = \frac{16.475}{16} = 1.0297$ ;  $\tilde{x} = \frac{(1.007 + 1.011)}{2} = 1.009$

- b. 1.394 can be decreased until it reaches 1.011 (i.e. by  $1.394 - 1.011 = 0.383$ ), the largest of the 2 middle values. If it is decreased by more than 0.383, the median will change.

40.

$\tilde{x} = 60.8$ ,  $\bar{x}_{tr(25)} = 59.3083$ ,  $\bar{x}_{tr(10)} = 58.3475$ ,  $\bar{x} = 58.54$ . All four measures of center have about the same value.

41.

- a.  $x/n = 7/10 = .7$
- b.  $\bar{x} = .70 =$  the sample proportion of successes
- c. To have  $x/n$  equal .80 requires  $x/25 = .80$  or  $x = (.80)(25) = 20$ . There are 7 successes (S) already, so another  $20 - 7 = 13$  would be required.

42.

- a.  $\bar{y} = \frac{\Sigma y_i}{n} = \frac{\Sigma(x_i + c)}{n} = \frac{\Sigma x_i}{n} + \frac{nc}{n} = \bar{x} + c$

$\tilde{y} =$  the median of  $(x_1 + c, x_2 + c, \dots, x_n + c) =$  median of  $(x_1, x_2, \dots, x_n) + c = \tilde{x} + c$

- b.  $\bar{y} = \frac{\Sigma y_i}{n} = \frac{\Sigma(x_i \cdot c)}{n} = \frac{c \Sigma x_i}{n} = c\bar{x}$

$\tilde{y} =$  the median of  $(cx_1, cx_2, \dots, cx_n) = c \cdot$  the median of  $(x_1, x_2, \dots, x_n) = c\tilde{x}$

43.

The median and certain trimmed means can be calculated, while the mean cannot — the exact values of the “100+” observations are required to calculate the mean.  $\tilde{x} = \frac{(57 + 79)}{2} = 68.0$ ,

$\bar{x}_{tr(20)} = 66.2$ ,  $\bar{x}_{tr(30)} = 67.5$ .



**Section 1.4**

44.

- a. The maximum and minimum values are 182.6 and 180.3, respectively, so the range is  $182.6 - 180.3 = 2.3^\circ\text{C}$ .
- b. Note: If we apply the hint and subtract 180 from each observation, the mean will be 1.41, and the middle two columns will not change. The sum and sum of squares will change, but those effects will cancel and the answer below will stay the same.

	$x_i$	$(x_i - \bar{x})$	$(x_i - \bar{x})^2$	$x_i^2$
	180.5	-0.90833	0.82507	32580.3
	181.7	0.29167	0.08507	33014.9
	180.9	-0.50833	0.25840	32724.8
	181.6	0.19167	0.03674	32978.6
	182.6	1.19167	1.42007	33342.8
	181.6	0.19167	0.03674	32978.6
	181.3	-0.10833	0.01174	32869.7
	182.1	0.69167	0.47840	33160.4
	182.1	0.69167	0.47840	33160.4
	180.3	-1.10833	1.22840	32508.1
	181.7	0.29167	0.08507	33014.9
	180.5	-0.90833	0.82507	32580.3
sums:	2176.9	0	5.769167	394913.6
	$\bar{x} = 181.41$			

$$s^2 = \sum_{i=1}^n (x_i - \bar{x})^2 / (n-1) = 5.769167 / (12 - 1) = 0.52447.$$

c.  $s = \sqrt{0.52447} = 0.724.$

d.  $s^2 = \frac{\sum x^2 - (\sum x)^2 / n}{n-1} = \frac{394913.6 - (2176.9)^2 / 12}{11} = 0.52447.$


45.

- a.  $\bar{x} = 115.58$ . The deviations from the mean are  $116.4 - 115.58 = .82$ ,  $115.9 - 115.58 = .32$ ,  $114.6 - 115.58 = -.98$ ,  $115.2 - 115.58 = -.38$ , and  $115.8 - 115.58 = .22$ . Notice that the deviations from the mean sum to zero, as they should.
- b.  $s^2 = [(.82)^2 + (.32)^2 + (-.98)^2 + (-.38)^2 + (.22)^2] / (5 - 1) = 1.928 / 4 = .482$ , so  $s = .694$ .
- c.  $\sum x_i^2 = 66795.61$ , so  $s^2 = S_{xx} / (n - 1) = (\sum x_i^2 - (\sum x_i)^2 / n) / (n - 1) = (66795.61 - (577.9)^2 / 5) / 4 = 1.928 / 4 = .482$ .
- d. The new sample values are: 16.4 15.9 14.6 15.2 15.8. While the new mean is 15.58, all the deviations are the same as in part (a), and the variance of the transformed data is identical to that of part (b).

## Chapter 1: Overview and Descriptive Statistics

46.

- a.** Since all three distributions are somewhat skewed and two contain outliers (see **d**), medians are the more appropriate central measures. The medians are  
Cooler:  $1.760^{\circ}\text{C}$       Control:  $1.900^{\circ}\text{C}$       Warmer:  $2.305^{\circ}\text{C}$   
The median difference between air and soil temperature increases as the conditions of the minichambers transition from cooler to warmer ( $1.76 < 1.9 < 2.305$ ).
- b.** With the aid of software, the standard deviations are  
Cooler:  $0.401^{\circ}\text{C}$       Control:  $0.531^{\circ}\text{C}$       Warmer:  $0.778^{\circ}\text{C}$   
For the 15 observations under the “cooler” conditions, the typical deviation between an observed temperature difference and the mean temperature difference ( $1.760^{\circ}\text{C}$ ) is roughly  $0.4^{\circ}\text{C}$ . A similar interpretation applies to the other two standard deviations. We see that, according to the standard deviations, variability increases as the conditions of the minichambers transition from cooler to warmer ( $0.401 < 0.531 < 0.778$ ).
- c.** Apply the definitions of lower fourth, upper fourth, and fourth spread to the sorted data within each condition.  
Cooler: lower fourth =  $(1.43 + 1.57)/2 = 1.50$ , upper fourth =  $(1.88 + 1.90)/2 = 1.89$ ,  
 $f_s = 1.89 - 1.50 = 0.39^{\circ}\text{C}$   
Control: lower fourth =  $(1.52 + 1.78)/2 = 1.65$ , upper fourth =  $(2.00 + 2.03)/2 = 2.015$ ,  
 $f_s = 2.015 - 1.65 = 0.365^{\circ}\text{C}$   
Warmer: lower fourth =  $1.91$ , upper fourth =  $2.60$ ,  
 $f_s = 2.60 - 1.91 = 0.69^{\circ}\text{C}$   
The fourth spreads do not communicate the same message as the standard deviations did. The fourth spreads indicate that variability is quite similar under the cooler and control settings, while variability is much larger under the warmer setting. The disparity between the results of **b** and **c** can be partly attributed to the skewness and outliers in the data, which unduly affect the standard deviations.
- d.** As noted earlier, the temperature difference distributions are negatively skewed under all three conditions. The control and warmer data sets each have a single outlier. The boxplots confirm that median temperature difference increases as we transition from cooler to warmer, that cooler and control variability are similar, and that variability under the warmer condition is quite a bit larger.



## Chapter 1: Overview and Descriptive Statistics

47.


- a. From software,  $\tilde{x} = 14.7\%$  and  $\bar{x} = 14.88\%$ . The sample average alcohol content of these 10 wines was 14.88%. Half the wines have alcohol content below 14.7% and half are above 14.7% alcohol.
- b. Working long-hand,  $\sum(x_i - \bar{x})^2 = (14.8 - 14.88)^2 + \dots + (15.0 - 14.88)^2 = 7.536$ . The sample variance equals  $s^2 = \sum(x_i - \bar{x})^2 / (10 - 1) = 0.837$ .
- c. Subtracting 13 from each value will not affect the variance. The 10 new observations are 1.8, 1.5, 3.1, 1.2, 2.9, 0.7, 3.2, 1.6, 0.8, and 2.0. The sum and sum of squares of these 10 new numbers are  $\sum y_i = 18.8$  and  $\sum y_i^2 = 42.88$ . Using the sample variance shortcut, we obtain  $s^2 = [42.88 - (18.8)^2/10]/(10 - 1) = 7.536/9 = 0.837$  again.

48.

- a. Using the sums provided for urban homes,  $S_{xx} = 10,079 - (237.0)^2/11 = 4972.73$ , so  $s = \sqrt{\frac{4972.73}{11-1}} = 22.3$  EU/mg. Similarly for farm homes,  $S_{xx} = 518.836$  and  $s = 6.09$  EU/mg. The endotoxin concentration in an urban home “typically” deviates from the average of 21.55 by about 22.3 EU/mg. The endotoxin concentration in a farm home “typically” deviates from the average of 8.56 by about 6.09 EU/mg. (These interpretations are very loose, especially since the distributions are not symmetric.) In any case, the variability in endotoxin concentration is far greater in urban homes than in farm homes.
- b. The upper and lower fourths of the urban data are 28.0 and 5.5, respectively, for a fourth spread of 22.5 EU/mg. The upper and lower fourths of the farm data are 10.1 and 4, respectively, for a fourth spread of 6.1 EU/mg. Again, we see that the variability in endotoxin concentration is much greater for urban homes than for farm homes.

## Chapter 1: Overview and Descriptive Statistics

- c. Consider the box plots below. The endotoxin concentration in urban homes generally exceeds that in farm homes, whether measured by settled dust or bag dust. The endotoxin concentration in bag dust generally exceeds that of settled dust, both in urban homes and in farm homes. Settled dust in farm homes shows far less variability than any other scenario.



49.

a.  $\sum x_i = 2.75 + \dots + 3.01 = 56.80$ ,  $\sum x_i^2 = 2.75^2 + \dots + 3.01^2 = 197.8040$

b.  $s^2 = \frac{197.8040 - (56.80)^2 / 17}{16} = \frac{8.0252}{16} = .5016$ ,  $s = .708$

50.

From software or from the sums provided,  $\bar{x} = 20179/27 = 747.37$  and

$s = \sqrt{\frac{24657511 - (20179)^2 / 27}{26}} = 606.89$ . The maximum award should be  $\bar{x} + 2s = 747.37 + 2(606.89) = 1961.16$ , or \$1,961,160. This is quite a bit less than the \$3.5 million that was awarded originally.

## Chapter 1: Overview and Descriptive Statistics

51.

- a. From software,  $s^2 = 1264.77 \text{ min}^2$  and  $s = 35.56 \text{ min}$ . Working by hand,  $\Sigma x = 2563$  and  $\Sigma x^2 = 368501$ , so

$$s^2 = \frac{368501 - (2563)^2 / 19}{19 - 1} = 1264.766 \text{ and } s = \sqrt{1264.766} = 35.564$$


- b. If  $y = \text{time in hours}$ , then  $y = cx$  where  $c = \frac{1}{60}$ . So,  $s_y^2 = c^2 s_x^2 = \left(\frac{1}{60}\right)^2 1264.766 = .351 \text{ hr}^2$  and  $s_y = cs_x = \left(\frac{1}{60}\right) 35.564 = .593 \text{ hr}$ .

52.

Let  $d$  denote the fifth deviation. Then  $.3 + .9 + 1.0 + 1.3 + d = 0$  or  $3.5 + d = 0$ , so  $d = -3.5$ . One sample for which these are the deviations is  $x_1 = 3.8$ ,  $x_2 = 4.4$ ,  $x_3 = 4.5$ ,  $x_4 = 4.8$ ,  $x_5 = 0$ . (These were obtained by adding 3.5 to each deviation; adding any other number will produce a different sample with the desired property.)

53.

- a. Using software, for the sample of balanced funds we have  $\bar{x} = 1.121$ ,  $\tilde{x} = 1.050$ ,  $s = 0.536$ ; for the sample of growth funds we have  $\bar{x} = 1.244$ ,  $\tilde{x} = 1.100$ ,  $s = 0.448$ .
- b. The distribution of expense ratios for this sample of balanced funds is fairly symmetric, while the distribution for growth funds is positively skewed. These balanced and growth mutual funds have similar median expense ratios (1.05% and 1.10%, respectively), but expense ratios are generally higher for growth funds. The lone exception is a balanced fund with a 2.86% expense ratio. (There is also one unusually low expense ratio in the sample of balanced funds, at 0.09%.)



## Chapter 1: Overview and Descriptive Statistics

54.


- a. Minitab provides the stem-and-leaf display below. Grip strengths for this sample of 42 individuals are positively skewed, and there is one high outlier at 403 N.

```

6   0  111234
14  0  55668999
(10) 1  0011223444      Stem = 100s
18  1  567889          Leaf = 10s
12  2  01223334
4   2  59
2   3  2
1   3
1   4  0
    
```

- b. Each half has 21 observations. The lower fourth is the 11<sup>th</sup> observation, 87 N. The upper fourth is the 32<sup>nd</sup> observation (11<sup>th</sup> from the top), 210 N. The fourth spread is the difference:  $f_s = 210 - 87 = 123$  N.
- c. min = 16; lower fourth = 87; median = 140; upper fourth = 210; max = 403

The boxplot tells a similar story: grip strengths are slightly positively skewed, with a median of 140N and a fourth spread of 123 N.



- d. inner fences:  $87 - 1.5(123) = -97.5$ ,  $210 + 1.5(123) = 394.5$   
 outer fences:  $87 - 3(123) = -282$ ,  $210 + 3(123) = 579$   
 Grip strength can't be negative, so low outliers are impossible here. A mild high outlier is above 394.5 N and an extreme high outlier is above 579 N. The value 403 N is a mild outlier by this criterion. (Note: some software uses slightly different rules to define outliers — using quartiles and interquartile range — which result in 403 N not being classified as an outlier.)
- e. The fourth spread is unaffected unless that observation drops below the current upper fourth, 210. That's a decrease of  $403 - 210 = 193$  N.

## Chapter 1: Overview and Descriptive Statistics


55.

- a. Lower half of the data set: 325 325 334 339 356 356 359 359 363 364 364 366 369, whose median, and therefore the lower fourth, is 359 (the 7<sup>th</sup> observation in the sorted list).

Upper half of the data set: 370 373 373 374 375 389 392 393 394 397 402 403 424, whose median, and therefore the upper fourth is 392.

So,  $f_s = 392 - 359 = 33$ .


- b. inner fences:  $359 - 1.5(33) = 309.5$ ,  $392 + 1.5(33) = 441.5$   
To be a mild outlier, an observation must be below 309.5 or above 441.5. There are none in this data set. Clearly, then, there are also no extreme outliers.
- c. A boxplot of this data appears below. The distribution of escape times is roughly symmetric with no outliers. Notice the box plot “hides” the fact that the distribution contains two gaps, which can be seen in the stem-and-leaf display.



- d. Not until the value  $x = 424$  is lowered below the upper fourth value of 392 would there be any change in the value of the upper fourth (and, thus, of the fourth spread). That is, the value  $x = 424$  could not be decreased by more than  $424 - 392 = 32$  seconds.


## Chapter 1: Overview and Descriptive Statistics

- 56.** The alcohol content distribution of this sample of 35 port wines is roughly symmetric except for two high outliers. The median alcohol content is 19.2% and the fourth spread is 1.42%. [upper fourth =  $(19.90 + 19.62)/2 = 19.76$ ; lower fourth =  $(18.00 + 18.68)/2 = 18.34$ ] The two outliers were 23.25% and 23.78%, indicating two port wines with unusually high alcohol content.



**57.**

- a.**  $f_s = 216.8 - 196.0 = 20.8$   
 inner fences:  $196 - 1.5(20.8) = 164.6$ ,  $216.8 + 1.5(20.8) = 248$   
 outer fences:  $196 - 3(20.8) = 133.6$ ,  $216.8 + 3(20.8) = 279.2$   
 Of the observations listed, 125.8 is an extreme low outlier and 250.2 is a mild high outlier.
- b.** A boxplot of this data appears below. There is a bit of positive skew to the data but, except for the two outliers identified in part (a), the variation in the data is relatively small.



- 58.** The most noticeable feature of the comparative boxplots is that machine 2's sample values have considerably more variation than does machine 1's sample values. However, a typical value, as measured by the median, seems to be about the same for the two machines. The only outlier that exists is from machine 1.



## Chapter 1: Overview and Descriptive Statistics

59.

- a. If you aren't using software, don't forget to *sort* the data first!


*ED*: median = .4, lower fourth =  $(.1 + .1)/2 = .1$ , upper fourth =  $(2.7 + 2.8)/2 = 2.75$ , fourth spread =  $2.75 - .1 = 2.65$

*Non-ED*: median =  $(1.5 + 1.7)/2 = 1.6$ , lower fourth = .3, upper fourth = 7.9, fourth spread =  $7.9 - .3 = 7.6$ .

- b. *ED*: mild outliers are less than  $.1 - 1.5(2.65) = -3.875$  or greater than  $2.75 + 1.5(2.65) = 6.725$ . Extreme outliers are less than  $.1 - 3(2.65) = -7.85$  or greater than  $2.75 + 3(2.65) = 10.7$ . So, the two largest observations (11.7, 21.0) are extreme outliers and the next two largest values (8.9, 9.2) are mild outliers. There are no outliers at the lower end of the data.


*Non-ED*: mild outliers are less than  $.3 - 1.5(7.6) = -11.1$  or greater than  $7.9 + 1.5(7.6) = 19.3$ . Note that there are no mild outliers in the data, hence there cannot be any extreme outliers, either.

- c. A comparative boxplot appears below. The outliers in the ED data are clearly visible. There is noticeable positive skewness in both samples; the Non-ED sample has more variability than the ED sample; the typical values of the ED sample tend to be smaller than those for the Non-ED sample.



## Chapter 1: Overview and Descriptive Statistics

- 60.** A comparative boxplot (created in Minitab) of this data appears below. The burst strengths for the test nozzle closure welds are quite different from the burst strengths of the production canister nozzle welds. The test welds have much higher burst strengths and the burst strengths are much more variable. The production welds have more consistent burst strength and are consistently lower than the test welds. The production welds data does contain 2 outliers.



- 61.** Outliers occur in the 6a.m. data. The distributions at the other times are fairly symmetric. Variability and the “typical” gasoline-vapor coefficient values increase somewhat until 2p.m., then decrease slightly.

### Supplementary Exercises

62. To simplify the math, subtract the mean from each observation; i.e., let  $y_i = x_i - \bar{x} = x_i - 76831$ . Then  $y_1 = 76683 - 76831 = -148$  and  $y_4 = 77048 - 76831 = 217$ ; by rescaling,  $\bar{y} = \bar{x} - 76831 = 0$ , so  $y_2 + y_3 = -(y_1 + y_4) = -69$ . Also,


$$180 = s = \sqrt{\frac{\sum(x_i - \bar{x})^2}{n-1}} = \sqrt{\frac{\sum y_i^2}{3}} \Rightarrow \sum y_i^2 = 3(180)^2 = 97200$$

so  $y_2^2 + y_3^2 = 97200 - (y_1^2 + y_4^2) = 97200 - ((-148)^2 + (217)^2) = 28207$ .

To solve the equations  $y_2 + y_3 = -69$  and  $y_2^2 + y_3^2 = 28207$ , substitute  $y_3 = -69 - y_2$  into the second equation and use the quadratic formula to solve. This gives  $y_2 = 79.14$  or  $-148.14$  (one is  $y_2$  and one is  $y_3$ ).

Finally,  $x_2$  and  $x_3$  are given by  $y_2 + 76831$  and  $y_3 + 76831$ , or 79,610 and 76,683.

63. As seen in the histogram below, this noise distribution is bimodal (but close to unimodal) with a positive skew and no outliers. The mean noise level is 64.89 dB and the median noise level is 64.7 dB. The fourth spread of the noise measurements is about  $70.4 - 57.8 = 12.6$  dB.




## Chapter 1: Overview and Descriptive Statistics

64.

- a. The sample coefficient of variation is similar for the three highest oil viscosity levels (29.66, 32.30, 27.86) but is much higher for the lowest viscosity (56.01). At low viscosity, it appears that there is much more variation in volume wear relative to the average or “typical” amount of wear.


Viscosity	Wear		
	$\bar{x}$	$s$	$cv$
20.4	40.17	22.50	56.01
30.2	38.83	11.52	29.66
89.4	84.10	27.20	32.30
252.6	27.10	7.55	27.86

- b. Volume wear varies dramatically by viscosity level. At very high viscosity, wear is typically the least and the least variable. Volume wear is actually by far the highest at a “medium” viscosity level and also has the greatest variability at this viscosity level. “Lower” viscosity levels correspond to less wear than a medium level, though there is much greater (relative) variation at a very low viscosity level.



65.


- a. The histogram appears below. A representative value for this data would be around 90 MPa. The histogram is reasonably symmetric, unimodal, and somewhat bell-shaped with a fair amount of variability ( $s \approx 3$  or 4 MPa).




- b. The proportion of the observations that are at least 85 is  $1 - (6+7)/169 = .9231$ . The proportion less than 95 is  $1 - (13+3)/169 = .9053$ .
- c. 90 is the midpoint of the class  $89 < x < 91$ , which contains 43 observations (a relative frequency of  $43/169 = .2544$ ). Therefore about half of this frequency, .1272, should be added to the relative frequencies for the classes to the left of  $x = 90$ . That is, the approximate proportion of observations that are less than 90 is  $.0355 + .0414 + .1006 + .1775 + .1272 = .4822$ .

66.

- a. The initial Se concentrations in the treatment and control groups are not that different. The differences in the box plots below are minor. The median initial Se concentrations for the treatment and control groups are 10.3 mg/L and 10.5 mg/L, respectively, each with fourth spread of about 1.25 mg/L. So, the two groups of cows are comparable at the beginning of the study.




- b. The final Se concentrations of the two groups are extremely different, as evidenced by the box plots below. Whereas the median final Se concentration for the control group is 9.3 mg/L (actually slightly lower than the initial concentration), the median Se concentration in the treatment group is now 103.9 mg/L, nearly a 10-fold increase.



67.

- a. Aortic root diameters for males have mean 3.64 cm, median 3.70 cm, standard deviation 0.269 cm, and fourth spread 0.40. The corresponding values for females are  $\bar{x} = 3.28$  cm,  $\tilde{x} = 3.15$  cm,  $s = 0.478$  cm, and  $f_s = 0.50$  cm. Aortic root diameters are typically (though not universally) somewhat smaller for females than for males, and females show more variability. The distribution for males is negatively skewed, while the distribution for females is positively skewed (see graphs below).



- b. For females ( $n = 10$ ), the 10% trimmed mean is the average of the middle 8 observations:  $\bar{x}_{tr(10)} = 3.24$  cm. For males ( $n = 13$ ), the 1/13 trimmed mean is  $40.2/11 = 3.6545$ , and the 2/13 trimmed mean is  $32.8/9 = 3.6444$ . Interpolating, the 10% trimmed mean is  $\bar{x}_{tr(10)} = 0.7(3.6545) + 0.3(3.6444) = 3.65$  cm. (10% is three-tenths of the way from 1/13 to 2/13).

68.

- a. 
$$\frac{d}{dc} \left\{ \sum (x_i - c)^2 \right\} = \sum \frac{d}{dc} (x_i - c)^2 = -2 \sum (x_i - c) = 0 \Rightarrow \sum (x_i - c) = 0 \Rightarrow$$
  

$$\sum x_i - \sum c = 0 \Rightarrow \sum x_i - nc = 0 \Rightarrow nc = \sum x_i \Rightarrow c = \frac{\sum x_i}{n} = \bar{x}$$
- b. Since  $c = \bar{x}$  minimizes  $\sum (x_i - c)^2$ ,  $\sum (x_i - \bar{x})^2 < \sum (x_i - \mu)^2$ .

## Chapter 1: Overview and Descriptive Statistics

69.

a.

$$\begin{aligned}\bar{y} &= \frac{\sum y_i}{n} = \frac{\sum (ax_i + b)}{n} = \frac{a \sum x_i + \sum b}{n} = \frac{a \sum x_i + nb}{n} = a\bar{x} + b \\ s_y^2 &= \frac{\sum (y_i - \bar{y})^2}{n-1} = \frac{\sum (ax_i + b - (a\bar{x} + b))^2}{n-1} = \frac{\sum (ax_i - a\bar{x})^2}{n-1} \\ &= \frac{a^2 \sum (x_i - \bar{x})^2}{n-1} = a^2 s_x^2\end{aligned}$$

b.


$$x = ^\circ\text{C}, y = ^\circ\text{F}$$

$$\bar{y} = \frac{9}{5}(87.3) + 32 = 189.14^\circ\text{F}$$

$$s_y = \sqrt{s_y^2} = \sqrt{\left(\frac{9}{5}\right)^2 (1.04)^2} = \sqrt{3.5044} = 1.872^\circ\text{F}$$

70.


- a. There is a significant difference in the variability of the two samples. The weight training produced much higher oxygen consumption, on average, than the treadmill exercise, with the median consumptions being approximately 20 and 11 liters, respectively.






## Chapter 1: Overview and Descriptive Statistics

- b.** The differences in oxygen consumption (weight minus treadmill) for the 15 subjects are 3.3, 9.1, 10.4, 9.1, 6.2, 2.5, 2.2, 8.4, 8.7, 14.4, 2.5, -2.8, -0.4, 5.0, and 11.5. The majority of the differences are positive, which suggests that the weight training produced higher oxygen consumption for most subjects than the treadmill did. The median difference is about 6 liters.



**71.**


- a.** The mean, median, and trimmed mean are virtually identical, which suggests symmetry. If there are outliers, they are balanced. The range of values is only 25.5, but half of the values are between 132.95 and 138.25.
- b.** See the comments for (a). In addition, using  $1.5(Q3 - Q1)$  as a yardstick, the two largest and three smallest observations are mild outliers.



## Chapter 1: Overview and Descriptive Statistics

72. A table of summary statistics, a stem and leaf display, and a comparative boxplot are below. The healthy individuals have higher receptor binding measure, on average, than the individuals with PTSD. There is also more variation in the healthy individuals' values. The distribution of values for the healthy is reasonably symmetric, while the distribution for the PTSD individuals is negatively skewed. The box plot indicates that there are no outliers, and confirms the above comments regarding symmetry and skewness.


	PTSD	Healthy		Healthy	PTSD	
Mean	32.92	52.23		1	0	stem = tens
Median	37	51		3	2	leaf = ones
Std Dev	9.93	14.86		9	3	
Min	10	23		7310	4	
Max	46	72		81	5	
				9763	6	
				2	7	



## Chapter 1: Overview and Descriptive Statistics

73. From software,  $\bar{x} = .9255$ ,  $s = .0809$ ;  $\tilde{x} = .93$ ,  $f_s = .1$ . The cadence observations are slightly skewed (mean = .9255 strides/sec, median = .93 strides/sec) and show a small amount of variability (standard deviation = .0809, fourth spread = .1). There are no apparent outliers in the data.

7 8	stem = tenths
8 11556	leaf = hundredths
9 2233335566	
0 0566	




- 74.
- a. The mode is .93. It occurs four times in the data set.
  - b. The *modal category* is the one in which the most observations occur; i.e., the modal category has the highest frequency. In a survey, the modal category is the most popular answer.


## Chapter 1: Overview and Descriptive Statistics

75.

- a. The median is the same (371) in each plot and all three data sets are very symmetric. In addition, all three have the same minimum value (350) and same maximum value (392). Moreover, all three data sets have the same lower (364) and upper quartiles (378). So, all three boxplots will be *identical*. (Slight differences in the boxplots below are due to the way Minitab software interpolates to calculate the quartiles.)



- b. A comparative dotplot is shown below. These graphs show that there are differences in the variability of the three data sets. They also show differences in the way the values are distributed in the three data sets, especially big differences in the presence of gaps and clusters.



- c. The boxplot in (a) is not capable of detecting the differences among the data sets. The primary reason is that boxplots give up some detail in describing data because they use only five summary numbers for comparing data sets.

## Chapter 1: Overview and Descriptive Statistics

- 76.** The measures that are sensitive to outliers are: the mean and the midrange. The mean is sensitive because all values are used in computing it. The midrange is sensitive because it uses only the most extreme values in its computation.

The median, the trimmed mean, and the midfourth are not sensitive to outliers.

The median is the most resistant to outliers because it uses only the middle value (or values) in its computation.

The trimmed mean is somewhat resistant to outliers. The larger the trimming percentage, the more resistant the trimmed mean becomes.

The midfourth, which uses the quartiles, is reasonably resistant to outliers because both quartiles are resistant to outliers.

**77.**


**a.**

```

0  4444444444577888999          leaf = 1.0
1  000111111111124455669999    stem = 0.1
2  1234457
3  11355
4  17
5  3
6
7  67
8  1

HI  10.44, 13.41
    
```

**b.** Since the intervals have unequal width, you must use a *density scale*.



## Chapter 1: Overview and Descriptive Statistics

- c. Representative depths are quite similar for the three types of soils — between 1.5 and 2. Data from the C and CL soils shows much more variability than for the other two types. The boxplots for the first three types show substantial positive skewness both in the middle 50% and overall. The boxplot for the SYCL soil shows negative skewness in the middle 50% and mild positive skewness overall. Finally, there are multiple outliers for the first three types of soils, including extreme outliers.

78.

- a. Since the constant  $\bar{x}$  is subtracted from each  $x$  value to obtain each  $y$  value, and addition or subtraction of a constant doesn't affect variability,  $s_y^2 = s_x^2$  and  $s_y = s_x$ .
- b. Let  $c = 1/s$ , where  $s$  is the sample standard deviation of the  $x$ 's (and also, by part (a), of the  $y$ 's). Then  $z_i = cy_i \Rightarrow s_z^2 = c^2 s_y^2 = (1/s)^2 s^2 = 1$  and  $s_z = 1$ . That is, the "standardized" quantities  $z_1, \dots, z_n$  have a sample variance and standard deviation of 1.

79.

a. 
$$\sum_{i=1}^{n+1} x_i = \sum_{i=1}^n x_i + x_{n+1} = n\bar{x}_n + x_{n+1}, \text{ so } \bar{x}_{n+1} = \frac{1}{n+1} \sum_{i=1}^{n+1} x_i = \frac{n\bar{x}_n + x_{n+1}}{n+1}.$$

- b. In the second line below, we artificially add and subtract  $n\bar{x}_n^2$  to create the term needed for the sample variance:

$$\begin{aligned} ns_{n+1}^2 &= \sum_{i=1}^{n+1} (x_i - \bar{x}_{n+1})^2 = \sum_{i=1}^{n+1} x_i^2 - (n+1)\bar{x}_{n+1}^2 \\ &= \sum_{i=1}^n x_i^2 + x_{n+1}^2 - (n+1)\bar{x}_{n+1}^2 = \left[ \sum_{i=1}^n x_i^2 - n\bar{x}_n^2 \right] + n\bar{x}_n^2 + x_{n+1}^2 - (n+1)\bar{x}_{n+1}^2 \\ &= (n-1)s_n^2 + \left\{ x_{n+1}^2 + n\bar{x}_n^2 - (n+1)\bar{x}_{n+1}^2 \right\} \end{aligned}$$

Substitute the expression for  $\bar{x}_{n+1}$  from part (a) into the expression in braces, and it

simplifies to  $\frac{n}{n+1}(x_{n+1} - \bar{x}_n)^2$ , as desired.


- c. First,  $\bar{x}_{16} = \frac{15(12.58) + 11.8}{16} = \frac{200.5}{16} = 12.53$ . Then, solving (b) for  $s_{n+1}^2$  gives

$$s_{n+1}^2 = \frac{n-1}{n} s_n^2 + \frac{1}{n+1} (x_{n+1} - \bar{x}_n)^2 = \frac{14}{15} (.512)^2 + \frac{1}{16} (11.8 - 12.58)^2 = .238. \text{ Finally, the standard deviation is } s_{16} = \sqrt{.238} = .532.$$

## Chapter 1: Overview and Descriptive Statistics

80.

a.



b. There are 391 observations. The proportion of route lengths less than 20 km is  $(6 + 23 + \dots + 42)/391 = 216/391 = .552$ . The proportion of route lengths at least 30 km is  $(27 + 11 + 2)/391 = 40/391 = .102$ .

c. First compute  $(.90)(391 + 1) = 352.8$ . Thus, the 90<sup>th</sup> percentile should be about the 352<sup>nd</sup> ordered value. The 352<sup>nd</sup> ordered value is the first value in the interval 30–<35. We do not know how the values in that interval are distributed, however, the smallest value (i.e., the 352<sup>nd</sup> value in the data set) cannot be smaller than 30. So, the 90<sup>th</sup> percentile is roughly 30.


d. First compute  $(.50)(391 + 1) = 196$ . Thus the median (50<sup>th</sup> percentile) should be the 196<sup>th</sup> ordered value. The 196<sup>th</sup> observation lies in the interval 18–<20, which includes observations #175 to #216. The 196<sup>th</sup> observation is about in the middle of these. Thus, we would say the median is roughly 19.

81. Assuming that the histogram is unimodal, then there is evidence of positive skewness in the data since the median lies to the left of the mean (for a symmetric distribution, the mean and median would coincide).

For more evidence of skewness, compare the distances of the 5<sup>th</sup> and 95<sup>th</sup> percentiles from the median: median – 5<sup>th</sup> %ile = 500 – 400 = 100, while 95<sup>th</sup> %ile – median = 720 – 500 = 220. Thus, the largest 5% of the values (above the 95th percentile) are further from the median than are the lowest 5%. The same skewness is evident when comparing the 10<sup>th</sup> and 90<sup>th</sup> percentiles to the median, or comparing the maximum and minimum to the median.

82.

- a. There is some evidence of a cyclical pattern.



- b. A complete listing of the smoothed values appears below. To illustrate, with  $\alpha = .1$  we have  $\bar{x}_2 = .1x_2 + .9\bar{x}_1 = (.1)(54) + (.9)(47) = 47.7$ ,  $\bar{x}_3 = .1x_3 + .9\bar{x}_2 = (.1)(53) + (.9)(47.7) = 48.23 \approx 48.2$ , etc. It's clear from the values below that  $\alpha = .1$  gives a smoother series.

$t$	$\bar{x}_t$ for $\alpha = .1$	$\bar{x}_t$ for $\alpha = .5$
1	47.0	47.0
2	47.7	50.5
3	48.2	51.8
4	48.4	50.9
5	48.2	48.4
6	48.0	47.2
7	47.9	47.1
8	48.1	48.6
9	48.4	49.8
10	48.5	49.9
11	48.3	47.9
12	48.6	50.0
13	48.8	50.0
14	48.9	50.0

- c. As seen below,  $\bar{x}_t$  depends on  $x_t$  and all previous values. As  $k$  increases, the coefficient on  $x_{t-k}$  decreases (further back in time implies less weight).

$$\begin{aligned} \bar{x}_t &= \alpha x_t + (1-\alpha)\bar{x}_{t-1} = \alpha x_t + (1-\alpha)[\alpha x_{t-1} + (1-\alpha)\bar{x}_{t-2}] \\ &= \alpha x_t + \alpha(1-\alpha)x_{t-1} + (1-\alpha)^2[\alpha x_{t-2} + (1-\alpha)\bar{x}_{t-3}] = \dots \\ &= \alpha x_t + \alpha(1-\alpha)x_{t-1} + \alpha(1-\alpha)^2 x_{t-2} + \dots + \alpha(1-\alpha)^{t-2} x_2 + (1-\alpha)^{t-1} x_1 \end{aligned}$$

- d. For large  $t$ , the smoothed series is not very sensitive to the initial value  $x_1$ , since the coefficient  $(1-\alpha)^{t-1}$  will be very small.




## Chapter 1: Overview and Descriptive Statistics

83.

- a. When there is perfect symmetry, the smallest observation  $y_1$  and the largest observation  $y_n$  will be equidistant from the median, so  $y_n - \tilde{x} = \tilde{x} - y_1$ . Similarly, the second-smallest and second-largest will be equidistant from the median, so  $y_{n-1} - \tilde{x} = \tilde{x} - y_2$ , and so on. Thus, the first and second numbers in each pair will be equal, so that each point in the plot will fall exactly on the  $45^\circ$  line.

When the data is positively skewed,  $y_n$  will be much further from the median than is  $y_1$ , so  $y_n - \tilde{x}$  will considerably exceed  $\tilde{x} - y_1$  and the point  $(y_n - \tilde{x}, \tilde{x} - y_1)$  will fall considerably below the  $45^\circ$  line, as will the other points in the plot.

- b. The median of these  $n = 26$  observations is 221.6 (the midpoint of the 13<sup>th</sup> and 14<sup>th</sup> ordered values). The first point in the plot is  $(2745.6 - 221.6, 221.6 - 4.1) = (2524.0, 217.5)$ . The others are: (1476.2, 213.9), (1434.4, 204.1), (756.4, 190.2), (481.8, 188.9), (267.5, 181.0), (208.4, 129.2), (112.5, 106.3), (81.2, 103.3), (53.1, 102.6), (53.1, 92.0), (33.4, 23.0), and (20.9, 20.9). The first number in each of the first seven pairs greatly exceeds the second number, so each of those points falls well below the  $45^\circ$  line. A substantial positive skew (stretched upper tail) is indicated.



## Chapter 1: Overview and Descriptive Statistics

- 84.** As suggested in the hint, split the sum into two “halves” corresponding to the lowest  $n/2$  observations and the highest  $n/2$  observations (we’ll use  $L$  and  $U$  to denote these).

$$\begin{aligned} \sum |x_i - \tilde{x}| &= \sum_L |x_i - \tilde{x}| + \sum_U |x_i - \tilde{x}| \\ &= \sum_L (\tilde{x} - x_i) + \sum_U (x_i - \tilde{x}) \\ &= \sum_L \tilde{x} - \sum_L x_i + \sum_U x_i - \sum_U \tilde{x} \end{aligned}$$

Each of these four sums covers exactly  $n/2$  terms. The first and fourth sums are, therefore, both equal to  $(n/2)\tilde{x}$ ; these cancel. The inner two sums may be re-written in terms of averages:

$$\begin{aligned} \sum |x_i - \tilde{x}| &= \sum_L \tilde{x} - \sum_L x_i + \sum_U x_i - \sum_U \tilde{x} = -\sum_L x_i + \sum_U x_i \\ &= -(n/2)\bar{x}_L + (n/2)\bar{x}_U \quad \Rightarrow \\ \sum |x_i - \tilde{x}| / n &= (\bar{x}_U - \bar{x}_L) / 2 \end{aligned}$$

When  $n$  is odd, the middle (ordered) value is exactly  $\tilde{x}$ . Using  $L$  and  $U$  to denote the lowest  $(n-1)/2$  observations and largest  $(n-1)/2$  observations, respectively, we may write

$$\sum |x_i - \tilde{x}| = \sum_L |x_i - \tilde{x}| + 0 + \sum_U |x_i - \tilde{x}|, \text{ where the } 0 \text{ comes from the middle (ordered) value,}$$

viz.  $|\tilde{x} - \tilde{x}| = 0$ . The rest of the derivation proceeds exactly as before, except that the surviving sums each have  $(n-1)/2$  terms in them, not  $n/2$ . As a result, for  $n$  odd we have

$$\sum |x_i - \tilde{x}| / (n-1) = (\bar{x}_U - \bar{x}_L) / 2.$$

## CHAPTER 2

### Section 2.1

1.

- a.  $S = \{1324, 1342, 1423, 1432, 2314, 2341, 2413, 2431, 3124, 3142, 4123, 4132, 3214, 3241, 4213, 4231\}$ .
- b. Event  $A$  contains the outcomes where 1 is first in the list:  
 $A = \{1324, 1342, 1423, 1432\}$ .
- c. Event  $B$  contains the outcomes where 2 is first or second:  
 $B = \{2314, 2341, 2413, 2431, 3214, 3241, 4213, 4231\}$ .
- d. The event  $A \cup B$  contains the outcomes in  $A$  or  $B$  or both:  
 $A \cup B = \{1324, 1342, 1423, 1432, 2314, 2341, 2413, 2431, 3214, 3241, 4213, 4231\}$ .  
 $A \cap B = \emptyset$ , since 1 and 2 can't both get into the championship game.  
 $A' = S - A = \{2314, 2341, 2413, 2431, 3124, 3142, 4123, 4132, 3214, 3241, 4213, 4231\}$ .

2.

- a.  $A = \{RRR, LLL, SSS\}$ .
- b.  $B = \{RLS, RSL, LRS, LSR, SRL, SLR\}$ .
- c.  $C = \{RRL, RRS, RLR, RSR, LRR, SRR\}$ .
- d.  $D = \{RRL, RRS, RLR, RSR, LRR, SRR, LLR, LLS, LRL, LSL, RLL, SLL, SSR, SSL, SRS, SLS, RSS, LSS\}$
- e. Event  $D'$  contains outcomes where either all cars go the same direction or they all go different directions:  
 $D' = \{RRR, LLL, SSS, RLS, RSL, LRS, LSR, SRL, SLR\}$ .  
Because event  $D$  totally encloses event  $C$  (see the lists above), the compound event  $C \cup D$  is just event  $D$ :  
 $C \cup D = D = \{RRL, RRS, RLR, RSR, LRR, SRR, LLR, LLS, LRL, LSL, RLL, SLL, SSR, SSL, SRS, SLS, RSS, LSS\}$ .  
Using similar reasoning, we see that the compound event  $C \cap D$  is just event  $C$ :  
 $C \cap D = C = \{RRL, RRS, RLR, RSR, LRR, SRR\}$ .

## Chapter 2: Probability

3.

- a.  $A = \{SSF, SFS, FSS\}$ .
- b.  $B = \{SSS, SSF, SFS, FSS\}$ .
- c. For event  $C$  to occur, the system must have component 1 working ( $S$  in the first position), then at least one of the other two components must work (at least one  $S$  in the second and third positions):  $C = \{SSS, SSF, SFS\}$ .
- d.  $C' = \{SFF, FSS, FSF, FFS, FFF\}$ .  
 $A \cup C = \{SSS, SSF, SFS, FSS\}$ .  
 $A \cap C = \{SSF, SFS\}$ .  
 $B \cup C = \{SSS, SSF, SFS, FSS\}$ . Notice that  $B$  contains  $C$ , so  $B \cup C = B$ .  
 $B \cap C = \{SSS, SSF, SFS\}$ . Since  $B$  contains  $C$ ,  $B \cap C = C$ .

4.

- a. The  $2^4 = 16$  possible outcomes have been numbered here for later reference.

Outcome	Home Mortgage Number			
	1	2	3	4
1	<i>F</i>	<i>F</i>	<i>F</i>	<i>F</i>
2	<i>F</i>	<i>F</i>	<i>F</i>	<i>V</i>
3	<i>F</i>	<i>F</i>	<i>V</i>	<i>F</i>
4	<i>F</i>	<i>F</i>	<i>V</i>	<i>V</i>
5	<i>F</i>	<i>V</i>	<i>F</i>	<i>F</i>
6	<i>F</i>	<i>V</i>	<i>F</i>	<i>V</i>
7	<i>F</i>	<i>V</i>	<i>V</i>	<i>F</i>
8	<i>F</i>	<i>V</i>	<i>V</i>	<i>V</i>
9	<i>V</i>	<i>F</i>	<i>F</i>	<i>F</i>
10	<i>V</i>	<i>F</i>	<i>F</i>	<i>V</i>
11	<i>V</i>	<i>F</i>	<i>V</i>	<i>F</i>
12	<i>V</i>	<i>F</i>	<i>V</i>	<i>V</i>
13	<i>V</i>	<i>V</i>	<i>F</i>	<i>F</i>
14	<i>V</i>	<i>V</i>	<i>F</i>	<i>V</i>
15	<i>V</i>	<i>V</i>	<i>V</i>	<i>F</i>
16	<i>V</i>	<i>V</i>	<i>V</i>	<i>V</i>

- b. Outcome numbers 2, 3, 5, 9 above.
- c. Outcome numbers 1, 16 above.
- d. Outcome numbers 1, 2, 3, 5, 9 above.
- e. In words, the union of (c) and (d) is the event that either all of the mortgages are variable, or that at most one of them is variable-rate: outcomes 1, 2, 3, 5, 9, 16. The intersection of (c) and (d) is the event that all of the mortgages are fixed-rate: outcome 1.
- f. The union of (b) and (c) is the event that either exactly three are fixed, or that all four are the same: outcomes 1, 2, 3, 5, 9, 16. The intersection of (b) and (c) is the event that exactly three are fixed and all four are the same type. This cannot happen (the events have no outcomes in common), so the intersection of (b) and (c) is  $\emptyset$ .

## Chapter 2: Probability

5.

- a. The  $3^3 = 27$  possible outcomes are numbered below for later reference.

Outcome Number	Outcome	Outcome Number	Outcome
1	111	15	223
2	112	16	231
3	113	17	232
4	121	18	233
5	122	19	311
6	123	20	312
7	131	21	313
8	132	22	321
9	133	23	322
10	211	24	323
11	212	25	331
12	213	26	332
13	221	27	333
14	222		

- b. Outcome numbers 1, 14, 27 above.  
 c. Outcome numbers 6, 8, 12, 16, 20, 22 above.  
 d. Outcome numbers 1, 3, 7, 9, 19, 21, 25, 27 above.

6.


- a.  $S = \{123, 124, 125, 213, 214, 215, 13, 14, 15, 23, 24, 25, 3, 4, 5\}$ .  
 b.  $A = \{3, 4, 5\}$ .  
 c.  $B = \{125, 215, 15, 25, 5\}$ .  
 d.  $C = \{23, 24, 25, 3, 4, 5\}$ .

7.


- a.  $S = \{BBBAAA, BBABAAA, BBAABAA, BBAAABA, BBAAAAB, BABBAAA, BABABAA, BABAABA, BABAAAAB, BAABBAA, BAABABA, BAABAAB, BAAABBA, BAAABAB, BAAAABB, ABBBAAA, ABBABAA, ABBAABA, ABBAAAAB, ABABBAA, ABABABA, ABABAAB, ABAABBA, ABAABAB, ABAABBB, AAABBAA, AABBBAA, AABBBAA, AABBAAB, AABABBA, AABABAB, AABAABB, AAABBBA, AAABBAB, AAABABB, AAAABBB\}$ .  
 b.  $AAAABBB, AAABABB, AAABBAB, AABAABB, AABABAB$ .

## Chapter 2: Probability


8.




a.  $A_1 \cup A_2 \cup A_3$




b.  $A_1 \cap A_2 \cap A_3$



c.  $A_1 \cap A_2' \cap A_3'$



d.  $(A_1 \cap A_2' \cap A_3') \cup (A_1' \cap A_2 \cap A_3') \cup (A_1' \cap A_2' \cap A_3)$





e.  $A_1 \cup (A_2 \cap A_3)$


## Chapter 2: Probability

9.

- a. In the diagram on the left, the shaded area is  $(A \cup B)'$ . On the right, the shaded area is  $A'$ , the striped area is  $B'$ , and the intersection  $A' \cap B'$  occurs where there is both shading and stripes. These two diagrams display the same area.



- b. In the diagram below, the shaded area represents  $(A \cap B)'$ . Using the right-hand diagram from (a), the union of  $A'$  and  $B'$  is represented by the areas that have either shading or stripes (or both). Both of the diagrams display the same area.



10.

- a. Many examples exist; e.g.,  $A = \{\text{Chevy, Buick}\}$ ,  $B = \{\text{Ford, Lincoln}\}$ ,  $C = \{\text{Toyota}\}$  are three mutually exclusive events.
- b. No. Let  $E = \{\text{Chevy, Buick}\}$ ,  $F = \{\text{Buick, Ford}\}$ ,  $G = \{\text{Toyota}\}$ . These events are not mutually exclusive ( $E$  and  $F$  have an outcome in common), yet there is no outcome common to all three events.

**Section 2.2**

11.

- a. .07.
- b.  $.15 + .10 + .05 = .30$ .
- c. Let  $A$  = the selected individual owns shares in a stock fund. Then  $P(A) = .18 + .25 = .43$ . The desired probability, that a selected customer does not shares in a stock fund, equals  $P(A') = 1 - P(A) = 1 - .43 = .57$ . This could also be calculated by adding the probabilities for all the funds that are not stocks.

12.

- a. No, this is not possible. Since event  $A \cap B$  is contained within event  $B$ , it must be the case that  $P(A \cap B) \leq P(B)$ . However,  $.5 > .4$ .
- b. By the addition rule,  $P(A \cup B) = .5 + .4 - .3 = .6$ .
- c.  $P(\text{neither } A \text{ nor } B) = P(A' \cap B') = P((A \cup B)') = 1 - P(A \cup B) = 1 - .6 = .4$ .
- d. The event of interest is  $A \cap B'$ ; from a Venn diagram, we see  $P(A \cap B') = P(A) - P(A \cap B) = .5 - .3 = .2$ .
- e. From a Venn diagram, we see that the probability of interest is  $P(\text{exactly one}) = P(\text{at least one}) - P(\text{both}) = P(A \cup B) - P(A \cap B) = .6 - .3 = .3$ .


13.

- a.  $A_1 \cup A_2$  = “awarded either #1 or #2 (or both)”: from the addition rule,  $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) = .22 + .25 - .11 = .36$ .
- b.  $A_1' \cap A_2'$  = “awarded neither #1 or #2”: using the hint and part (a),  $P(A_1' \cap A_2') = P((A_1 \cup A_2)') = 1 - P(A_1 \cup A_2) = 1 - .36 = .64$ .
- c.  $A_1 \cup A_2 \cup A_3$  = “awarded at least one of these three projects”: using the addition rule for 3 events,  $P(A_1 \cup A_2 \cup A_3) = P(A_1) + P(A_2) + P(A_3) - P(A_1 \cap A_2) - P(A_1 \cap A_3) - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3) = .22 + .25 + .28 - .11 - .05 - .07 + .01 = .53$ .
- d.  $A_1' \cap A_2' \cap A_3'$  = “awarded none of the three projects”:  $P(A_1' \cap A_2' \cap A_3') = 1 - P(\text{awarded at least one}) = 1 - .53 = .47$ .




## Chapter 2: Probability


- e.  $A_1' \cap A_2' \cap A_3 =$  “awarded #3 but neither #1 nor #2”: from a Venn diagram,  
 $P(A_1' \cap A_2' \cap A_3) = P(A_3) - P(A_1 \cap A_3) - P(A_2 \cap A_3) + P(A_1 \cap A_2 \cap A_3) =$   
 $.28 - .05 - .07 + .01 = .17$ . The last term addresses the “double counting” of the two subtractions.



- f.  $(A_1' \cap A_2') \cup A_3 =$  “awarded neither of #1 and #2, or awarded #3”: from a Venn diagram,  
 $P((A_1' \cap A_2') \cup A_3) = P(\text{none awarded}) + P(A_3) = .47$  (from **d**)  $+ .28 = .75$ .



Alternatively, answers to **a-f** can be obtained from probabilities on the accompanying Venn diagram:



14. Let  $A =$  an adult consumes coffee and  $B =$  an adult consumes carbonated soda. We’re told that  $P(A) = .55$ ,  $P(B) = .45$ , and  $P(A \cup B) = .70$ .
- a. The addition rule says  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ , so  $.70 = .55 + .45 - P(A \cap B)$  or  $P(A \cap B) = .55 + .45 - .70 = .30$ .
- b. There are two ways to read this question. We can read “does not (consume at least one),” which means the adult consumes neither beverage. The probability is then  $P(\text{neither } A \text{ nor } B) = P(A' \cap B') = 1 - P(A \cup B) = 1 - .70 = .30$ .

The other reading, and this is presumably the intent, is “there is at least one beverage the adult does not consume, i.e.  $A' \cup B'$ ”. The probability is  $P(A' \cup B') = 1 - P(A \cap B) = 1 - .30$  from **a** = .70. (It’s just a coincidence this equals  $P(A \cup B)$ .)

Both of these approaches use *deMorgan’s laws*, which say that  $P(A' \cap B') = 1 - P(A \cup B)$  and  $P(A' \cup B') = 1 - P(A \cap B)$ .

## Chapter 2: Probability

15.

- a. Let  $E$  be the event that at most one purchases an electric dryer. Then  $E'$  is the event that at least two purchase electric dryers, and  $P(E') = 1 - P(E) = 1 - .428 = .572$ .
- b. Let  $A$  be the event that all five purchase gas, and let  $B$  be the event that all five purchase electric. All other possible outcomes are those in which at least one of each type of clothes dryer is purchased. Thus, the desired probability is  $1 - [P(A) - P(B)] = 1 - [.116 + .005] = .879$ .

16.

- a. There are six simple events, corresponding to the outcomes  $CDP$ ,  $CPD$ ,  $DCP$ ,  $DPC$ ,  $PCD$ , and  $PDC$ . Since the same cola is in every glass, these six outcomes are equally likely to occur, and the probability assigned to each is  $\frac{1}{6}$ .
- b.  $P(C \text{ ranked first}) = P(\{CPD, CDP\}) = \frac{1}{6} + \frac{1}{6} = \frac{2}{6} = .333$ .
- c.  $P(C \text{ ranked first and } D \text{ last}) = P(\{CPD\}) = \frac{1}{6}$ .

17.

- a. The probabilities do not add to 1 because there are other software packages besides SPSS and SAS for which requests could be made.
- b.  $P(A') = 1 - P(A) = 1 - .30 = .70$ .
- c. Since  $A$  and  $B$  are mutually exclusive events,  $P(A \cup B) = P(A) + P(B) = .30 + .50 = .80$ .
- d. By deMorgan's law,  $P(A' \cap B') = P((A \cup B)') = 1 - P(A \cup B) = 1 - .80 = .20$ . In this example, deMorgan's law says the event "neither  $A$  nor  $B$ " is the complement of the event "either  $A$  or  $B$ ." (That's true regardless of whether they're mutually exclusive.)

18. The only reason we'd need at least two selections to find a \$10 bill is if the first selection was not a \$10 bill. There are  $4 + 6 = 10$  non-\$10 bills out of  $5 + 4 + 6 = 15$  bills in the wallet, so the probability of this event is simply  $10/15$ , or  $2/3$ .

19. Let  $A$  be that the selected joint was found defective by inspector  $A$ , so  $P(A) = \frac{724}{10,000}$ . Let  $B$  be analogous for inspector  $B$ , so  $P(B) = \frac{751}{10,000}$ . The event "at least one of the inspectors judged a joint to be defective is  $A \cup B$ , so  $P(A \cup B) = \frac{1159}{10,000}$ .


- a. By deMorgan's law,  $P(\text{neither } A \text{ nor } B) = P(A' \cap B') = 1 - P(A \cup B) = 1 - \frac{1159}{10,000} = \frac{8841}{10,000} = .8841$ .
- b. The desired event is  $B \cap A'$ . From a Venn diagram, we see that  $P(B \cap A') = P(B) - P(A \cap B)$ . From the addition rule,  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$  gives  $P(A \cap B) = .0724 + .0751 - .1159 = .0316$ . Finally,  $P(B \cap A') = P(B) - P(A \cap B) = .0751 - .0316 = .0435$ .

## Chapter 2: Probability

- 20.**
- a. Let  $S_1, S_2$  and  $S_3$  represent day, swing, and night shifts, respectively. Let  $C_1$  and  $C_2$  represent unsafe conditions and unrelated to conditions, respectively. Then the simple events are  $S_1C_1, S_1C_2, S_2C_1, S_2C_2, S_3C_1, S_3C_2$ .
  - b.  $P(C_1) = P(\{S_1C_1, S_2C_1, S_3C_1\}) = .10 + .08 + .05 = .23$ .
  - c.  $P(S'_1) = 1 - P(\{S_1C_1, S_1C_2\}) = 1 - (.10 + .35) = .55$ .
- 21.** In what follows, the first letter refers to the auto deductible and the second letter refers to the homeowner's deductible.
- a.  $P(MH) = .10$ .
  - b.  $P(\text{low auto deductible}) = P(\{LN, LL, LM, LH\}) = .04 + .06 + .05 + .03 = .18$ . Following a similar pattern,  $P(\text{low homeowner's deductible}) = .06 + .10 + .03 = .19$ .
  - c.  $P(\text{same deductible for both}) = P(\{LL, MM, HH\}) = .06 + .20 + .15 = .41$ .
  - d.  $P(\text{deductibles are different}) = 1 - P(\text{same deductible for both}) = 1 - .41 = .59$ .
  - e.  $P(\text{at least one low deductible}) = P(\{LN, LL, LM, LH, ML, HL\}) = .04 + .06 + .05 + .03 + .10 + .03 = .31$ .
  - f.  $P(\text{neither deductible is low}) = 1 - P(\text{at least one low deductible}) = 1 - .31 = .69$ .
- 22.** Let  $A$  = motorist must stop at first signal and  $B$  = motorist must stop at second signal. We're told that  $P(A) = .4, P(B) = .5$ , and  $P(A \cup B) = .6$ .
- a. From the addition rule,  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ , so  $.6 = .4 + .5 - P(A \cap B)$ , from which  $P(A \cap B) = .4 + .5 - .6 = .3$ .
  - b. From a Venn diagram,  $P(A \cap B') = P(A) - P(A \cap B) = .4 - .3 = .1$ .
  - c. From a Venn diagram,  $P(\text{stop at exactly one signal}) = P(A \cup B) - P(A \cap B) = .6 - .3 = .3$ . Or,  $P(\text{stop at exactly one signal}) = P([A \cap B'] \cup [A' \cap B]) = P(A \cap B') + P(A' \cap B) = [P(A) - P(A \cap B)] + [P(B) - P(A \cap B)] = [.4 - .3] + [.5 - .3] = .1 + .2 = .3$ .
- 23.** Assume that the computers are numbered 1-6 as described and that computers 1 and 2 are the two laptops. There are 15 possible outcomes: (1,2) (1,3) (1,4) (1,5) (1,6) (2,3) (2,4) (2,5) (2,6) (3,4) (3,5) (3,6) (4,5) (4,6) and (5,6).
- a.  $P(\text{both are laptops}) = P(\{(1,2)\}) = \frac{1}{15} = .067$ .
  - b.  $P(\text{both are desktops}) = P(\{(3,4) (3,5) (3,6) (4,5) (4,6) (5,6)\}) = \frac{6}{15} = .40$ .
  - c.  $P(\text{at least one desktop}) = 1 - P(\text{no desktops}) = 1 - P(\text{both are laptops}) = 1 - .067 = .933$ .
  - d.  $P(\text{at least one of each type}) = 1 - P(\text{both are the same}) = 1 - [P(\text{both are laptops}) + P(\text{both are desktops})] = 1 - [.067 + .40] = .533$ .

## Chapter 2: Probability


- 24.** Since  $A$  is contained in  $B$ , we may write  $B = A \cup (B \cap A')$ , the union of two mutually exclusive events. (See diagram for these two events.) Apply the axioms:  
 $P(B) = P(A \cup (B \cap A')) = P(A) + P(B \cap A')$  by Axiom 3. Then, since  $P(B \cap A') \geq 0$  by Axiom 1,  $P(B) = P(A) + P(B \cap A') \geq P(A) + 0 = P(A)$ . This proves the statement.



For general events  $A$  and  $B$  (i.e., not necessarily those in the diagram), it's always the case that  $A \cap B$  is contained in  $A$  as well as in  $B$ , while  $A$  and  $B$  are both contained in  $A \cup B$ . Therefore,  $P(A \cap B) \leq P(A) \leq P(A \cup B)$  and  $P(A \cap B) \leq P(B) \leq P(A \cup B)$ .

- 25.** By rearranging the addition rule,  $P(A \cap B) = P(A) + P(B) - P(A \cup B) = .40 + .55 - .63 = .32$ . By the same method,  $P(A \cap C) = .40 + .70 - .77 = .33$  and  $P(B \cap C) = .55 + .70 - .80 = .45$ . Finally, rearranging the addition rule for 3 events gives  
 $P(A \cap B \cap C) = P(A \cup B \cup C) - P(A) - P(B) - P(C) + P(A \cap B) + P(A \cap C) + P(B \cap C) = .85 - .40 - .55 - .70 + .32 + .33 + .45 = .30$ .

These probabilities are reflected in the Venn diagram below.



- a.  $P(A \cup B \cup C) = .85$ , as given.
- b.  $P(\text{none selected}) = 1 - P(\text{at least one selected}) = 1 - P(A \cup B \cup C) = 1 - .85 = .15$ .
- c. From the Venn diagram,  $P(\text{only automatic transmission selected}) = .22$ .
- d. From the Venn diagram,  $P(\text{exactly one of the three}) = .05 + .08 + .22 = .35$ .



## Section 2.3

29.

- a. There are 26 letters, so allowing repeats there are  $(26)(26) = (26)^2 = 676$  possible 2-letter domain names. Add in the 10 digits, and there are 36 characters available, so allowing repeats there are  $(36)(36) = (36)^2 = 1296$  possible 2-character domain names.
- b. By the same logic as part a, the answers are  $(26)^3 = 17,576$  and  $(36)^3 = 46,656$ .
- c. Continuing,  $(26)^4 = 456,976$ ;  $(36)^4 = 1,679,616$ .
- d.  $P(4\text{-character sequence is already owned}) = 1 - P(4\text{-character sequence still available}) = 1 - 97,786/(36)^4 = .942$ .

30.

- a. Because order is important, we'll use  $P_{3,8} = (8)(7)(6) = 336$ .
- b. Order doesn't matter here, so we use  $\binom{30}{6} = 593,775$ .
- c. The number of ways to choose 2 zinfandels from the 8 available is  $\binom{8}{2}$ . Similarly, the number of ways to choose the merlots and cabernets are  $\binom{10}{2}$  and  $\binom{12}{2}$ , respectively. Hence, the total number of options (using the Fundamental Counting Principle) equals  $\binom{8}{2}\binom{10}{2}\binom{12}{2} = (28)(45)(66) = 83,160$ .
- d. The numerator comes from part c and the denominator from part b:  $\frac{83,160}{593,775} = .140$ .
- e. We use the same denominator as in part d. The number of ways to choose all zinfandel is  $\binom{8}{6}$ , with similar answers for all merlot and all cabernet. Since these are disjoint events,  $P(\text{all same}) = P(\text{all zin}) +$

$$P(\text{all merlot}) + P(\text{all cab}) = \frac{\binom{8}{6} + \binom{10}{6} + \binom{12}{6}}{\binom{30}{6}} = \frac{1162}{593,775} = .002.$$

31.

- a. Use the Fundamental Counting Principle:  $(9)(5) = 45$ .
- b. By the same reasoning, there are  $(9)(5)(32) = 1440$  such sequences, so such a policy could be carried out for 1440 successive nights, or almost 4 years, without repeating exactly the same program.

## Chapter 2: Probability

32.

- a. Since there are 5 receivers, 4 CD players, 3 speakers, and 4 turntables, the total number of possible selections is  $(5)(4)(3)(4) = 240$ .
- b. We now only have 1 choice for the receiver and CD player:  $(1)(1)(3)(4) = 12$ .
- c. Eliminating Sony leaves 4, 3, 3, and 3 choices for the four pieces of equipment, respectively:  $(4)(3)(3)(3) = 108$ .
- d. From **a**, there are 240 possible configurations. From **c**, 108 of them involve zero Sony products. So, the number of configurations with at least one Sony product is  $240 - 108 = 132$ .
- e. Assuming all 240 arrangements are equally likely,  $P(\text{at least one Sony}) = \frac{132}{240} = .55$ .

Next,  $P(\text{exactly one component Sony}) = P(\text{only the receiver is Sony}) + P(\text{only the CD player is Sony}) + P(\text{only the turntable is Sony})$ . Counting from the available options gives

$$P(\text{exactly one component Sony}) = \frac{(1)(3)(3)(3) + (4)(1)(3)(3) + (4)(3)(3)(1)}{240} = \frac{99}{240} = .413.$$

33.

- a. Since there are 15 players and 9 positions, and order matters in a line-up (catcher, pitcher, shortstop, etc. are different positions), the number of possibilities is  $P_{9,15} = (15)(14)\dots(7)$  or  $15!/(15-9)! = 1,816,214,440$ .
- b. For each of the starting line-ups in part (a), there are 9! possible batting orders. So, multiply the answer from (a) by 9! to get  $(1,816,214,440)(362,880) = 659,067,881,472,000$ .
- c. Order still matters: There are  $P_{3,5} = 60$  ways to choose three left-handers for the outfield and  $P_{6,10} = 151,200$  ways to choose six right-handers for the other positions. The total number of possibilities is  $= (60)(151,200) = 9,072,000$ .

34.

- a. Since order doesn't matter, the number of ways to randomly select 5 keyboards from the 25 available is  $\binom{25}{5} = 53,130$ .
- b. Sample in two stages. First, there are 6 keyboards with an electrical defect, so the number of ways to select exactly 2 of them is  $\binom{6}{2}$ . Next, the remaining  $5 - 2 = 3$  keyboards in the sample must have mechanical defects; as there are 19 such keyboards, the number of ways to randomly select 3 is  $\binom{19}{3}$ . So, the number of ways to achieve both of these in the sample of 5 is the product of these two counting numbers:  $\binom{6}{2} \binom{19}{3} = (15)(969) = 14,535$ .

## Chapter 2: Probability

- c. Following the analogy from **b**, the number of samples with exactly 4 mechanical defects is  $\binom{19}{4}\binom{6}{1}$ , and the number with exactly 5 mechanical defects is  $\binom{19}{5}\binom{6}{0}$ . So, the number of samples with at least 4 mechanical defects is  $\binom{19}{4}\binom{6}{1} + \binom{19}{5}\binom{6}{0}$ , and the probability of this event is
- $$\frac{\binom{19}{4}\binom{6}{1} + \binom{19}{5}\binom{6}{0}}{\binom{25}{5}} = \frac{34,884}{53,130} = .657. \text{ (The denominator comes from a.)}$$

**35.**

- a. There are  $\binom{10}{5} = 252$  ways to select 5 workers from the day shift. In other words, of all the ways to select 5 workers from among the 24 available, 252 such selections result in 5 day-shift workers. Since the grand total number of possible selections is  $\binom{24}{5} = 42504$ , the probability of randomly selecting 5 day-shift workers (and, hence, no swing or graveyard workers) is  $252/42504 = .00593$ .
- b. Similar to **a**, there are  $\binom{8}{5} = 56$  ways to select 5 swing-shift workers and  $\binom{6}{5} = 6$  ways to select 5 graveyard-shift workers. So, there are  $252 + 56 + 6 = 314$  ways to pick 5 workers from the same shift. The probability of this randomly occurring is  $314/42504 = .00739$ .
- c.  $P(\text{at least two shifts represented}) = 1 - P(\text{all from same shift}) = 1 - .00739 = .99261$ .
- d. There are several ways to approach this question. For example, let  $A_1 = \text{"day shift is unrepresented,"}$   $A_2 = \text{"swing shift is unrepresented,"}$  and  $A_3 = \text{"graveyard shift is unrepresented."}$  Then we want  $P(A_1 \cup A_2 \cup A_3)$ .

$$N(A_1) = N(\text{day shift unrepresented}) = N(\text{all from swing/graveyard}) = \binom{8+6}{5} = 2002,$$

since there are  $8 + 6 = 14$  total employees in the swing and graveyard shifts. Similarly,

$$N(A_2) = \binom{10+6}{5} = 4368 \text{ and } N(A_3) = \binom{10+8}{5} = 8568. \text{ Next, } N(A_1 \cap A_2) = N(\text{all from graveyard}) = 6$$

from **b**. Similarly,  $N(A_1 \cap A_3) = 56$  and  $N(A_2 \cap A_3) = 252$ . Finally,  $N(A_1 \cap A_2 \cap A_3) = 0$ , since at least one shift must be represented. Now, apply the addition rule for 3 events:

$$P(A_1 \cup A_2 \cup A_3) = \frac{2002 + 4368 + 8568 - 6 - 56 - 252 + 0}{42504} = \frac{14624}{42504} = .3441.$$

- 36.** There are  $\binom{5}{2} = 10$  possible ways to select the positions for  $B$ 's votes:  $BBAAA, BABAA, BAABA, BAAAB, ABBA, ABABA, ABAAB, AABBA, AABAB, \text{ and } AAABB$ . Only the last two have  $A$  ahead of  $B$  throughout the vote count. Since the outcomes are equally likely, the desired probability is  $2/10 = .20$ .



## Chapter 2: Probability

37.

- a. By the Fundamental Counting Principle, with  $n_1 = 3$ ,  $n_2 = 4$ , and  $n_3 = 5$ , there are  $(3)(4)(5) = 60$  runs.
- b. With  $n_1 = 1$  (just one temperature),  $n_2 = 2$ , and  $n_3 = 5$ , there are  $(1)(2)(5) = 10$  such runs.
- c. For each of the 5 specific catalysts, there are  $(3)(4) = 12$  pairings of temperature and pressure. Imagine we separate the 60 possible runs into those 5 sets of 12. The number of ways to select exactly one run from each of these 5 sets of 12 is  $\binom{12}{1}^5 = 12^5$ . Since there are  $\binom{60}{5}$  ways to select the 5 runs overall, the desired probability is  $\binom{12}{1}^5 / \binom{60}{5} = 12^5 / \binom{60}{5} = .0456$ .

38.

- a. A sonnet has 14 lines, each of which may come from any of the 10 pages. Order matters, and we're sampling with replacement, so the number of possibilities is  $10 \times 10 \times \dots \times 10 = 10^{14}$ .
- b. Similarly, the number of sonnets you could create avoiding the first and last pages (so, only using lines from the middle 8 sonnets) is  $8^{14}$ . Thus, the probability that a randomly-created sonnet would not use any lines from the first or last page is  $8^{14}/10^{14} = .8^{14} = .044$ .

39.

In **a-c**, the size of the sample space is  $N = \binom{5+6+4}{3} = \binom{15}{3} = 455$ .

- a. There are four 23W bulbs available and  $5+6 = 11$  non-23W bulbs available. The number of ways to select exactly two of the former (and, thus, exactly one of the latter) is  $\binom{4}{2}\binom{11}{1} = 6(11) = 66$ . Hence, the probability is  $66/455 = .145$ .
- b. The number of ways to select three 13W bulbs is  $\binom{5}{3} = 10$ . Similarly, there are  $\binom{6}{3} = 20$  ways to select three 18W bulbs and  $\binom{4}{3} = 4$  ways to select three 23W bulbs. Put together, there are  $10 + 20 + 4 = 34$  ways to select three bulbs of the same wattage, and so the probability is  $34/455 = .075$ .
- c. The number of ways to obtain one of each type is  $\binom{5}{1}\binom{6}{1}\binom{4}{1} = (5)(6)(4) = 120$ , and so the probability is  $120/455 = .264$ .
- d. Rather than consider many different options (choose 1, choose 2, etc.), re-frame the problem this way: at least 6 draws are required to get a 23W bulb iff a random sample of five bulbs fails to produce a 23W bulb. Since there are 11 non-23W bulbs, the chance of getting no 23W bulbs in a sample of size 5 is  $\binom{11}{5} / \binom{15}{5} = 462/3003 = .154$ .

## Chapter 2: Probability

40.

- a. If the  $A$ 's were distinguishable from one another, and similarly for the  $B$ 's,  $C$ 's and  $D$ 's, then there would be  $12!$  possible chain molecules. Six of these are:

$$A_1A_2A_3B_2C_3C_1D_3C_2D_1D_2B_3B_1 \quad A_1A_3A_2B_2C_3C_1D_3C_2D_1D_2B_3B_1$$

$$A_2A_1A_3B_2C_3C_1D_3C_2D_1D_2B_3B_1 \quad A_2A_3A_1B_2C_3C_1D_3C_2D_1D_2B_3B_1$$

$$A_3A_1A_2B_2C_3C_1D_3C_2D_1D_2B_3B_1 \quad A_3A_2A_1B_2C_3C_1D_3C_2D_1D_2B_3B_1$$

These 6 ( $=3!$ ) differ only with respect to ordering of the 3  $A$ 's. In general, groups of 6 chain molecules can be created such that within each group only the ordering of the  $A$ 's is different. When the  $A$  subscripts are suppressed, each group of 6 "collapses" into a single molecule ( $B$ 's,  $C$ 's and  $D$ 's are still distinguishable).

At this point there are  $(12!/3!)$  different molecules. Now suppressing subscripts on the  $B$ 's,  $C$ 's, and  $D$ 's in turn gives  $\frac{12!}{(3!)^4} = 369,600$  chain molecules.

- b. Think of the group of 3  $A$ 's as a single entity, and similarly for the  $B$ 's,  $C$ 's, and  $D$ 's. Then there are  $4! = 24$  ways to order these triplets, and thus 24 molecules in which the  $A$ 's are contiguous, the  $B$ 's,  $C$ 's, and  $D$ 's also. The desired probability is  $\frac{24}{369,600} = .00006494$ .

41.

- a.  $(10)(10)(10)(10) = 10^4 = 10,000$ . These are the strings 0000 through 9999.
- b. Count the number of prohibited sequences. There are (i) 10 with all digits identical (0000, 1111, ..., 9999); (ii) 14 with sequential digits (0123, 1234, 2345, 3456, 4567, 5678, 6789, and 7890, plus these same seven descending); (iii) 100 beginning with 19 (1900 through 1999). That's a total of  $10 + 14 + 100 = 124$  impermissible sequences, so there are a total of  $10,000 - 124 = 9876$  permissible sequences. The chance of randomly selecting one is just  $\frac{9876}{10,000} = .9876$ .
- c. All PINs of the form 8xx1 are legitimate, so there are  $(10)(10) = 100$  such PINs. With someone randomly selecting 3 such PINs, the chance of guessing the correct sequence is  $3/100 = .03$ .
- d. Of all the PINs of the form 1xx1, eleven is prohibited: 1111, and the ten of the form 19x1. That leaves 89 possibilities, so the chances of correctly guessing the PIN in 3 tries is  $3/89 = .0337$ .

42.

- a. If Player X sits out, the number of possible teams is  $\binom{3}{1}\binom{4}{2}\binom{4}{2} = 108$ . If Player X plays guard, we need one more guard, and the number of possible teams is  $\binom{3}{1}\binom{4}{1}\binom{4}{2} = 72$ . Finally, if Player X plays forward, we need one more forward, and the number of possible teams is  $\binom{3}{1}\binom{4}{2}\binom{4}{1} = 72$ . So, the total possible number of teams from this group of 12 players is  $108 + 72 + 72 = 252$ .
- b. Using the idea in a, consider all possible scenarios. If Players X and Y both sit out, the number of possible teams is  $\binom{3}{1}\binom{5}{2}\binom{5}{2} = 300$ . If Player X plays while Player Y sits out, the number of possible

## Chapter 2: Probability

teams is  $\binom{3}{1}\binom{5}{1}\binom{5}{2} + \binom{3}{1}\binom{5}{2}\binom{5}{1} = 150 + 150 = 300$ . Similarly, there are 300 teams with Player X

benched and Player Y in. Finally, there are three cases when X and Y both play: they're both guards, they're both forwards, or they split duties. The number of ways to select the rest of the team under

these scenarios is  $\binom{3}{1}\binom{5}{0}\binom{5}{2} + \binom{3}{1}\binom{5}{2}\binom{5}{0} + \binom{3}{1}\binom{5}{1}\binom{5}{1} = 30 + 30 + 75 = 135$ .

Since there are  $\binom{15}{5} = 3003$  ways to randomly select 5 players from a 15-person roster, the probability

of randomly selecting a legitimate team is  $\frac{300 + 300 + 135}{3003} = \frac{735}{3003} = .245$ .

**43.** There are  $\binom{52}{5} = 2,598,960$  five-card hands. The number of 10-high straights is  $(4)(4)(4)(4)(4) = 4^5 = 1024$

(any of four 6s, any of four 7s, etc.). So,  $P(10 \text{ high straight}) = \frac{1024}{2,598,960} = .000394$ . Next, there ten "types

of straight: A2345, 23456, ..., 910JQK, 10JQKA. So,  $P(\text{straight}) = 10 \times \frac{1024}{2,598,960} = .00394$ . Finally, there

are only 40 straight flushes: each of the ten sequences above in each of the 4 suits makes  $(10)(4) = 40$ . So,

$P(\text{straight flush}) = \frac{40}{2,598,960} = .00001539$ .

**44.** 
$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n!}{(n-k)!k!} = \binom{n}{n-k}$$

The number of subsets of size  $k$  equals the number of subsets of size  $n - k$ , because to each subset of size  $k$  there corresponds exactly one subset of size  $n - k$ : the  $n - k$  objects not in the subset of size  $k$ . The combinations formula counts the number of ways to split  $n$  objects into two subsets: one of size  $k$ , and one of size  $n - k$ .

Section 2.4

45.

a.  $P(A) = .106 + .141 + .200 = .447$ ,  $P(C) = .215 + .200 + .065 + .020 = .500$ , and  $P(A \cap C) = .200$ .

b.  $P(A/C) = \frac{P(A \cap C)}{P(C)} = \frac{.200}{.500} = .400$ . If we know that the individual came from ethnic group 3, the

probability that he has Type A blood is .40.  $P(C/A) = \frac{P(A \cap C)}{P(A)} = \frac{.200}{.447} = .447$ . If a person has Type A blood, the probability that he is from ethnic group 3 is .447.

c. Define  $D =$  "ethnic group 1 selected." We are asked for  $P(D/B')$ . From the table,  $P(D \cap B') = .082 + .106 + .004 = .192$  and  $P(B') = 1 - P(B) = 1 - [.008 + .018 + .065] = .909$ . So, the desired probability is  $P(D/B') = \frac{P(D \cap B')}{P(B')} = \frac{.192}{.909} = .211$ .

46.

Let  $A$  be that the individual is more than 6 feet tall. Let  $B$  be that the individual is a professional basketball player. Then  $P(A/B)$  = the probability of the individual being more than 6 feet tall, knowing that the individual is a professional basketball player, while  $P(B/A)$  = the probability of the individual being a professional basketball player, knowing that the individual is more than 6 feet tall.  $P(A/B)$  will be larger. Most professional basketball players are tall, so the probability of an individual in that reduced sample space being more than 6 feet tall is very large. On the other hand, the number of individuals that are pro basketball players is small in relation to the number of males more than 6 feet tall.

47.

a. Apply the addition rule for three events:  $P(A \cup B \cup C) = .6 + .4 + .2 - .3 - .15 - .1 + .08 = .73$ .

b.  $P(A \cap B \cap C) = P(A \cap B) - P(A \cap B \cap C) = .3 - .08 = .22$ .

c.  $P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{.3}{.6} = .50$  and  $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{.3}{.4} = .75$ . Half of students with Visa cards also have a MasterCard, while three-quarters of students with a MasterCard also have a Visa card.

d.  $P(A \cap B | C) = \frac{P([A \cap B] \cap C)}{P(C)} = \frac{P(A \cap B \cap C)}{P(C)} = \frac{.08}{.2} = .40$ .

e.  $P(A \cup B | C) = \frac{P([A \cup B] \cap C)}{P(C)} = \frac{P([A \cap C] \cup [B \cap C])}{P(C)}$ . Use a distributive law:  

$$= \frac{P(A \cap C) + P(B \cap C) - P([A \cap C] \cap [B \cap C])}{P(C)} = \frac{P(A \cap C) + P(B \cap C) - P(A \cap B \cap C)}{P(C)} =$$

$$\frac{.15 + .1 - .08}{.2} = .85.$$

## Chapter 2: Probability

48.

- a.  $P(A_2 | A_1) = \frac{P(A_2 \cap A_1)}{P(A_1)} = \frac{.06}{.12} = .50$ . The numerator comes from Exercise 26.
- b.  $P(A_1 \cap A_2 \cap A_3 | A_1) = \frac{P([A_1 \cap A_2 \cap A_3] \cap A_1)}{P(A_1)} = \frac{P(A_1 \cap A_2 \cap A_3)}{P(A_1)} = \frac{.01}{.12} = .0833$ . The numerator simplifies because  $A_1 \cap A_2 \cap A_3$  is a subset of  $A_1$ , so their intersection is just the smaller event.

- c. For this example, you definitely need a Venn diagram. The seven pieces of the partition inside the three circles have probabilities .04, .05, .00, .02, .01, .01, and .01. Those add to .14 (so the chance of no defects is .86).

Let  $E =$  “exactly one defect.” From the Venn diagram,  $P(E) = .04 + .00 + .01 = .05$ . From the addition above,  $P(\text{at least one defect}) = P(A_1 \cup A_2 \cup A_3) = .14$ . Finally, the answer to the question is

$$P(E | A_1 \cup A_2 \cup A_3) = \frac{P(E \cap [A_1 \cup A_2 \cup A_3])}{P(A_1 \cup A_2 \cup A_3)} = \frac{P(E)}{P(A_1 \cup A_2 \cup A_3)} = \frac{.05}{.14} = .3571. \text{ The numerator}$$

simplifies because  $E$  is a subset of  $A_1 \cup A_2 \cup A_3$ .

- d.  $P(A'_3 | A_1 \cap A_2) = \frac{P(A'_3 \cap [A_1 \cap A_2])}{P(A_1 \cap A_2)} = \frac{.05}{.06} = .8333$ . The numerator is Exercise 26(c), while the denominator is Exercise 26(b).

49.

- a.  $P(\text{small cup}) = .14 + .20 = .34$ .  $P(\text{decaf}) = .20 + .10 + .10 = .40$ .
- b.  $P(\text{decaf} | \text{small}) = \frac{P(\text{small} \cap \text{decaf})}{P(\text{small})} = \frac{.20}{.34} = .588$ . 58.8% of all people who purchase a small cup of coffee choose decaf.
- c.  $P(\text{small} | \text{decaf}) = \frac{P(\text{small} \cap \text{decaf})}{P(\text{decaf})} = \frac{.20}{.40} = .50$ . 50% of all people who purchase decaf coffee choose the small size.

50.

- a.  $P(\mathbf{M} \cap \mathbf{LS} \cap \mathbf{PR}) = .05$ , directly from the table of probabilities.
- b.  $P(\mathbf{M} \cap \mathbf{Pr}) = P(\mathbf{M} \cap \mathbf{LS} \cap \mathbf{PR}) + P(\mathbf{M} \cap \mathbf{SS} \cap \mathbf{PR}) = .05 + .07 = .12$ .
- c.  $P(\mathbf{SS}) =$  sum of 9 probabilities in the **SS** table  $= .56$ .  $P(\mathbf{LS}) = 1 - .56 = .44$ .
- d. From the two tables,  $P(\mathbf{M}) = .08 + .07 + .12 + .10 + .05 + .07 = .49$ .  $P(\mathbf{Pr}) = .02 + .07 + .07 + .02 + .05 + .02 = .25$ .
- e.  $P(\mathbf{M} | \mathbf{SS} \cap \mathbf{PI}) = \frac{P(\mathbf{M} \cap \mathbf{SS} \cap \mathbf{PI})}{P(\mathbf{SS} \cap \mathbf{PI})} = \frac{.08}{.04 + .08 + .03} = .533$ .
- f.  $P(\mathbf{SS} | \mathbf{M} \cap \mathbf{PI}) = \frac{P(\mathbf{SS} \cap \mathbf{M} \cap \mathbf{PI})}{P(\mathbf{M} \cap \mathbf{PI})} = \frac{.08}{.08 + .10} = .444$ .  $P(\mathbf{LS} | \mathbf{M} \cap \mathbf{PI}) = 1 - P(\mathbf{SS} | \mathbf{M} \cap \mathbf{PI}) = 1 - .444 = .556$ .

## Chapter 2: Probability

51.

- a. Let  $A$  = child has a food allergy, and  $R$  = child has a history of severe reaction. We are told that  $P(A) = .08$  and  $P(R | A) = .39$ . By the multiplication rule,  $P(A \cap R) = P(A) \times P(R | A) = (.08)(.39) = .0312$ .
- b. Let  $M$  = the child is allergic to multiple foods. We are told that  $P(M | A) = .30$ , and the goal is to find  $P(M)$ . But notice that  $M$  is actually a subset of  $A$ : you can't have multiple food allergies without having at least one such allergy! So, apply the multiplication rule again:  
 $P(M) = P(M \cap A) = P(A) \times P(M | A) = (.08)(.30) = .024$ .

52.

We know that  $P(A_1 \cup A_2) = .07$  and  $P(A_1 \cap A_2) = .01$ , and that  $P(A_1) = P(A_2)$  because the pumps are identical. There are two solution methods. The first doesn't require explicit reference to  $q$  or  $r$ : Let  $A_1$  be the event that #1 fails and  $A_2$  be the event that #2 fails.

Apply the addition rule:  $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) \Rightarrow .07 = 2P(A_1) - .01 \Rightarrow P(A_1) = .04$ .

Otherwise, we assume that  $P(A_1) = P(A_2) = q$  and that  $P(A_1 | A_2) = P(A_2 | A_1) = r$  (the goal is to find  $q$ ).

Proceed as follows:  $.01 = P(A_1 \cap A_2) = P(A_1) P(A_2 | A_1) = qr$  and  $.07 = P(A_1 \cup A_2) =$

$$P(A_1 \cap A_2) + P(A_1' \cap A_2) + P(A_1 \cap A_2') = .01 + q(1 - r) + q(1 - r) \Rightarrow q(1 - r) = .03.$$

These two equations give  $2q - .01 = .07$ , from which  $q = .04$  (and  $r = .25$ ).

53.

$$\begin{aligned} P(B|A) &= \frac{P(A \cap B)}{P(A)} = \frac{P(B)}{P(A)} && \text{(since } B \text{ is contained in } A, A \cap B = B) \\ &= \frac{.05}{.60} = .0833 \end{aligned}$$

54.

- a.  $P(A_2 | A_1) = \frac{P(A_1 \cap A_2)}{P(A_1)} = \frac{.11}{.22} = .50$ . If the firm is awarded project 1, there is a 50% chance they will also be awarded project 2.
- b.  $P(A_2 \cap A_3 | A_1) = \frac{P(A_1 \cap A_2 \cap A_3)}{P(A_1)} = \frac{.01}{.22} = .0455$ . If the firm is awarded project 1, there is a 4.55% chance they will also be awarded projects 2 and 3.
- c.  $P(A_2 \cup A_3 | A_1) = \frac{P[A_1 \cap (A_2 \cup A_3)]}{P(A_1)} = \frac{P[(A_1 \cap A_2) \cup (A_1 \cap A_3)]}{P(A_1)}$   
 $= \frac{P(A_1 \cap A_2) + P(A_1 \cap A_3) - P(A_1 \cap A_2 \cap A_3)}{P(A_1)} = \frac{.15}{.22} = .682$ . If the firm is awarded project 1, there is a 68.2% chance they will also be awarded at least one of the other two projects.
- d.  $P(A_1 \cap A_2 \cap A_3 | A_1 \cup A_2 \cup A_3) = \frac{P(A_1 \cap A_2 \cap A_3)}{P(A_1 \cup A_2 \cup A_3)} = \frac{.01}{.53} = .0189$ . If the firm is awarded at least one of the projects, there is a 1.89% chance they will be awarded all three projects.

## Chapter 2: Probability


55. Let  $A = \{\text{carries Lyme disease}\}$  and  $B = \{\text{carries HGE}\}$ . We are told  $P(A) = .16$ ,  $P(B) = .10$ , and  $P(A \cap B | A \cup B) = .10$ . From this last statement and the fact that  $A \cap B$  is contained in  $A \cup B$ ,
- $$.10 = \frac{P(A \cap B)}{P(A \cup B)} \Rightarrow P(A \cap B) = .10P(A \cup B) = .10[P(A) + P(B) - P(A \cap B)] = .10[.10 + .16 - P(A \cap B)] \Rightarrow$$
- $$1.1P(A \cap B) = .026 \Rightarrow P(A \cap B) = .02364.$$
- Finally, the desired probability is  $P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{.02364}{.10} = .2364$ .

56. 
$$P(A | B) + P(A' | B) = \frac{P(A \cap B)}{P(B)} + \frac{P(A' \cap B)}{P(B)} = \frac{P(A \cap B) + P(A' \cap B)}{P(B)} = \frac{P(B)}{P(B)} = 1$$

57.  $P(B | A) > P(B)$  iff  $P(B | A) + P(B' | A) > P(B) + P(B' | A)$  iff  $1 > P(B) + P(B' | A)$  by Exercise 56 (with the letters switched). This holds iff  $1 - P(B) > P(B' | A)$  iff  $P(B') > P(B' | A)$ , QED.

58. 
$$P(A \cup B | C) = \frac{P[(A \cup B) \cap C]}{P(C)} = \frac{P[(A \cap C) \cup (B \cap C)]}{P(C)} = \frac{P(A \cap C) + P(B \cap C) - P(A \cap B \cap C)}{P(C)} = P(A | C) + P(B | C) - P(A \cap B | C)$$


59. The required probabilities appear in the tree diagram below.



- a.  $P(A_2 \cap B) = .21$ .
- b. By the law of total probability,  $P(B) = P(A_1 \cap B) + P(A_2 \cap B) + P(A_3 \cap B) = .455$ .
- c. Using Bayes' theorem,  $P(A_1 | B) = \frac{P(A_1 \cap B)}{P(B)} = \frac{.12}{.455} = .264$ ;  $P(A_2 | B) = \frac{.21}{.455} = .462$ ;  $P(A_3 | B) = 1 - .264 - .462 = .274$ . Notice the three probabilities sum to 1.

## Chapter 2: Probability

60. The tree diagram below shows the probability for the four disjoint options; e.g.,  $P(\text{the flight is discovered and has a locator}) = P(\text{discovered})P(\text{locator} \mid \text{discovered}) = (.7)(.6) = .42$ .



a. 
$$P(\text{not discovered} \mid \text{has locator}) = \frac{P(\text{not discovered} \cap \text{has locator})}{P(\text{has locator})} = \frac{.03}{.03 + .42} = .067.$$

b. 
$$P(\text{discovered} \mid \text{no locator}) = \frac{P(\text{discovered} \cap \text{no locator})}{P(\text{no locator})} = \frac{.28}{.55} = .509.$$

61. The initial (“prior”) probabilities of 0, 1, 2 defectives in the batch are .5, .3, .2. Now, let’s determine the probabilities of 0, 1, 2 defectives in the sample based on these three cases.

- If there are 0 defectives in the batch, clearly there are 0 defectives in the sample.

$$P(0 \text{ def in sample} \mid 0 \text{ def in batch}) = 1.$$

- If there is 1 defective in the batch, the chance it’s discovered in a sample of 2 equals  $2/10 = .2$ , and the probability it isn’t discovered is  $8/10 = .8$ .

$$P(0 \text{ def in sample} \mid 1 \text{ def in batch}) = .8, P(1 \text{ def in sample} \mid 1 \text{ def in batch}) = .2.$$

- If there are 2 defectives in the batch, the chance both are discovered in a sample of 2 equals


$$\frac{2}{10} \times \frac{1}{9} = .022; \text{ the chance neither is discovered equals } \frac{8}{10} \times \frac{7}{9} = .622; \text{ and the chance exactly 1 is}$$

$$\text{discovered equals } 1 - (.022 + .622) = .356.$$

$$P(0 \text{ def in sample} \mid 2 \text{ def in batch}) = .622, P(1 \text{ def in sample} \mid 2 \text{ def in batch}) = .356,$$

$$P(2 \text{ def in sample} \mid 2 \text{ def in batch}) = .022.$$

These calculations are summarized in the tree diagram below. Probabilities at the endpoints are intersectional probabilities, e.g.  $P(2 \text{ def in batch} \cap 2 \text{ def in sample}) = (.2)(.022) = .0044$ .





## Chapter 2: Probability

- a. Using the tree diagram and Bayes' rule,

$$P(0 \text{ def in batch} \mid 0 \text{ def in sample}) = \frac{.5}{.5 + .24 + .1244} = .578$$

$$P(1 \text{ def in batch} \mid 0 \text{ def in sample}) = \frac{.24}{.5 + .24 + .1244} = .278$$

$$P(2 \text{ def in batch} \mid 0 \text{ def in sample}) = \frac{.1244}{.5 + .24 + .1244} = .144$$

- b.  $P(0 \text{ def in batch} \mid 1 \text{ def in sample}) = 0$

$$P(1 \text{ def in batch} \mid 1 \text{ def in sample}) = \frac{.06}{.06 + .0712} = .457$$

$$P(2 \text{ def in batch} \mid 1 \text{ def in sample}) = \frac{.0712}{.06 + .0712} = .543$$

62. Let  $B$  = blue cab was involved,  $G = B'$  = green cab was involved, and  $W$  = witness claims to have seen a blue cab. Before any witness statements,  $P(B) = .15$  and  $P(G)$ . The witness' reliability can be coded as follows:  $P(W \mid B) = .8$  (correctly identify blue),  $P(W' \mid G) = .8$  (correctly identify green), and by taking complements  $P(W' \mid B) = P(W \mid G) = .2$  (the two ways to mis-identify a color at night).


The goal is to determine  $P(B \mid W)$ , the chance a blue cab was involved given that's what the witness claims to have seen. Apply Bayes' Theorem:

$$P(B \mid W) = \frac{P(B)P(W \mid B)}{P(B)P(W \mid B) + P(B')P(W \mid B')} = \frac{(.15)(.8)}{(.15)(.8) + (.85)(.2)} = .4138.$$

The "posterior" probability that the cab was really blue is actually less than 50%. That's because there are so many more green cabs on the street, that it's more likely the witness mis-identified a green cab ( $.85 \times .2$ ) than that the witness correctly identified a blue cab ( $.15 \times .8$ ).

- 63.

- a.



- b. From the top path of the tree diagram,  $P(A \cap B \cap C) = (.75)(.9)(.8) = .54$ .

- c. Event  $B \cap C$  occurs twice on the diagram:  $P(B \cap C) = P(A \cap B \cap C) + P(A' \cap B \cap C) = .54 + (.25)(.8)(.7) = .68$ .

## Chapter 2: Probability

- d.  $P(C) = P(A \cap B \cap C) + P(A' \cap B \cap C) + P(A \cap B' \cap C) + P(A' \cap B' \cap C) = .54 + .045 + .14 + .015 = .74.$
- e. Rewrite the conditional probability first:  $P(A | B \cap C) = \frac{P(A \cap B \cap C)}{P(B \cap C)} = \frac{.54}{.68} = .7941.$
- 64.** A tree diagram can help. We know that  $P(\text{short}) = .6$ ,  $P(\text{medium}) = .3$ ,  $P(\text{long}) = .1$ ; also,  $P(\text{Word} | \text{short}) = .8$ ,  $P(\text{Word} | \text{medium}) = .5$ ,  $P(\text{Word} | \text{long}) = .3$ .
- a. Use the law of total probability:  $P(\text{Word}) = (.6)(.8) + (.3)(.5) + (.1)(.3) = .66.$
- b.  $P(\text{small} | \text{Word}) = \frac{P(\text{small} \cap \text{Word})}{P(\text{Word})} = \frac{(.6)(.8)}{.66} = .727.$  Similarly,  $P(\text{medium} | \text{Word}) = \frac{(.3)(.5)}{.66} = .227,$   
and  $P(\text{long} | \text{Word}) = .045.$  (These sum to .999 due to rounding error.)
- 65.** A tree diagram can help. We know that  $P(\text{day}) = .2$ ,  $P(\text{1-night}) = .5$ ,  $P(\text{2-night}) = .3$ ; also,  $P(\text{purchase} | \text{day}) = .1$ ,  $P(\text{purchase} | \text{1-night}) = .3$ , and  $P(\text{purchase} | \text{2-night}) = .2.$
- Apply Bayes' rule: e.g.,  $P(\text{day} | \text{purchase}) = \frac{P(\text{day} \cap \text{purchase})}{P(\text{purchase})} = \frac{(.2)(.1)}{(.2)(.1) + (.5)(.3) + (.3)(.2)} = \frac{.02}{.23} = .087.$
- Similarly,  $P(\text{1-night} | \text{purchase}) = \frac{(.5)(.3)}{.23} = .652$  and  $P(\text{2-night} | \text{purchase}) = .261.$
- 66.** Let  $E$ ,  $C$ , and  $L$  be the events associated with e-mail, cell phones, and laptops, respectively. We are told  $P(E) = 40\%$ ,  $P(C) = 30\%$ ,  $P(L) = 25\%$ ,  $P(E \cap C) = 23\%$ ,  $P(E' \cap C' \cap L) = 51\%$ ,  $P(E | L) = 88\%$ , and  $P(L | C) = 70\%$ .
- a.  $P(C | E) = P(E \cap C) / P(E) = .23 / .40 = .575.$
- b. Use Bayes' rule:  $P(C | L) = P(C \cap L) / P(L) = P(C)P(L | C) / P(L) = .30(.70) / .25 = .84.$
- c.  $P(C | E \cap L) = P(C \cap E \cap L) / P(E \cap L).$   
For the denominator,  $P(E \cap L) = P(L)P(E | L) = (.25)(.88) = .22.$   
For the numerator, use  $P(E \cup C \cup L) = 1 - P(E' \cap C' \cap L) = 1 - .51 = .49$  and write  
 $P(E \cup C \cup L) = P(C) + P(E) + P(L) - P(E \cap C) - P(C \cap L) - P(E \cap L) + P(C \cap E \cap L)$   
 $\Rightarrow .49 = .30 + .40 + .25 - .23 - .30(.70) - .22 + P(C \cap E \cap L) \Rightarrow P(C \cap E \cap L) = .20.$   
So, finally,  $P(C | E \cap L) = .20 / .22 = .9091.$
- 67.** Let  $T$  denote the event that a randomly selected person is, in fact, a terrorist. Apply Bayes' theorem, using  $P(T) = 1,000 / 300,000,000 = .0000033:$
- $$P(T | +) = \frac{P(T)P(+|T)}{P(T)P(+|T) + P(T')P(+|T')} = \frac{(.0000033)(.99)}{(.0000033)(.99) + (1 - .0000033)(1 - .999)} = .003289.$$
- That is to say, roughly 0.3% of all people "flagged" as terrorists would be actual terrorists in this scenario.

## Chapter 2: Probability

- 68.** Let's see how we can implement the hint. If she's flying airline #1, the chance of 2 late flights is  $(30\%)(10\%) = 3\%$ ; the two flights being "unaffected" by each other means we can multiply their probabilities. Similarly, the chance of 0 late flights on airline #1 is  $(70\%)(90\%) = 63\%$ . Since percents add to 100%, the chance of exactly 1 late flight on airline #1 is  $100\% - (3\% + 63\%) = 34\%$ . A similar approach works for the other two airlines: the probability of exactly 1 late flight on airline #2 is 35%, and the chance of exactly 1 late flight on airline #3 is 45%.

The initial ("prior") probabilities for the three airlines are  $P(A_1) = 50\%$ ,  $P(A_2) = 30\%$ , and  $P(A_3) = 20\%$ . Given that she had exactly 1 late flight (call that event  $B$ ), the conditional ("posterior") probabilities of the three airlines can be calculated using Bayes' Rule:

$$P(A_1 | B) = \frac{P(A_1)P(B | A_1)}{P(A_1)P(B | A_1) + P(A_2)P(B | A_2) + P(A_3)P(B | A_3)} = \frac{(.5)(.34)}{(.5)(.34) + (.3)(.35) + (.2)(.45)} = \frac{.170}{.365} =$$


.4657;

$$P(A_2 | B) = \frac{P(A_2)P(B | A_2)}{P(A_1)P(B | A_1) + P(A_2)P(B | A_2) + P(A_3)P(B | A_3)} = \frac{(.3)(.35)}{.365} = .2877; \text{ and}$$

$$P(A_3 | B) = \frac{P(A_3)P(B | A_3)}{P(A_1)P(B | A_1) + P(A_2)P(B | A_2) + P(A_3)P(B | A_3)} = \frac{(.2)(.45)}{.365} = .2466.$$


Notice that, except for rounding error, these three posterior probabilities add to 1.

The tree diagram below shows these probabilities.



## Chapter 2: Probability

69. The tree diagram below summarizes the information in the exercise (plus the previous information in Exercise 59). Probabilities for the branches corresponding to paying with credit are indicated at the far right. (“extra” = “plus”)



- a.  $P(\text{plus} \cap \text{fill} \cap \text{credit}) = (.35)(.6)(.6) = .1260$ .
- b.  $P(\text{premium} \cap \text{no fill} \cap \text{credit}) = (.25)(.5)(.4) = .05$ .
- c. From the tree diagram,  $P(\text{premium} \cap \text{credit}) = .0625 + .0500 = .1125$ .
- d. From the tree diagram,  $P(\text{fill} \cap \text{credit}) = .0840 + .1260 + .0625 = .2725$ .
- e.  $P(\text{credit}) = .0840 + .1400 + .1260 + .0700 + .0625 + .0500 = .5325$ .
- f.  $P(\text{premium} | \text{credit}) = \frac{P(\text{premium} \cap \text{credit})}{P(\text{credit})} = \frac{.1125}{.5325} = .2113$ .

## Chapter 2: Probability

### Section 2.5

- 70.** Using the definition, two events  $A$  and  $B$  are independent if  $P(A | B) = P(A)$ ;  
 $P(A | B) = .6125$ ;  $P(A) = .50$ ;  $.6125 \neq .50$ , so  $A$  and  $B$  are not independent.  
 Using the multiplication rule, the events are independent if  $P(A \cap B) = P(A)P(B)$ ;  
 $P(A \cap B) = .25$ ;  $P(A)P(B) = (.5)(.4) = .2$ .  $.25 \neq .2$ , so  $A$  and  $B$  are not independent.
- 71.**
- a.** Since the events are independent, then  $A'$  and  $B'$  are independent, too. (See the paragraph below Equation 2.7.) Thus,  $P(B'|A') = P(B') = 1 - .7 = .3$ .
- b.** Using the addition rule,  $P(A \cup B) = P(A) + P(B) - P(A \cap B) = .4 + .7 - (.4)(.7) = .82$ . Since  $A$  and  $B$  are independent, we are permitted to write  $P(A \cap B) = P(A)P(B) = (.4)(.7)$ .
- c.** 
$$P(AB' | A \cup B) = \frac{P(AB' \cap (A \cup B))}{P(A \cup B)} = \frac{P(AB')}{P(A \cup B)} = \frac{P(A)P(B')}{P(A \cup B)} = \frac{(.4)(1-.7)}{.82} = \frac{.12}{.82} = .146.$$
- 72.**  $P(A_1 \cap A_2) = .11$  while  $P(A_1)P(A_2) = .055$ , so  $A_1$  and  $A_2$  are not independent.  
 $P(A_1 \cap A_3) = .05$  while  $P(A_1)P(A_3) = .0616$ , so  $A_1$  and  $A_3$  are not independent.  
 $P(A_2 \cap A_3) = .07$  and  $P(A_2)P(A_3) = .07$ , so  $A_2$  and  $A_3$  are independent.
- 73.** From a Venn diagram,  $P(B) = P(A' \cap B) + P(A \cap B) = P(B) \Rightarrow P(A' \cap B) = P(B) - P(A \cap B)$ . If  $A$  and  $B$  are independent, then  $P(A' \cap B) = P(B) - P(A)P(B) = [1 - P(A)]P(B) = P(A')P(B)$ . Thus,  $A'$  and  $B$  are independent.  
 Alternatively, 
$$P(A' | B) = \frac{P(A' \cap B)}{P(B)} = \frac{P(B) - P(A \cap B)}{P(B)} = \frac{P(B) - P(A)P(B)}{P(B)} = 1 - P(A) = P(A').$$
- 74.** Using subscripts to differentiate between the selected individuals,  
 $P(O_1 \cap O_2) = P(O_1)P(O_2) = (.45)(.45) = .2025$ .  
 $P(\text{two individuals match}) = P(A_1 \cap A_2) + P(B_1 \cap B_2) + P(AB_1 \cap AB_2) + P(O_1 \cap O_2) =$   
 $.40^2 + .11^2 + .04^2 + .45^2 = .3762$ .
- 75.** Let event  $E$  be the event that an error was signaled incorrectly.  
 We want  $P(\text{at least one signaled incorrectly}) = P(E_1 \cup \dots \cup E_{10})$ . To use independence, we need intersections, so apply deMorgan's law:  $= P(E_1 \cup \dots \cup E_{10}) = 1 - P(E'_1 \cap \dots \cap E'_{10})$ .  $P(E') = 1 - .05 = .95$ , so for 10 independent points,  $P(E'_1 \cap \dots \cap E'_{10}) = (.95) \dots (.95) = (.95)^{10}$ . Finally,  $P(E_1 \cup E_2 \cup \dots \cup E_{10}) = 1 - (.95)^{10} = .401$ . Similarly, for 25 points, the desired probability is  $1 - (P(E'))^{25} = 1 - (.95)^{25} = .723$ .

## Chapter 2: Probability

- 76.** Follow the same logic as in Exercise 75: If the probability of an event is  $p$ , and there are  $n$  independent “trials,” the chance this event never occurs is  $(1 - p)^n$ , while the chance of at least one occurrence is  $1 - (1 - p)^n$ . With  $p = 1/9,000,000,000$  and  $n = 1,000,000,000$ , this calculates to  $1 - .9048 = .0952$ .

Note: For extremely small values of  $p$ ,  $(1 - p)^n \approx 1 - np$ . So, the probability of at least one occurrence under these assumptions is roughly  $1 - (1 - np) = np$ . Here, that would equal  $1/9$ .

- 77.** Let  $p$  denote the probability that a rivet is defective.

**a.**  $.15 = P(\text{seam needs reworking}) = 1 - P(\text{seam doesn't need reworking}) =$   
 $1 - P(\text{no rivets are defective}) = 1 - P(1^{\text{st}} \text{ isn't def} \cap \dots \cap 25^{\text{th}} \text{ isn't def}) =$   
 $1 - (1 - p) \dots (1 - p) = 1 - (1 - p)^{25}.$


Solve for  $p$ :  $(1 - p)^{25} = .85 \Rightarrow 1 - p = (.85)^{1/25} \Rightarrow p = 1 - .99352 = .00648.$

**b.** The desired condition is  $.10 = 1 - (1 - p)^{25}$ . Again, solve for  $p$ :  $(1 - p)^{25} = .90 \Rightarrow$   
 $p = 1 - (.90)^{1/25} = 1 - .99579 = .00421.$

- 78.**  $P(\text{at least one opens}) = 1 - P(\text{none open}) = 1 - (.04)^5 = .999999897.$

$P(\text{at least one fails to open}) = 1 - P(\text{all open}) = 1 - (.96)^5 = .1846.$

- 79.** Let  $A_1 =$  older pump fails,  $A_2 =$  newer pump fails, and  $x = P(A_1 \cap A_2)$ . The goal is to find  $x$ . From the Venn diagram below,  $P(A_1) = .10 + x$  and  $P(A_2) = .05 + x$ . Independence implies that  $x = P(A_1 \cap A_2) = P(A_1)P(A_2) = (.10 + x)(.05 + x)$ . The resulting quadratic equation,  $x^2 - .85x + .005 = 0$ , has roots  $x = .0059$  and  $x = .8441$ . The latter is impossible, since the probabilities in the Venn diagram would then exceed 1. Therefore,  $x = .0059$ .



## Chapter 2: Probability


- 80.** Let  $A_i$  denote the event that component # $i$  works ( $i = 1, 2, 3, 4$ ). Based on the design of the system, the event “the system works” is  $(A_1 \cup A_2) \cup (A_3 \cap A_4)$ . We’ll eventually need  $P(A_1 \cup A_2)$ , so work that out first:  $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) = (.9) + (.9) - (.9)(.9) = .99$ . The third term uses independence of events. Also,  $P(A_3 \cap A_4) = (.8)(.8) = .64$ , again using independence.

Now use the addition rule and independence for the system:

$$\begin{aligned} P((A_1 \cup A_2) \cup (A_3 \cap A_4)) &= P(A_1 \cup A_2) + P(A_3 \cap A_4) - P((A_1 \cup A_2) \cap (A_3 \cap A_4)) \\ &= P(A_1 \cup A_2) + P(A_3 \cap A_4) - P(A_1 \cup A_2) \times P(A_3 \cap A_4) \\ &= (.99) + (.64) - (.99)(.64) = .9964 \end{aligned}$$

(You could also use deMorgan’s law in a couple of places.)

- 81.** Using the hints, let  $P(A_i) = p$ , and  $x = p^2$ . Following the solution provided in the example,  $P(\text{system lifetime exceeds } t_0) = p^2 + p^2 - p^4 = 2p^2 - p^4 = 2x - x^2$ . Now, set this equal to .99:  $2x - x^2 = .99 \Rightarrow x^2 - 2x + .99 = 0 \Rightarrow x = 0.9$  or  $1.1 \Rightarrow p = 1.049$  or  $.9487$ . Since the value we want is a probability and cannot exceed 1, the correct answer is  $p = .9487$ .



- 82.**  $A = \{(3,1)(3,2)(3,3)(3,4)(3,5)(3,6)\} \Rightarrow P(A) = \frac{6}{36} = \frac{1}{6}$ ;  $B = \{(1,4)(2,4)(3,4)(4,4)(5,4)(6,4)\} \Rightarrow P(B) = \frac{1}{6}$ ; and  $C = \{(1,6)(2,5)(3,4)(4,3)(5,2)(6,1)\} \Rightarrow P(C) = \frac{1}{6}$ .  
 $A \cap B = \{(3,4)\} \Rightarrow P(A \cap B) = \frac{1}{36} = P(A)P(B)$ ;  $A \cap C = \{(3,4)\} \Rightarrow P(A \cap C) = \frac{1}{36} = P(A)P(C)$ ; and  $B \cap C = \{(3,4)\} \Rightarrow P(B \cap C) = \frac{1}{36} = P(B)P(C)$ . Therefore, these three events are pairwise independent.  
 However,  $A \cap B \cap C = \{(3,4)\} \Rightarrow P(A \cap B \cap C) = \frac{1}{36}$ , while  $P(A)P(B)P(C) = \frac{1}{6} \cdot \frac{1}{6} \cdot \frac{1}{6} = \frac{1}{216}$ , so  $P(A \cap B \cap C) \neq P(A)P(B)P(C)$  and these three events are not mutually independent.

- 83.** We’ll need to know  $P(\text{both detect the defect}) = 1 - P(\text{at least one doesn't}) = 1 - .2 = .8$ .
- a.**  $P(1^{\text{st}} \text{ detects} \cap 2^{\text{nd}} \text{ doesn't}) = P(1^{\text{st}} \text{ detects}) - P(1^{\text{st}} \text{ does} \cap 2^{\text{nd}} \text{ does}) = .9 - .8 = .1$ .  
 Similarly,  $P(1^{\text{st}} \text{ doesn't} \cap 2^{\text{nd}} \text{ does}) = .1$ , so  $P(\text{exactly one does}) = .1 + .1 = .2$ .
- b.**  $P(\text{neither detects a defect}) = 1 - [P(\text{both do}) + P(\text{exactly 1 does})] = 1 - [.8 + .2] = 0$ . That is, under this model there is a 0% probability neither inspector detects a defect. As a result,  $P(\text{all 3 escape}) = (0)(0)(0) = 0$ .

## Chapter 2: Probability

- 84.** We'll make repeated use of the independence of the  $A_i$ s and their complements.
- a.**  $P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2)P(A_3) = (.95)(.98)(.80) = .7448$ .
  - b.** This is the complement of part **a**, so the answer is  $1 - .7448 = .2552$ .
  - c.**  $P(A'_1 \cap A'_2 \cap A'_3) = P(A'_1)P(A'_2)P(A'_3) = (.05)(.02)(.20) = .0002$ .
  - d.**  $P(A'_1 \cap A_2 \cap A_3) = P(A'_1)P(A_2)P(A_3) = (.05)(.98)(.80) = .0392$ .
  - e.**  $P([A'_1 \cap A_2 \cap A_3] \cup [A_1 \cap A'_2 \cap A_3] \cup [A_1 \cap A_2 \cap A'_3]) = (.05)(.98)(.80) + (.95)(.02)(.80) + (.95)(.98)(.20) = .07302$ .
  - f.** This is just a little joke — we've all had the experience of electronics dying right after the warranty expires! ☺
- 85.**
- a.** Let  $D_1 =$  detection on 1<sup>st</sup> fixation,  $D_2 =$  detection on 2<sup>nd</sup> fixation.  
 $P(\text{detection in at most 2 fixations}) = P(D_1) + P(D'_1 \cap D_2)$ ; since the fixations are independent,  
 $P(D_1) + P(D'_1 \cap D_2) = P(D_1) + P(D'_1)P(D_2) = p + (1-p)p = p(2-p)$ .
  - b.** Define  $D_1, D_2, \dots, D_n$  as in **a**. Then  $P(\text{at most } n \text{ fixations}) =$   
 $P(D_1) + P(D'_1 \cap D_2) + P(D'_1 \cap D'_2 \cap D_3) + \dots + P(D'_1 \cap D'_2 \cap \dots \cap D'_{n-1} \cap D_n) =$   
 $p + (1-p)p + (1-p)^2p + \dots + (1-p)^{n-1}p = p[1 + (1-p) + (1-p)^2 + \dots + (1-p)^{n-1}] =$   
 $p \cdot \frac{1-(1-p)^n}{1-(1-p)} = 1 - (1-p)^n$ .  
 Alternatively,  $P(\text{at most } n \text{ fixations}) = 1 - P(\text{at least } n+1 \text{ fixations are required}) =$   
 $1 - P(\text{no detection in 1<sup>st } n \text{ fixations}) = 1 - P(D'_1 \cap D'_2 \cap \dots \cap D'_n) = 1 - (1-p)^n</sup>$ .
  - c.**  $P(\text{no detection in 3 fixations}) = (1-p)^3$ .
  - d.**  $P(\text{passes inspection}) = P(\{\text{not flawed}\} \cup \{\text{flawed and passes}\})$   
 $= P(\text{not flawed}) + P(\text{flawed and passes})$   
 $= .9 + P(\text{flawed})P(\text{passes} \mid \text{flawed}) = .9 + (.1)(1-p)^3$ .
  - e.** Borrowing from **d**,  $P(\text{flawed} \mid \text{passed}) = \frac{P(\text{flawed} \cap \text{passed})}{P(\text{passed})} = \frac{.1(1-p)^3}{.9 + .1(1-p)^3}$ . For  $p = .5$ ,  
 $P(\text{flawed} \mid \text{passed}) = \frac{.1(1-.5)^3}{.9 + .1(1-.5)^3} = .0137$ .



## Chapter 2: Probability

86.

- a.  $P(A) = \frac{2,000}{10,000} = .2$ . Using the law of total probability,  $P(B) = P(A)P(B|A) + P(A')P(B|A') =$   
 $(.2)\frac{1,999}{9,999} + (.8)\frac{2,000}{9,999} = .2$  exactly. That is,  $P(B) = P(A) = .2$ . Finally, use the multiplication rule:  
 $P(A \cap B) = P(A) \times P(B|A) = (.2)\frac{1,999}{9,999} = .039984$ . Events  $A$  and  $B$  are *not* independent, since  $P(B) =$   
 $.2$  while  $P(B|A) = \frac{1,999}{9,999} = .19992$ , and these are not equal.
- b. If  $A$  and  $B$  were independent, we'd have  $P(A \cap B) = P(A) \times P(B) = (.2)(.2) = .04$ . This is very close to the answer .039984 from part a. This suggests that, for most practical purposes, we could treat events  $A$  and  $B$  in this example as if they were independent.
- c. Repeating the steps in part a, you again get  $P(A) = P(B) = .2$ . However, using the multiplication rule,  
 $P(A \cap B) = P(A) \times P(B|A) = \frac{2}{10} \times \frac{1}{9} = .0222$ . This is very different from the value of .04 that we'd get if  $A$  and  $B$  were independent!

The critical difference is that the population size in parts a-b is huge, and so the probability a second board is green *almost* equals .2 (i.e.,  $1,999/9,999 = .19992 \approx .2$ ). But in part c, the conditional probability of a green board shifts a lot:  $2/10 = .2$ , but  $1/9 = .1111$ .

87.

- a. Use the information provided and the addition rule:  
 $P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) \Rightarrow P(A_1 \cap A_2) = P(A_1) + P(A_2) - P(A_1 \cup A_2) = .55 + .65 - .80 = .40$ .
- b. By definition,  $P(A_2 | A_3) = \frac{P(A_2 \cap A_3)}{P(A_3)} = \frac{.40}{.70} = .5714$ . If a person likes vehicle #3, there's a 57.14% chance s/he will also like vehicle #2.
- c. No. From b,  $P(A_2 | A_3) = .5714 \neq P(A_2) = .65$ . Therefore,  $A_2$  and  $A_3$  are not independent. Alternatively,  $P(A_2 \cap A_3) = .40 \neq P(A_2)P(A_3) = (.65)(.70) = .455$ .
- d. The goal is to find  $P(A_2 \cup A_3 | A_1')$ , i.e.  $\frac{P([A_2 \cup A_3] \cap A_1')}{P(A_1')}$ . The denominator is simply  $1 - .55 = .45$ .  
 There are several ways to calculate the numerator; the simplest approach using the information provided is to draw a Venn diagram and observe that  $P([A_2 \cup A_3] \cap A_1') = P(A_1' \cap A_2 \cup A_3) - P(A_1' \cap A_2 \cap A_3) =$   
 $.88 - .55 = .33$ . Hence,  $P(A_2 \cup A_3 | A_1') = \frac{.33}{.45} = .7333$ .

## Chapter 2: Probability

- 88.** Let  $D$  = patient has disease, so  $P(D) = .05$ . Let  $++$  denote the event that the patient gets two independent, positive tests. Given the sensitivity and specificity of the test,  $P(++ | D) = (.98)(.98) = .9604$ , while  $P(++ | D') = (1 - .99)(1 - .99) = .0001$ . (That is, there's a 1-in-10,000 chance of a healthy person being misdiagnosed with the disease twice.) Apply Bayes' Theorem:

$$P(D | ++) = \frac{P(D)P(++ | D)}{P(D)P(++ | D) + P(D')P(++ | D')} = \frac{(.05)(.9604)}{(.05)(.9604) + (.95)(.0001)} = .9980$$

- 89.** The question asks for  $P(\text{exactly one tag lost} | \text{at most one tag lost}) = P((C_1 \cap C_2') \cup (C_1' \cap C_2) | (C_1 \cap C_2)')$ .

Since the first event is contained in (a subset of) the second event, this equals

$$\frac{P((C_1 \cap C_2') \cup (C_1' \cap C_2))}{P((C_1 \cap C_2)')} = \frac{P(C_1 \cap C_2') + P(C_1' \cap C_2)}{1 - P(C_1 \cap C_2)} = \frac{P(C_1)P(C_2') + P(C_1')P(C_2)}{1 - P(C_1)P(C_2)} \text{ by independence} =$$

$$\frac{\pi(1-\pi) + (1-\pi)\pi}{1-\pi^2} = \frac{2\pi(1-\pi)}{1-\pi^2} = \frac{2\pi}{1+\pi}.$$

## Supplementary Exercises

90.

a.  $\binom{10}{3} = 120.$

b. There are 9 other senators from whom to choose the other two subcommittee members, so the answer is  $1 \times \binom{9}{2} = 36.$

c. There are 120 possible subcommittees. Among those, the number which would include none of the 5 most senior senators (i.e., all 3 members are chosen from the 5 most junior senators) is  $\binom{5}{3} = 10.$

Hence, the number of subcommittees with at least one senior senator is  $120 - 10 = 110$ , and the chance of this randomly occurring is  $110/120 = .9167.$

d. The number of subcommittees that can form from the 8 “other” senators is  $\binom{8}{3} = 56$ , so the probability of this event is  $56/120 = .4667.$

91.

a.  $P(\text{line 1}) = \frac{500}{1500} = .333;$

$$P(\text{crack}) = \frac{.50(500) + .44(400) + .40(600)}{1500} = \frac{666}{1500} = .444.$$

b. This is one of the percentages provided:  $P(\text{blemish} | \text{line 1}) = .15.$

c.  $P(\text{surface defect}) = \frac{.10(500) + .08(400) + .15(600)}{1500} = \frac{172}{1500};$

$$P(\text{line 1} \cap \text{surface defect}) = \frac{.10(500)}{1500} = \frac{50}{1500};$$

$$\text{so, } P(\text{line 1} | \text{surface defect}) = \frac{50/1500}{172/1500} = \frac{50}{172} = .291.$$

92.

a. He will have one type of form left if either 4 withdrawals or 4 course substitutions remain. This means the first six were either 2 withdrawals and 4 subs or 6 withdrawals and 0 subs; the desired probability

$$\text{is } \frac{\binom{6}{2}\binom{4}{4} + \binom{6}{6}\binom{4}{0}}{\binom{10}{6}} = \frac{16}{210} = .0762.$$

## Chapter 2: Probability


- b.** He can start with the withdrawal forms or the course substitution forms, allowing two sequences: W-C-W-C or C-W-C-W. The number of ways the first sequence could arise is  $(6)(4)(5)(3) = 360$ , and the number of ways the second sequence could arise is  $(4)(6)(3)(5) = 360$ , for a total of 720 such possibilities. The total number of ways he could select four forms one at a time is  $P_{4,10} = (10)(9)(8)(7) = 5040$ . So, the probability of a perfectly alternating sequence is  $720/5040 = .143$ .

- 93.** Apply the addition rule:  $P(A \cup B) = P(A) + P(B) - P(A \cap B) \Rightarrow .626 = P(A) + P(B) - .144$ . Apply independence:  $P(A \cap B) = P(A)P(B) = .144$ . So,  $P(A) + P(B) = .770$  and  $P(A)P(B) = .144$ . Let  $x = P(A)$  and  $y = P(B)$ . Using the first equation,  $y = .77 - x$ , and substituting this into the second equation yields  $x(.77 - x) = .144$  or  $x^2 - .77x + .144 = 0$ . Use the quadratic formula to solve:  

$$x = \frac{.77 \pm \sqrt{(-.77)^2 - (4)(1)(.144)}}{2(1)} = \frac{.77 \pm .13}{2} = .32 \text{ or } .45$$
. Since  $x = P(A)$  is assumed to be the larger probability,  $x = P(A) = .45$  and  $y = P(B) = .32$ .

- 94.** The probability of a bit reversal is .2, so the probability of maintaining a bit is .8.

- a.** Using independence,  $P(\text{all three relays correctly send } 1) = (.8)(.8)(.8) = .512$ .
- b.** In the accompanying tree diagram, each .2 indicates a bit reversal (and each .8 its opposite). There are several paths that maintain the original bit: no reversals or exactly two reversals (e.g.,  $1 \rightarrow 1 \rightarrow 0 \rightarrow 1$ , which has reversals at relays 2 and 3). The total probability of these options is  $.512 + (.8)(.2)(.2) + (.2)(.8)(.2) + (.2)(.2)(.8) = .512 + 3(.032) = .608$ .



- c.** Using the answer from **b**,  $P(1 \text{ sent} \mid 1 \text{ received}) = \frac{P(1 \text{ sent} \cap 1 \text{ received})}{P(1 \text{ received})} = \frac{P(1 \text{ sent})P(1 \text{ received} \mid 1 \text{ sent})}{P(1 \text{ sent})P(1 \text{ received} \mid 1 \text{ sent}) + P(0 \text{ sent})P(1 \text{ received} \mid 0 \text{ sent})} = \frac{(.7)(.608)}{(.7)(.608) + (.3)(.392)} = \frac{.4256}{.5432} = .7835$ .

In the denominator,  $P(1 \text{ received} \mid 0 \text{ sent}) = 1 - P(0 \text{ received} \mid 0 \text{ sent}) = 1 - .608$ , since the answer from **b** also applies to a 0 being relayed as a 0.

## Chapter 2: Probability

95.

- a. There are  $5! = 120$  possible orderings, so  $P(\text{BCDEF}) = \frac{1}{120} = .0833$ .
- b. The number of orderings in which F is third equals  $4 \times 3 \times 1 \times 2 \times 1 = 24$  (\*because F must be here), so  $P(\text{F is third}) = \frac{24}{120} = .2$ . Or more simply, since the five friends are ordered completely at random, there is a  $\frac{1}{5}$  chance F is specifically in position three.
- c. Similarly,  $P(\text{F last}) = \frac{4 \times 3 \times 2 \times 1 \times 1}{120} = .2$ .
- d.  $P(\text{F hasn't heard after 10 times}) = P(\text{not on \#1} \cap \text{not on \#2} \cap \dots \cap \text{not on \#10}) = \frac{4}{5} \times \dots \times \frac{4}{5} = \left(\frac{4}{5}\right)^{10} = .1074$ .

96.

Palmberg equation:  $P_d(c) = \frac{(c/c^*)^\beta}{1 + (c/c^*)^\beta}$

a.  $P_d(c^*) = \frac{(c^*/c^*)^\beta}{1 + (c^*/c^*)^\beta} = \frac{1^\beta}{1 + 1^\beta} = \frac{1}{1 + 1} = .5$ .

b. The probability of detecting a crack that is twice the size of the “50-50” size  $c^*$  equals

$$P_d(2c^*) = \frac{(2c^*/c^*)^\beta}{1 + (2c^*/c^*)^\beta} = \frac{2^\beta}{1 + 2^\beta}. \text{ When } \beta = 4, P_d(2c^*) = \frac{2^4}{1 + 2^4} = \frac{16}{17} = .9412.$$

c. Using the answers from **a** and **b**,  $P(\text{exactly one of two detected}) = P(\text{first is, second isn't}) + P(\text{first isn't, second is}) = (.5)(1 - .9412) + (1 - .5)(.9412) = .5$ .

d. If  $c = c^*$ , then  $P_d(c) = .5$  irrespective of  $\beta$ . If  $c < c^*$ , then  $c/c^* < 1$  and  $P_d(c) \rightarrow \frac{0}{0+1} = 0$  as  $\beta \rightarrow \infty$ .  
Finally, if  $c > c^*$  then  $c/c^* > 1$  and, from calculus,  $P_d(c) \rightarrow 1$  as  $\beta \rightarrow \infty$ .

97.

When three experiments are performed, there are 3 different ways in which detection can occur on exactly 2 of the experiments: (i) #1 and #2 and not #3; (ii) #1 and not #2 and #3; and (iii) not #1 and #2 and #3. If the impurity is present, the probability of exactly 2 detections in three (independent) experiments is  $(.8)(.8)(.2) + (.8)(.2)(.8) + (.2)(.8)(.8) = .384$ . If the impurity is absent, the analogous probability is  $3(.1)(.1)(.9) = .027$ . Thus, applying Bayes' theorem,  $P(\text{impurity is present} \mid \text{detected in exactly 2 out of 3}) = \frac{P(\text{detected in exactly 2} \cap \text{present})}{P(\text{detected in exactly 2})} = \frac{(.384)(.4)}{(.384)(.4) + (.027)(.6)} = .905$ .

## Chapter 2: Probability

98. Our goal is to find  $P(A \cup B \cup C \cup D \cup E)$ . We'll need all of the following probabilities:

$P(A) = P(\text{Allison gets her calculator back}) = 1/5$ . This is intuitively obvious; you can also see it by writing out the  $5! = 120$  orderings in which the friends could get calculators (ABCDE, ABCED, ..., EDCBA) and observe that 24 of the 120 have A in the first position. So,  $P(A) = 24/120 = 1/5$ . By the same reasoning,  $P(B) = P(C) = P(D) = P(E) = 1/5$ .

$P(A \cap B) = P(\text{Allison and Beth get their calculators back}) = 1/20$ . This can be computed by considering all 120 orderings and noticing that six — those of the form ABxyz — have A and B in the correct positions. Or, you can use the multiplication rule:  $P(A \cap B) = P(A)P(B/A) = (1/5)(1/4) = 1/20$ . All other pairwise intersection probabilities are also  $1/20$ .

$P(A \cap B \cap C) = P(\text{Allison and Beth and Carol get their calculators back}) = 1/60$ , since this can only occur if two ways — ABCDE and ABCED — and  $2/120 = 1/60$ . So, all three-wise intersections have probability  $1/60$ .

$P(A \cap B \cap C \cap D) = 1/120$ , since this can only occur if all 5 girls get their own calculators back. In fact, all four-wise intersections have probability  $1/120$ , as does  $P(A \cap B \cap C \cap D \cap E)$  — they're the same event.

Finally, put all the parts together, using a general inclusion-exclusion rule for unions:

$$\begin{aligned} P(A \cup B \cup C \cup D \cup E) &= P(A) + P(B) + P(C) + P(D) + P(E) \\ &\quad - P(A \cap B) - P(A \cap C) - \dots - P(D \cap E) \\ &\quad + P(A \cap B \cap C) + \dots + P(C \cap D \cap E) \\ &\quad - P(A \cap B \cap C \cap D) - \dots - P(B \cap C \cap D \cap E) \\ &\quad + P(A \cap B \cap C \cap D \cap E) \\ &= 5 \cdot \frac{1}{5} - 10 \cdot \frac{1}{20} + 10 \cdot \frac{1}{60} - 5 \cdot \frac{1}{120} + \frac{1}{120} = 1 - \frac{1}{2} + \frac{1}{6} - \frac{1}{24} + \frac{1}{120} = \frac{76}{120} = .633 \end{aligned}$$

The final answer has the form  $1 - \frac{1}{2} + \frac{1}{6} - \frac{1}{24} = \frac{1}{1!} - \frac{1}{2!} + \frac{1}{3!} - \frac{1}{4!} + \frac{1}{5!}$ . Generalizing to  $n$  friends, the probability at least one will get her own calculator back is  $\frac{1}{1!} - \frac{1}{2!} + \frac{1}{3!} - \frac{1}{4!} + \dots + (-1)^{n-1} \frac{1}{n!}$ .


When  $n$  is large, we can relate this to the power series for  $e^x$  evaluated at  $x = -1$ :

$$\begin{aligned} e^x &= \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \Rightarrow \\ e^{-1} &= 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots = 1 - \left[ \frac{1}{1!} - \frac{1}{2!} + \frac{1}{3!} - \dots \right] \Rightarrow \\ 1 - e^{-1} &= \frac{1}{1!} - \frac{1}{2!} + \frac{1}{3!} - \dots \end{aligned}$$

So, for large  $n$ ,  $P(\text{at least one friend gets her own calculator back}) \approx 1 - e^{-1} = .632$ . Contrary to intuition, the chance of this event does not converge to 1 (because “someone is bound to get hers back”) or to 0 (because “there are just too many possible arrangements”). Rather, in a large group, there's about a 63.2% chance someone will get her own item back (a match), and about a 36.8% chance that nobody will get her own item back (no match).

## Chapter 2: Probability

99. Refer to the tree diagram below.



- a.  $P(\text{pass inspection}) = P(\text{pass initially} \cup \text{passes after recrimping}) = P(\text{pass initially}) + P(\text{fails initially} \cap \text{goes to recrimping} \cap \text{is corrected after recrimping}) = .95 + (.05)(.80)(.60)$  (following path “bad-good-good” on tree diagram)  $= .974$ .
- b.  $P(\text{needed no recrimping} \mid \text{passed inspection}) = \frac{P(\text{passed initially})}{P(\text{passed inspection})} = \frac{.95}{.974} = .9754$ .

100.

- a. First, the probabilities of the  $A_i$  are  $P(A_1) = P(JJ) = (.6)^2 = .36$ ;  $P(A_2) = P(MM) = (.4)^2 = .16$ ; and  $P(A_3) = P(JM \text{ or } MJ) = (.6)(.4) + (.4)(.6) = .48$ .  
Second,  $P(\text{Jay wins} \mid A_1) = 1$ , since Jay is two points ahead and, thus has won;  $P(\text{Jay wins} \mid A_2) = 0$ , since Maurice is two points ahead and, thus, Jay has lost; and  $P(\text{Jay wins} \mid A_3) = p$ , since at that moment the score has returned to deuce and the game has effectively started over. Apply the law of total probability:

$$P(\text{Jay wins}) = P(A_1)P(\text{Jay wins} \mid A_1) + P(A_2)P(\text{Jay wins} \mid A_2) + P(A_3)P(\text{Jay wins} \mid A_3)$$

$$p = (.36)(1) + (.16)(0) + (.48)(p)$$

Therefore,  $p = .36 + .48p$ ; solving for  $p$  gives  $p = \frac{.36}{1 - .48} = .6923$ .

- b. Apply Bayes' rule:  $P(JJ \mid \text{Jay wins}) = \frac{P(JJ)P(\text{Jay wins} \mid JJ)}{P(\text{Jay wins})} = \frac{(.36)(1)}{.6923} = .52$ .

101. Let  $A = 1^{\text{st}}$  functions,  $B = 2^{\text{nd}}$  functions, so  $P(B) = .9$ ,  $P(A \cup B) = .96$ ,  $P(A \cap B) = .75$ . Use the addition rule:  $P(A \cup B) = P(A) + P(B) - P(A \cap B) \Rightarrow .96 = P(A) + .9 - .75 \Rightarrow P(A) = .81$ .

$$\text{Therefore, } P(B \mid A) = \frac{P(B \cap A)}{P(A)} = \frac{.75}{.81} = .926.$$

## Chapter 2: Probability

**102.**

- a.  $P(F) = 919/2026 = .4536$ .  $P(C) = 308/2026 = .1520$ .
- b.  $P(F \cap C) = 110/2026 = .0543$ . Since  $P(F) \times P(C) = .4536 \times .1520 = .0690 \neq .0543$ , we find that events  $F$  and  $C$  are not independent.
- c.  $P(F | C) = P(F \cap C)/P(C) = 110/308 = .3571$ .
- d.  $P(C | F) = P(C \cap F)/P(F) = 110/919 = .1197$ .
- e. Divide each of the two rows, Male and Female, by its row total.

	Blue	Brown	Green	Hazel
<b>Male</b>	.3342	.3180	.1789	.1689
<b>Female</b>	.3906	.3156	.1197	.1741

According to the data, brown and hazel eyes have similar likelihoods for males and females. However, females are much more likely to have blue eyes than males (39% versus 33%) and, conversely, males have a much greater propensity for green eyes than do females (18% versus 12%).

**103.** A tree diagram can help here.

- a.  $P(E_1 \cap L) = P(E_1)P(L | E_1) = (.40)(.02) = .008$ .
- b. The law of total probability gives  $P(L) = \sum P(E_i)P(L | E_i) = (.40)(.02) + (.50)(.01) + (.10)(.05) = .018$ .
- c.  $P(E'_1 | L') = 1 - P(E_1 | L') = 1 - \frac{P(E_1 \cap L')}{P(L')} = 1 - \frac{P(E_1)P(L' | E_1)}{1 - P(L)} = 1 - \frac{(.40)(.98)}{1 - .018} = .601$ .

**104.** Let  $B$  denote the event that a component needs rework. By the law of total probability,  $P(B) = \sum P(A_i)P(B | A_i) = (.50)(.05) + (.30)(.08) + (.20)(.10) = .069$ .

Thus,  $P(A_1 | B) = \frac{(.50)(.05)}{.069} = .362$ ,  $P(A_2 | B) = \frac{(.30)(.08)}{.069} = .348$ , and  $P(A_3 | B) = .290$ .

**105.** This is the famous “Birthday Problem” in probability.

- a. There are  $365^{10}$  possible lists of birthdays, e.g. (Dec 10, Sep 27, Apr 1, ...). Among those, the number with zero matching birthdays is  $P_{10,365}$  (sampling ten birthdays without replacement from 365 days. So,  $P(\text{all different}) = \frac{P_{10,365}}{365^{10}} = \frac{(365)(364) \cdots (356)}{(365)^{10}} = .883$ .  $P(\text{at least two the same}) = 1 - .883 = .117$ .



## Chapter 2: Probability

**b.** The general formula is  $P(\text{at least two the same}) = 1 - \frac{P_{k,365}}{365^k}$ . By trial and error, this probability equals .476 for  $k = 22$  and equals .507 for  $k = 23$ . Therefore, the smallest  $k$  for which  $k$  people have at least a 50-50 chance of a birthday match is 23.


**c.** There are 1000 possible 3-digit sequences to end a SS number (000 through 999). Using the idea from

**a.**  $P(\text{at least two have the same SS ending}) = 1 - \frac{P_{10,1000}}{1000^{10}} = 1 - .956 = .044$ .

Assuming birthdays and SS endings are independent,  $P(\text{at least one "coincidence"}) = P(\text{birthday coincidence} \cup \text{SS coincidence}) = .117 + .044 - (.117)(.044) = .156$ .

## Chapter 2: Probability

106. See the accompanying tree diagram.



- a.  $P(G | R_1 < R_2 < R_3) = \frac{.15}{.15 + .075} = .67$  while  $P(B | R_1 < R_2 < R_3) = .33$ , so classify the specimen as granite. Equivalently,  $P(G | R_1 < R_2 < R_3) = .67 > \frac{1}{2}$  so granite is more likely.
- b.  $P(G | R_1 < R_3 < R_2) = \frac{.0625}{.2125} = .2941 < \frac{1}{2}$ , so classify the specimen as basalt.  
 $P(G | R_3 < R_1 < R_2) = \frac{.0375}{.5625} = .0667 < \frac{1}{2}$ , so classify the specimen as basalt.
- c.  $P(\text{erroneous classification}) = P(B \text{ classified as } G) + P(G \text{ classified as } B) = P(B)P(\text{classified as } G | B) + P(G)P(\text{classified as } B | G) = (.75)P(R_1 < R_2 < R_3 | B) + (.25)P(R_1 < R_3 < R_2 \text{ or } R_3 < R_1 < R_2 | G) = (.75)(.10) + (.25)(.25 + .15) = .175.$
- d. For what values of  $p$  will  $P(G | R_1 < R_2 < R_3)$ ,  $P(G | R_1 < R_3 < R_2)$ , and  $P(G | R_3 < R_1 < R_2)$  all exceed  $\frac{1}{2}$ ? Replacing .25 and .75 with  $p$  and  $1 - p$  in the tree diagram,  

$$P(G | R_1 < R_2 < R_3) = \frac{.6p}{.6p + .1(1-p)} = \frac{.6p}{.1 + .5p} > .5 \text{ iff } p > \frac{1}{7};$$

$$P(G | R_1 < R_3 < R_2) = \frac{.25p}{.25p + .2(1-p)} > .5 \text{ iff } p > \frac{4}{9};$$

$$P(G | R_3 < R_1 < R_2) = \frac{.15p}{.15p + .7(1-p)} > .5 \text{ iff } p > \frac{14}{17} \text{ (most restrictive). Therefore, one would always classify a rock as granite iff } p > \frac{14}{17}.$$

## Chapter 2: Probability

**107.**  $P(\text{detection by the end of the } n\text{th glimpse}) = 1 - P(\text{not detected in first } n \text{ glimpses}) =$   
 $1 - P(G'_1 \cap G'_2 \cap \dots \cap G'_n) = 1 - P(G'_1)P(G'_2) \dots P(G'_n) = 1 - (1 - p_1)(1 - p_2) \dots (1 - p_n) = 1 - \prod_{i=1}^n (1 - p_i).$

**108.**

- a.**  $P(\text{walks on 4}^{\text{th}} \text{ pitch}) = P(\text{first 4 pitches are balls}) = (.5)^4 = .0625.$
- b.**  $P(\text{walks on 6}^{\text{th}} \text{ pitch}) = P(2 \text{ of the first 5 are strikes} \cap \#6 \text{ is a ball}) =$   
 $P(2 \text{ of the first 5 are strikes})P(\#6 \text{ is a ball}) = \binom{5}{2} (.5)^2 (.5)^3 (.5) = .15625.$
- c.** Following the pattern from **b**,  $P(\text{walks on 5}^{\text{th}} \text{ pitch}) = \binom{4}{1} (.5)^1 (.5)^3 (.5) = .125.$  Therefore,  $P(\text{batter walks}) = P(\text{walks on 4}^{\text{th}}) + P(\text{walks on 5}^{\text{th}}) + P(\text{walks on 6}^{\text{th}}) = .0625 + .125 + .15625 = .34375.$
- d.**  $P(\text{first batter scores while no one is out}) = P(\text{first four batters all walk}) = (.34375)^4 = .014.$

**109.**

- a.**  $P(\text{all in correct room}) = \frac{1}{4!} = \frac{1}{24} = .0417.$
- b.** The 9 outcomes which yield completely incorrect assignments are: 2143, 2341, 2413, 3142, 3412, 3421, 4123, 4321, and 4312, so  $P(\text{all incorrect}) = \frac{9}{24} = .375.$

**110.**

- a.**  $P(\text{all full}) = P(A \cap B \cap C) = (.9)(.7)(.8) = .504.$   
 $P(\text{at least one isn't full}) = 1 - P(\text{all full}) = 1 - .504 = .496.$
- b.**  $P(\text{only NY is full}) = P(A \cap B' \cap C') = P(A)P(B')P(C') = (.9)(1-.7)(1-.8) = .054.$   
 Similarly,  $P(\text{only Atlanta is full}) = .014$  and  $P(\text{only LA is full}) = .024.$   
 So,  $P(\text{exactly one full}) = .054 + .014 + .024 = .092.$

## Chapter 2: Probability

- 111.** Note:  $s = 0$  means that the very first candidate interviewed is hired. Each entry below is the candidate hired for the given policy and outcome.

Outcome	$s = 0$	$s = 1$	$s = 2$	$s = 3$	Outcome	$s = 0$	$s = 1$	$s = 2$	$s = 3$
1234	1	4	4	4	3124	3	1	4	4
1243	1	3	3	3	3142	3	1	4	2
1324	1	4	4	4	3214	3	2	1	4
1342	1	2	2	2	3241	3	2	1	1
1423	1	3	3	3	3412	3	1	1	2
1432	1	2	2	2	3421	3	2	2	1
2134	2	1	4	4	4123	4	1	3	3
2143	2	1	3	3	4132	4	1	2	2
2314	2	1	1	4	4213	4	2	1	3
2341	2	1	1	1	4231	4	2	1	1
2413	2	1	1	3	4312	4	3	1	2
2431	2	1	1	1	4321	4	3	2	1

From the table, we derive the following probability distribution based on  $s$ :

$s$	0	1	2	3
$P(\text{hire \#1})$	$\frac{6}{24}$	$\frac{11}{24}$	$\frac{10}{24}$	$\frac{6}{24}$

Therefore  $s = 1$  is the best policy.

- 112.**  $P(\text{at least one occurs}) = 1 - P(\text{none occur}) = 1 - (1 - p_1)(1 - p_2)(1 - p_3)(1 - p_4)$ .  
 $P(\text{at least two occur}) = 1 - P(\text{none or exactly one occur}) =$   
 $1 - [(1 - p_1)(1 - p_2)(1 - p_3)(1 - p_4) + p_1(1 - p_2)(1 - p_3)(1 - p_4) + (1 - p_1)p_2(1 - p_3)(1 - p_4) +$   
 $(1 - p_1)(1 - p_2)p_3(1 - p_4) + (1 - p_1)(1 - p_2)(1 - p_3)p_4]$ .
- 113.**  $P(A_1) = P(\text{draw slip 1 or 4}) = \frac{1}{2}$ ;  $P(A_2) = P(\text{draw slip 2 or 4}) = \frac{1}{2}$ ;  
 $P(A_3) = P(\text{draw slip 3 or 4}) = \frac{1}{2}$ ;  $P(A_1 \cap A_2) = P(\text{draw slip 4}) = \frac{1}{4}$ ;  
 $P(A_2 \cap A_3) = P(\text{draw slip 4}) = \frac{1}{4}$ ;  $P(A_1 \cap A_3) = P(\text{draw slip 4}) = \frac{1}{4}$ .  
Hence  $P(A_1 \cap A_2) = P(A_1)P(A_2) = \frac{1}{4}$ ;  $P(A_2 \cap A_3) = P(A_2)P(A_3) = \frac{1}{4}$ ; and  
 $P(A_1 \cap A_3) = P(A_1)P(A_3) = \frac{1}{4}$ . Thus, there exists pairwise independence. However,  
 $P(A_1 \cap A_2 \cap A_3) = P(\text{draw slip 4}) = \frac{1}{4} \neq \frac{1}{8} = P(A_1)P(A_2)P(A_3)$ , so the events are not mutually independent.
- 114.**  $P(A_1 | A_2 \cap A_3) = \frac{P(A_1 \cap A_2 \cap A_3)}{P(A_2 \cap A_3)} = \frac{P(A_1)P(A_2)P(A_3)}{P(A_2)P(A_3)} = P(A_1)$ .