

# C H A P T E R   1 1

## Vectors and the Geometry of Space

<b>Section 11.1</b>	Vectors in the Plane.....	<b>1091</b>
<b>Section 11.2</b>	Space Coordinates and Vectors in Space .....	<b>1101</b>
<b>Section 11.3</b>	The Dot Product of Two Vectors.....	<b>1110</b>
<b>Section 11.4</b>	The Cross Product of Two Vectors in Space .....	<b>1118</b>
<b>Section 11.5</b>	Lines and Planes in Space .....	<b>1125</b>
<b>Section 11.6</b>	Surfaces in Space.....	<b>1137</b>
<b>Section 11.7</b>	Cylindrical and Spherical Coordinates .....	<b>1141</b>
<b>Review Exercises</b>	.....	<b>1152</b>
<b>Problem Solving</b>	.....	<b>1159</b>

# C H A P T E R 11


## Vectors and the Geometry of Space

### Section 11.1 Vectors in the Plane


1. Answers will vary. *Sample answer:* A scalar is a real number such as 2. A vector is represented by a directed line segment. A vector has both magnitude and direction.

For example  $\langle \sqrt{3}, 1 \rangle$  has direction  $\frac{\pi}{6}$  and a magnitude of 2.

2. Notice that  $\mathbf{v} = \langle 6, -7 \rangle = \langle 2 - (-4), -1 - 6 \rangle = \overrightarrow{QP}$ . Hence,  $Q$  is the initial point and  $P$  is the terminal point.
3. (a)  $\mathbf{v} = \langle 5 - 1, 4 - 2 \rangle = \langle 4, 2 \rangle$



4. (a)  $\mathbf{v} = \langle -4 - 2, -3 - (-3) \rangle = \langle -6, 0 \rangle$



5.  $\mathbf{u} = \langle 5 - 3, 6 - 2 \rangle = \langle 2, 4 \rangle$

$\mathbf{v} = \langle 3 - 1, 8 - 4 \rangle = \langle 2, 4 \rangle$

$\mathbf{u} = \mathbf{v}$

6.  $\mathbf{u} = \langle 1 - (-4), 8 - 0 \rangle = \langle 5, 8 \rangle$

$\mathbf{v} = \langle 7 - 2, 7 - (-1) \rangle = \langle 5, 8 \rangle$

$\mathbf{u} = \mathbf{v}$

7.  $\mathbf{u} = \langle 6 - 0, -2 - 3 \rangle = \langle 6, -5 \rangle$

$\mathbf{v} = \langle 9 - 3, 5 - 10 \rangle = \langle 6, -5 \rangle$

$\mathbf{u} = \mathbf{v}$

8.  $\mathbf{u} = \langle 11 - (-4), -4 - (-1) \rangle = \langle 15, -3 \rangle$


$\mathbf{v} = \langle 25 - 0, 10 - 13 \rangle = \langle 15, -3 \rangle$

$\mathbf{u} = \mathbf{v}$

9. (b)  $\mathbf{v} = \langle 5 - 2, 5 - 0 \rangle = \langle 3, 5 \rangle$

(c)  $\mathbf{v} = 3\mathbf{i} + 5\mathbf{j}$


(a), (d)



10. (b)  $\mathbf{v} = \langle 3 - 4, 6 - (-6) \rangle = \langle -1, 12 \rangle$

(c)  $\mathbf{v} = -\mathbf{i} + 12\mathbf{j}$


(a), (d)



11. (b)  $\mathbf{v} = \langle 6 - 8, -1 - 3 \rangle = \langle -2, -4 \rangle$

(c)  $\mathbf{v} = -2\mathbf{i} - 4\mathbf{j}$


(a), (d)



12. (b)  $\mathbf{v} = \langle -5 - 0, -1 - (-4) \rangle = \langle -5, 3 \rangle$

(c)  $\mathbf{v} = -5\mathbf{i} + 3\mathbf{j}$


(a) and (d).



13. (b)  $\mathbf{v} = \langle 6 - 6, 6 - 2 \rangle = \langle 0, 4 \rangle$

(c)  $\mathbf{v} = 4\mathbf{j}$


(a) and (d).



14. (b)  $\mathbf{v} = \langle -3 - 7, -1 - (-1) \rangle = \langle -10, 0 \rangle$

(c)  $\mathbf{v} = -10\mathbf{i}$


(a) and (d).



15. (b)  $\mathbf{v} = \left\langle \frac{1}{2} - \frac{3}{2}, 3 - \frac{4}{3} \right\rangle = \left\langle -1, \frac{5}{3} \right\rangle$

(c)  $\mathbf{v} = -\mathbf{i} + \frac{5}{3}\mathbf{j}$


(a) and (d).



16. (b)  $\mathbf{v} = \langle 0.84 - 0.12, 1.25 - 0.60 \rangle = \langle 0.72, 0.65 \rangle$

(c)  $\mathbf{v} = 0.72\mathbf{i} + 0.65\mathbf{j}$

(a) and (d).



17.  $u_1 - 4 = -1$

$u_1 = 3$

$u_2 - 2 = 3$

$u_2 = 5$

$Q = (3, 5)$

Terminal point

18.  $u_1 - 5 = 4$

$u_1 = 9$

$u_2 - 3 = -9$

$u_2 = -6$

$Q = (9, -6)$

Terminal point

19.  $\mathbf{v} = 4\mathbf{i}$

$\|\mathbf{v}\| = \sqrt{4^2} = 4$

20.  $\mathbf{v} = -9\mathbf{j}$

$\|\mathbf{v}\| = \sqrt{(-9)^2} = 9$

21.  $\mathbf{v} = \langle 8, 15 \rangle$

$\|\mathbf{v}\| = \sqrt{8^2 + 15^2} = 17$

22.  $\mathbf{v} = \langle -24, 7 \rangle$

$\|\mathbf{v}\| = \sqrt{(-24)^2 + 7^2} = 25$


23.  $\mathbf{v} = -\mathbf{i} - 5\mathbf{j}$

$\|\mathbf{v}\| = \sqrt{(-1)^2 + (-5)^2} = \sqrt{26}$


24.  $\mathbf{v} = 3\mathbf{i} + 3\mathbf{j}$

$\|\mathbf{v}\| = \sqrt{3^2 + 3^2} = \sqrt{18} = 3\sqrt{2}$


25. (a)  $2\mathbf{v} = 2\langle 3, 5 \rangle = \langle 6, 10 \rangle$




(b)  $-3\mathbf{v} = \langle -9, -15 \rangle$




(c)  $\frac{7}{2}\mathbf{v} = \left\langle \frac{21}{2}, \frac{35}{2} \right\rangle$




(d)  $\frac{2}{3}\mathbf{v} = \left\langle 2, \frac{10}{3} \right\rangle$




26. (a)  $4\mathbf{v} = 4\langle -2, 3 \rangle = \langle -8, 12 \rangle$




(b)  $-\frac{1}{2}\mathbf{v} = \left\langle 1, -\frac{3}{2} \right\rangle$



(c)  $0\mathbf{v} = \langle 0, 0 \rangle$



(d)  $-6\mathbf{u} = \langle 12, -18 \rangle$



27.  $\mathbf{u} = \langle 4, 9 \rangle, \mathbf{v} = \langle 2, -5 \rangle$

(a)  $\frac{2}{3}\mathbf{u} = \frac{2}{3}\langle 4, 9 \rangle = \left\langle \frac{8}{3}, 6 \right\rangle$

(b)  $3\mathbf{v} = 3\langle 2, -5 \rangle = \langle 6, -15 \rangle$

(c)  $\mathbf{v} - \mathbf{u} = \langle 2, -5 \rangle - \langle 4, 9 \rangle = \langle -2, -14 \rangle$

$$\begin{aligned} (d) \quad 2\mathbf{u} + 5\mathbf{v} &= 2\langle 4, 9 \rangle + 5\langle 2, -5 \rangle \\ &= \langle 8, 18 \rangle + \langle 10, -25 \rangle \\ &= \langle 18, -7 \rangle \end{aligned}$$

28.  $\mathbf{u} = \langle -3, -8 \rangle, \mathbf{v} = \langle 8, 7 \rangle$


(a)  $\frac{2}{3}\mathbf{u} = \frac{2}{3}\langle -3, -8 \rangle = \left\langle -2, -\frac{16}{3} \right\rangle$

(b)  $3\mathbf{v} = 3\langle 8, 7 \rangle = \langle 24, 21 \rangle$


(c)  $\mathbf{v} - \mathbf{u} = \langle 8, 7 \rangle - \langle -3, -8 \rangle = \langle 11, 15 \rangle$

$$\begin{aligned} (d) \quad 2\mathbf{u} + 5\mathbf{v} &= 2\langle -3, -8 \rangle + 5\langle 8, 7 \rangle \\ &= \langle -6, -16 \rangle + \langle 40, 35 \rangle \\ &= \langle 34, 19 \rangle \end{aligned}$$


29.




30. Twice as long as given vector  $\mathbf{u}$ .




31.




32.



33.



34.



35.  $\mathbf{v} = \langle 3, 12 \rangle$

$$\|\mathbf{v}\| = \sqrt{3^2 + 12^2} = \sqrt{153}$$

$$\begin{aligned}\mathbf{u} &= \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{\langle 3, 12 \rangle}{\sqrt{153}} = \left\langle \frac{3}{\sqrt{153}}, \frac{12}{\sqrt{153}} \right\rangle \\ &= \left\langle \frac{\sqrt{17}}{17}, \frac{4\sqrt{17}}{17} \right\rangle \text{ unit vector}\end{aligned}$$

36.  $\mathbf{v} = \langle -5, 15 \rangle$

$$\|\mathbf{v}\| = \sqrt{25 + 225} = \sqrt{250} = 5\sqrt{10}$$

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{\langle -5, 15 \rangle}{5\sqrt{10}} = \left\langle -\frac{\sqrt{10}}{10}, \frac{3\sqrt{10}}{10} \right\rangle \text{ unit vector}$$

37.  $\mathbf{v} = \left\langle \frac{3}{2}, \frac{5}{2} \right\rangle$

$$\|\mathbf{v}\| = \sqrt{\left(\frac{3}{2}\right)^2 + \left(\frac{5}{2}\right)^2} = \frac{\sqrt{34}}{2}$$

$$\begin{aligned}\mathbf{u} &= \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{\left\langle \frac{3}{2}, \frac{5}{2} \right\rangle}{\frac{\sqrt{34}}{2}} = \left\langle \frac{3}{\sqrt{34}}, \frac{5}{\sqrt{34}} \right\rangle \\ &= \left\langle \frac{3\sqrt{34}}{34}, \frac{5\sqrt{34}}{34} \right\rangle \text{ unit vector}\end{aligned}$$

38.  $\mathbf{v} = \langle -6.2, 3.4 \rangle$

$$\|\mathbf{v}\| = \sqrt{(-6.2)^2 + (3.4)^2} = \sqrt{50} = 5\sqrt{2}$$

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{\langle -6.2, 3.4 \rangle}{5\sqrt{2}} = \left\langle -\frac{31\sqrt{2}}{50}, \frac{17\sqrt{2}}{50} \right\rangle \text{ unit vector}$$

39.  $\mathbf{u} = \langle 1, -1 \rangle, \mathbf{v} = \langle -1, 2 \rangle$

(a)  $\|\mathbf{u}\| = \sqrt{1+1} = \sqrt{2}$

(b)  $\|\mathbf{v}\| = \sqrt{1+4} = \sqrt{5}$

(c)  $\mathbf{u} + \mathbf{v} = \langle 0, 1 \rangle$

$$\|\mathbf{u} + \mathbf{v}\| = \sqrt{0+1} = 1$$

(d)  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{1}{\sqrt{2}} \langle 1, -1 \rangle$

$$\left\| \frac{\mathbf{u}}{\|\mathbf{u}\|} \right\| = 1$$

(e)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{\sqrt{5}} \langle -1, 2 \rangle$

$$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = 1$$

(f)  $\frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} = \langle 0, 1 \rangle$

$$\left\| \frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} \right\| = 1$$

40.  $\mathbf{u} = \langle 0, 1 \rangle, \mathbf{v} = \langle 3, -3 \rangle$

(a)  $\|\mathbf{u}\| = \sqrt{0+1} = 1$

(b)  $\|\mathbf{v}\| = \sqrt{9+9} = 3\sqrt{2}$

(c)  $\mathbf{u} + \mathbf{v} = \langle 3, -2 \rangle$

$$\|\mathbf{u} + \mathbf{v}\| = \sqrt{9+4} = \sqrt{13}$$

(d)  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \langle 0, 1 \rangle$

$$\left\| \frac{\mathbf{u}}{\|\mathbf{u}\|} \right\| = 1$$

(e)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{3\sqrt{2}} \langle 3, -3 \rangle$

$$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = 1$$

(f)  $\frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} = \frac{1}{\sqrt{13}} \langle 3, -2 \rangle$

$$\left\| \frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} \right\| = 1$$

41.  $\mathbf{u} = \left\langle 1, \frac{1}{2} \right\rangle, \mathbf{v} = \langle 2, 3 \rangle$

(a)  $\|\mathbf{u}\| = \sqrt{1 + \frac{1}{4}} = \frac{\sqrt{5}}{2}$

(b)  $\|\mathbf{v}\| = \sqrt{4 + 9} = \sqrt{13}$

(c)  $\mathbf{u} + \mathbf{v} = \left\langle 3, \frac{7}{2} \right\rangle$

$\|\mathbf{u} + \mathbf{v}\| = \sqrt{9 + \frac{49}{4}} = \frac{\sqrt{85}}{2}$

(d)  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{2}{\sqrt{5}} \left\langle 1, \frac{1}{2} \right\rangle$

$\left\| \frac{\mathbf{u}}{\|\mathbf{u}\|} \right\| = 1$

(e)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{\sqrt{13}} \langle 2, 3 \rangle$

$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = 1$

(f)  $\frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} = \frac{2}{\sqrt{85}} \left\langle 3, \frac{7}{2} \right\rangle$

$\left\| \frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} \right\| = 1$

42.  $\mathbf{u} = \langle 2, -4 \rangle, \mathbf{v} = \langle 5, 5 \rangle$

(a)  $\|\mathbf{u}\| = \sqrt{4 + 16} = 2\sqrt{5}$

(b)  $\|\mathbf{v}\| = \sqrt{25 + 25} = 5\sqrt{2}$

(c)  $\mathbf{u} + \mathbf{v} = \langle 7, 1 \rangle$

$\|\mathbf{u} + \mathbf{v}\| = \sqrt{49 + 1} = 5\sqrt{2}$

(d)  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{1}{2\sqrt{5}} \langle 2, -4 \rangle$

$\left\| \frac{\mathbf{u}}{\|\mathbf{u}\|} \right\| = 1$

(e)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{5\sqrt{2}} \langle 5, 5 \rangle$

$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = 1$

(f)  $\frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} = \frac{1}{5\sqrt{2}} \langle 7, 1 \rangle$

$\left\| \frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} \right\| = 1$

43.  $\mathbf{u} = \langle 2, 1 \rangle$

$\|\mathbf{u}\| = \sqrt{5} \approx 2.236$

$\mathbf{v} = \langle 5, 4 \rangle$


$\|\mathbf{v}\| = \sqrt{41} \approx 6.403$

$\mathbf{u} + \mathbf{v} = \langle 7, 5 \rangle$

$\|\mathbf{u} + \mathbf{v}\| = \sqrt{74} \approx 8.602$

$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$

$\sqrt{74} \leq \sqrt{5} + \sqrt{41}$



44.  $\mathbf{u} = \langle -3, 2 \rangle$

$\|\mathbf{u}\| = \sqrt{13} \approx 3.606$

$\mathbf{v} = \langle 1, -2 \rangle$


$\|\mathbf{v}\| = \sqrt{5} \approx 2.236$

$\mathbf{u} + \mathbf{v} = \langle -2, 0 \rangle$

$\|\mathbf{u} + \mathbf{v}\| = 2$

$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$

$2 \leq \sqrt{13} + \sqrt{5}$



45.  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{1}{3} \langle 0, 3 \rangle = \langle 0, 1 \rangle$

$6 \left( \frac{\mathbf{u}}{\|\mathbf{u}\|} \right) = 6 \langle 0, 1 \rangle = \langle 0, 6 \rangle$

$\mathbf{v} = \langle 0, 6 \rangle$

46.  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{1}{\sqrt{2}} \langle 1, 1 \rangle$

$4 \left( \frac{\mathbf{u}}{\|\mathbf{u}\|} \right) = 2\sqrt{2} \langle 1, 1 \rangle$

$\mathbf{v} = \langle 2\sqrt{2}, 2\sqrt{2} \rangle$

47.  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{1}{\sqrt{5}} \langle -1, 2 \rangle = \left\langle -\frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}} \right\rangle$

$5 \left( \frac{\mathbf{u}}{\|\mathbf{u}\|} \right) = 5 \left\langle -\frac{1}{\sqrt{5}}, \frac{2}{\sqrt{5}} \right\rangle = \langle -\sqrt{5}, 2\sqrt{5} \rangle$

$\mathbf{v} = \langle -\sqrt{5}, 2\sqrt{5} \rangle$

48.  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{1}{2\sqrt{3}} \langle \sqrt{3}, 3 \rangle$

$2 \left( \frac{\mathbf{u}}{\|\mathbf{u}\|} \right) = \frac{1}{\sqrt{3}} \langle \sqrt{3}, 3 \rangle$

$\mathbf{v} = \langle 1, \sqrt{3} \rangle$

49.  $\mathbf{v} = 3[(\cos 0^\circ)\mathbf{i} + (\sin 0^\circ)\mathbf{j}] = 3\mathbf{i} = \langle 3, 0 \rangle$

50.  $\mathbf{v} = 5[(\cos 120^\circ)\mathbf{i} + (\sin 120^\circ)\mathbf{j}]$   
 $= -\frac{5}{2}\mathbf{i} + \frac{5\sqrt{3}}{2}\mathbf{j} = \left\langle -\frac{5}{2}, \frac{5\sqrt{3}}{2} \right\rangle$

51.  $\mathbf{v} = 2[(\cos 150^\circ)\mathbf{i} + (\sin 150^\circ)\mathbf{j}]$   
 $= -\sqrt{3}\mathbf{i} + \mathbf{j} = \langle -\sqrt{3}, 1 \rangle$

52.  $\mathbf{v} = 4[(\cos 3.5^\circ)\mathbf{i} + (\sin 3.5^\circ)\mathbf{j}]$   
 $\approx 3.9925\mathbf{i} + 0.2442\mathbf{j}$   
 $= \langle 3.9925, 0.2442 \rangle$

53.  $\mathbf{u} = (\cos 0^\circ)\mathbf{i} + (\sin 0^\circ)\mathbf{j} = \mathbf{i}$   
 $\mathbf{v} = 3(\cos 45^\circ)\mathbf{i} + 3(\sin 45^\circ)\mathbf{j} = \frac{3\sqrt{2}}{2}\mathbf{i} + \frac{3\sqrt{2}}{2}\mathbf{j}$   
 $\mathbf{u} + \mathbf{v} = \left\langle \frac{2+3\sqrt{2}}{2}, \frac{3\sqrt{2}}{2} \right\rangle$

54.  $\mathbf{u} = 4(\cos 0^\circ)\mathbf{i} + 4(\sin 0^\circ)\mathbf{j} = 4\mathbf{i}$   
 $\mathbf{v} = 2(\cos 30^\circ)\mathbf{i} + 2(\sin 30^\circ)\mathbf{j} = \mathbf{i} + \sqrt{3}\mathbf{j}$   
 $\mathbf{u} + \mathbf{v} = 5\mathbf{i} + \sqrt{3}\mathbf{j} = \langle 5, \sqrt{3} \rangle$

60. (a) True.  $\mathbf{d}$  has the same magnitude as  $\mathbf{a}$  but is in the opposite direction.  
 (b) True.  $\mathbf{c}$  and  $\mathbf{s}$  have the same length and direction.  
 (c) True.  $\mathbf{a}$  and  $\mathbf{u}$  are the adjacent sides of a parallelogram. So, the resultant vector,  $\mathbf{a} + \mathbf{u}$ , is the diagonal of the parallelogram,  $\mathbf{c}$ .  
 (d) False. The negative of a vector has the opposite direction of the original vector.

- (e) True.  $\mathbf{a} + \mathbf{d} = \mathbf{a} + (-\mathbf{a}) = 0$   
 (f) False.  $\mathbf{u} - \mathbf{v} = \mathbf{u} - (-\mathbf{u}) = 2\mathbf{u}$

$$-2(\mathbf{b} + \mathbf{t}) = -2(\mathbf{b} + \mathbf{b}) = -2(2\mathbf{b}) = -2[2(-\mathbf{u})] = 4\mathbf{u}$$

61.  $\mathbf{v} = \langle 4, 5 \rangle = a\langle 1, 2 \rangle + b\langle 1, -1 \rangle$

$$4 = a + b$$

$$5 = 2a - b$$

Adding the equations,  $9 = 3a \Rightarrow a = 3$ .

Then you have  $b = 4 - a = 4 - 3 = 1$ .

$$a = 3, b = 1$$

62.  $\mathbf{v} = \langle -7, -2 \rangle = a\langle 1, 2 \rangle + b\langle 1, -1 \rangle$

$$-7 = a + b$$

$$-2 = 2a - b$$

Adding the equations,  $-9 = 3a \Rightarrow a = -3$ .

Then you have  $b = -7 - a = -7 - (-3) = -4$ .

$$a = -3, b = -4$$


55.  $\mathbf{u} = 2(\cos 4)\mathbf{i} + 2(\sin 4)\mathbf{j}$   
 $\mathbf{v} = (\cos 2)\mathbf{i} + (\sin 2)\mathbf{j}$   
 $\mathbf{u} + \mathbf{v} = (2 \cos 4 + \cos 2)\mathbf{i} + (2 \sin 4 + \sin 2)\mathbf{j}$   
 $= \langle 2 \cos 4 + \cos 2, 2 \sin 4 + \sin 2 \rangle$

56.  $\mathbf{u} = 5[\cos(-0.5)]\mathbf{i} + 5[\sin(-0.5)]\mathbf{j}$   
 $= 5(\cos 0.5)\mathbf{i} - 5(\sin 0.5)\mathbf{j}$   
 $\mathbf{v} = 5(\cos 0.5)\mathbf{i} + 5(\sin 0.5)\mathbf{j}$   
 $\mathbf{u} + \mathbf{v} = 10(\cos 0.5)\mathbf{i} = \langle 10 \cos 0.5, 0 \rangle$

57. The forces act along the same direction.  $\theta = 0^\circ$ .

58. The forces cancel out each other.  $\theta = 180^\circ$ .

59.



$$\overrightarrow{XY} + \overrightarrow{YZ} + \overrightarrow{ZX} = 0.$$

Vectors that start and end at the same point have a magnitude of 0.

63.  $\mathbf{v} = \langle -6, 0 \rangle = a\langle 1, 2 \rangle + 6\langle 1, -1 \rangle$

$$-6 = a + b$$

$$0 = 2a - b$$

Adding the equations,  $-6 = 3a \Rightarrow a = -2$ .

Then you have  $b = -6 - a = -6 - (-2) = -4$ .

$$a = -2, b = -4$$

64.  $\mathbf{v} = \langle 0, 6 \rangle = a\langle 1, 2 \rangle + b\langle 1, -1 \rangle$

$$0 = a + b$$

$$6 = 2a - b$$

Adding the equations,  $6 = 3a \Rightarrow a = 2$ .

Then you have  $b = -a = -2$ .

$$a = 2, b = -2$$

65.  $\mathbf{v} = \langle 1, -3 \rangle = a\langle 1, 2 \rangle + b\langle 1, -1 \rangle$

$$1 = a + b$$

$$-3 = 2a - b$$

Adding the equations,  $-2 = 3a \Rightarrow a = -\frac{2}{3}$ .

Then you have  $b = 1 - a = 1 - \left(-\frac{2}{3}\right) = \frac{5}{3}$ .

$$a = -\frac{2}{3}, b = \frac{5}{3}$$

67.  $f(x) = x^2, f'(x) = 2x, f'(3) = 6$

(a)  $m = 6$ . Let  $\mathbf{w} = \langle 1, 6 \rangle, \|\mathbf{w}\| = \sqrt{37}$ , then  $\pm \frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{37}}\langle 1, 6 \rangle$ .

(b)  $m = -\frac{1}{6}$ . Let  $\mathbf{w} = \langle -6, 1 \rangle, \|\mathbf{w}\| = \sqrt{37}$ , then  $\pm \frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{37}}\langle -6, 1 \rangle$ .

66.  $\mathbf{v} = \langle -1, 8 \rangle = a\langle 1, 2 \rangle + b\langle 1, -1 \rangle$


$$-1 = a + b$$

$$8 = 2a - b$$

Adding the equations,  $7 = 3a \Rightarrow a = \frac{7}{3}$ .

Then you have  $b = -1 - a = -1 - \frac{7}{3} = -\frac{10}{3}$ .


$$a = \frac{7}{3}, b = -\frac{10}{3}$$



68.  $f(x) = -x^2 + 5, f'(x) = -2x, f'(-1) = -2$

(a)  $m = -2$ . Let  $\mathbf{w} = \langle 1, -2 \rangle, \|\mathbf{w}\| = \sqrt{5}$ , then  $\pm \frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{5}}\langle 1, -2 \rangle$ .


(b)  $m = \frac{1}{2}$ . Let  $\mathbf{w} = \langle 2, 1 \rangle, \|\mathbf{w}\| = \sqrt{5}$ , then  $\pm \frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{5}}\langle 2, 1 \rangle$ .



69.  $f(x) = x^3, f'(x) = 3x^2 = 3$  at  $x = 1$ .

(a)  $m = 3$ . Let  $\mathbf{w} = \langle 1, 3 \rangle, \|\mathbf{w}\| = \sqrt{10}$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{10}}\langle 1, 3 \rangle$ .


(b)  $m = -\frac{1}{3}$ . Let  $\mathbf{w} = \langle 3, -1 \rangle, \|\mathbf{w}\| = \sqrt{10}$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{10}}\langle 3, -1 \rangle$ .



70.  $f(x) = x^3, f'(-2) = 3x^2 = 12$  at  $x = -2$ .

(a)  $m = 12$ . Let  $\mathbf{w} = \langle 1, 12 \rangle, \|\mathbf{w}\| = \sqrt{145}$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{145}}\langle 1, 12 \rangle$ .

(b)  $m = -\frac{1}{12}$ . Let  $\mathbf{w} = \langle 12, -1 \rangle, \|\mathbf{w}\| = \sqrt{145}$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{145}}\langle 12, -1 \rangle$ .




71.  $f(x) = \sqrt{25 - x^2}$

$$f'(x) = \frac{-x}{\sqrt{25 - x^2}} = \frac{-3}{4} \text{ at } x = 3.$$


(a)  $m = -\frac{3}{4}$ . Let  $\mathbf{w} = \langle -4, 3 \rangle, \|\mathbf{w}\| = 5$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{5}\langle -4, 3 \rangle$ .

(b)  $m = \frac{4}{3}$ . Let  $\mathbf{w} = \langle 3, 4 \rangle, \|\mathbf{w}\| = 5$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{5}\langle 3, 4 \rangle$ .



72.  $f(x) = \tan x$

$f'(x) = \sec^2 x = 2$  at  $x = \frac{\pi}{4}$

(a)  $m = 2$ . Let  $\mathbf{w} = \langle 1, 2 \rangle$ ,  $\|\mathbf{w}\| = \sqrt{5}$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{5}} \langle 1, 2 \rangle$ .(b)  $m = -\frac{1}{2}$ . Let  $\mathbf{w} = \langle -2, 1 \rangle$ ,  $\|\mathbf{w}\| = \sqrt{5}$ , then  $\frac{\mathbf{w}}{\|\mathbf{w}\|} = \pm \frac{1}{\sqrt{5}} \langle -2, 1 \rangle$ .

73.  $\mathbf{u} = \frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{j}$

$\mathbf{u} + \mathbf{v} = \sqrt{2} \mathbf{j}$

$\mathbf{v} = (\mathbf{u} + \mathbf{v}) - \mathbf{u} = -\frac{\sqrt{2}}{2} \mathbf{i} + \frac{\sqrt{2}}{2} \mathbf{j} = \left\langle -\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right\rangle$

74.  $\mathbf{u} = 2\sqrt{3} \mathbf{i} + 2\mathbf{j}$

$\mathbf{u} + \mathbf{v} = -3\mathbf{i} + 3\sqrt{3} \mathbf{j}$

$\mathbf{v} = (\mathbf{u} + \mathbf{v}) - \mathbf{u} = (-3 - 2\sqrt{3})\mathbf{i} + (3\sqrt{3} - 2)\mathbf{j}$   
 $= \langle -3 - 2\sqrt{3}, 3\sqrt{3} - 2 \rangle$

75.  $\mathbf{F}_1 + \mathbf{F}_2 = (500 \cos 30^\circ \mathbf{i} + 500 \sin 30^\circ \mathbf{j}) + (200 \cos(-45^\circ) \mathbf{i} + 200 \sin(-45^\circ) \mathbf{j}) = (250\sqrt{3} + 100\sqrt{2})\mathbf{i} + (250 - 100\sqrt{2})\mathbf{j}$

$\|\mathbf{F}_1 + \mathbf{F}_2\| = \sqrt{(250\sqrt{3} + 100\sqrt{2})^2 + (250 - 100\sqrt{2})^2} \approx 584.6 \text{ N}$

$\tan \theta = \frac{250 - 100\sqrt{2}}{250\sqrt{3} + 100\sqrt{2}} \Rightarrow \theta \approx 10.7^\circ$

76. (a)  $180(\cos 30^\circ \mathbf{i} + \sin 30^\circ \mathbf{j}) + 275\mathbf{i} \approx 430.88\mathbf{i} + 90\mathbf{j}$



Direction:  $\alpha \approx \arctan\left(\frac{90}{430.88}\right) \approx 0.206 \text{ } (\approx 11.8^\circ)$

Magnitude:  $\sqrt{430.88^2 + 90^2} \approx 440.18 \text{ newtons}$

(b)  $M = \sqrt{(275 + 180 \cos \theta)^2 + (180 \sin \theta)^2}$

$\alpha = \arctan\left[\frac{180 \sin \theta}{275 + 180 \cos \theta}\right]$

$\theta$	$0^\circ$	$30^\circ$	$60^\circ$	$90^\circ$	$120^\circ$	$150^\circ$	$180^\circ$
$M$	455	440.2	396.9	328.7	241.9	149.3	95
$\alpha$	$0^\circ$	$11.8^\circ$	$23.1^\circ$	$33.2^\circ$	$40.1^\circ$	$37.1^\circ$	$0^\circ$

(e)  $M$  decreases because the forces change from acting in the same direction to acting in the opposite direction as  $\theta$  increases from  $0^\circ$  to  $180^\circ$ .

77.  $\mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3 = (75 \cos 30^\circ \mathbf{i} + 75 \sin 30^\circ \mathbf{j}) + (100 \cos 45^\circ \mathbf{i} + 100 \sin 45^\circ \mathbf{j}) + (125 \cos 120^\circ \mathbf{i} + 125 \sin 120^\circ \mathbf{j})$

$= \left(\frac{75}{2}\sqrt{3} + 50\sqrt{2} - \frac{125}{2}\right)\mathbf{i} + \left(\frac{75}{2} + 50\sqrt{2} + \frac{125}{2}\sqrt{3}\right)\mathbf{j}$

$\|\mathbf{R}\| = \|\mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3\| \approx 228.5 \text{ N}$

$\theta_{\mathbf{R}} = \theta_{\mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3} \approx 71.3^\circ$


78.  $\mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3 = [400(\cos(-30^\circ) \mathbf{i} + \sin(-30^\circ) \mathbf{j})] + [280(\cos(45^\circ) \mathbf{i} + \sin(45^\circ) \mathbf{j})] + [350(\cos(135^\circ) \mathbf{i} + \sin(135^\circ) \mathbf{j})]$

$= [200\sqrt{3} + 140\sqrt{2} - 175\sqrt{2}]\mathbf{i} + [-200 + 140\sqrt{2} + 175\sqrt{2}]\mathbf{j}$

$\|\mathbf{R}\| = \sqrt{(200\sqrt{3} - 35\sqrt{2})^2 + (-200 + 315\sqrt{2})^2} \approx 385.2483 \text{ newtons}$

$\theta_{\mathbf{R}} = \arctan\left(\frac{-200 + 315\sqrt{2}}{200\sqrt{3} - 35\sqrt{2}}\right) \approx 0.6908 \approx 39.6^\circ$

79.  $\mathbf{u} = \overrightarrow{CB} = \|\mathbf{u}\|(\cos 30^\circ \mathbf{i} + \sin 30^\circ \mathbf{j})$   
 $\mathbf{v} = \overrightarrow{CA} = \|\mathbf{v}\|(\cos 130^\circ \mathbf{i} + \sin 130^\circ \mathbf{j})$



Vertical components:  $\|\mathbf{u}\| \sin 30^\circ + \|\mathbf{v}\| \sin 130^\circ = 3000$

Horizontal components:  $\|\mathbf{u}\| \cos 30^\circ + \|\mathbf{v}\| \cos 130^\circ = 0$

Solving this system, you obtain

$\|\mathbf{u}\| \approx 1958.1$  newtons,

$\|\mathbf{v}\| \approx 2638.2$  newtons.

80.  $\theta_1 = \arctan\left(\frac{24}{20}\right) \approx 0.8761$  or  $50.2^\circ$

$\theta_2 = \arctan\left(\frac{24}{-10}\right) + \pi \approx 1.9656$  or  $112.6^\circ$

$\mathbf{u} = \|\mathbf{u}\|(\cos \theta_1 \mathbf{i} + \sin \theta_1 \mathbf{j})$


$\mathbf{v} = \|\mathbf{v}\|(\cos \theta_2 \mathbf{i} + \sin \theta_2 \mathbf{j})$

Vertical components:  $\|\mathbf{u}\| \sin \theta_1 + \|\mathbf{v}\| \sin \theta_2 = 5000$

Horizontal components:  $\|\mathbf{u}\| \cos \theta_1 + \|\mathbf{v}\| \cos \theta_2 = 0$

Solving this system, you obtain

$\|\mathbf{u}\| \approx 2169.4$  and  $\|\mathbf{v}\| \approx 3611.2$ .



81. Horizontal component =  $\|\mathbf{v}\| \cos \theta$

=  $120 \cos 6^\circ \approx 119.343$  m/sec

Vertical component =  $\|\mathbf{v}\| \sin \theta$

=  $120 \sin 6^\circ \approx 12.543$  m/sec

82. To lift the weight vertically, the sum of the vertical components of  $\mathbf{u}$  and  $\mathbf{v}$  must be 100 and the sum of the horizontal components must be 0.

$\mathbf{u} = \|\mathbf{u}\|(\cos 60^\circ \mathbf{i} + \sin 60^\circ \mathbf{j})$

$\mathbf{v} = \|\mathbf{v}\|(\cos 110^\circ \mathbf{i} + \sin 110^\circ \mathbf{j})$

So,  $\|\mathbf{u}\| \sin 60^\circ + \|\mathbf{v}\| \sin 110^\circ = 100$ , or  $\|\mathbf{u}\| \left(\frac{\sqrt{3}}{2}\right) + \|\mathbf{v}\| \sin 110^\circ = 100$ .

And  $\|\mathbf{u}\| \cos 60^\circ + \|\mathbf{v}\| \cos 110^\circ = 0$  or  $\|\mathbf{u}\| \left(\frac{1}{2}\right) + \|\mathbf{v}\| \cos 110^\circ = 0$ .

Multiplying the last equation by  $(\sqrt{3})$  and adding to the first equation gives

$\|\mathbf{u}\| (\sin 110^\circ - \sqrt{3} \cos 110^\circ) = 100 \Rightarrow \|\mathbf{v}\| \approx 65.27$  newtons.


Then,  $\|\mathbf{u}\| \left(\frac{1}{2}\right) + 65.27 \cos 110^\circ = 0$  gives  $\|\mathbf{u}\| \approx 44.65$  newtons.

(a) The tension in each rope:  $\|\mathbf{u}\| = 44.65$  N,

$\|\mathbf{v}\| = 65.27$  N

(b) Vertical components:  $\|\mathbf{u}\| \sin 60^\circ \approx 38.67$  N,

$\|\mathbf{v}\| \sin 110^\circ \approx 61.33$  N



83.  $\mathbf{u} = 900(\cos 148^\circ \mathbf{i} + \sin 148^\circ \mathbf{j})$

$\mathbf{v} = 100(\cos 45^\circ \mathbf{i} + \sin 45^\circ \mathbf{j})$

$$\mathbf{u} + \mathbf{v} = (900 \cos 148^\circ + 100 \cos 45^\circ) \mathbf{i} + (900 \sin 148^\circ + 100 \sin 45^\circ) \mathbf{j}$$

$$\approx -692.53 \mathbf{i} + 547.64 \mathbf{j}$$

$$\theta \approx \arctan\left(\frac{547.64}{-692.53}\right) \approx -38.34^\circ; 38.34^\circ \text{ North of West}$$

$$\|\mathbf{u} + \mathbf{v}\| \approx \sqrt{(-692.53)^2 + (547.64)^2} \approx 882.9 \text{ km/h}$$

84.  $\mathbf{u} = 400\mathbf{i}$  (resultant)

$$\mathbf{v} = 50(\cos(-45^\circ)\mathbf{i} + \sin(-45^\circ)\mathbf{j}) = 25\sqrt{2}\mathbf{i} - 25\sqrt{2}\mathbf{j}$$
 (wind)

$$\mathbf{u} - \mathbf{v} = (400 - 25\sqrt{2})\mathbf{i} + 25\sqrt{2}\mathbf{j} \approx 364.64\mathbf{i} + 35.36\mathbf{j}$$
 (plane)

$$\tan \theta = \frac{35.36}{364.64} \Rightarrow \theta \approx 5.54^\circ$$

Compass direction:  $\approx$  N  $84.46^\circ$ E

Speed:  $\approx 366.35$  km/h

85. False. Weight has direction.

86. True

87. True

88. True

89. True

90. True

91. True

92. False

$$a = b = 0$$

93. False

$$\|a\mathbf{i} + b\mathbf{j}\| = \sqrt{2}|a|$$

94. True


95.  $\|\mathbf{u}\| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1$ ,

$$\|\mathbf{v}\| = \sqrt{\sin^2 \theta + \cos^2 \theta} = 1$$

96. Let the triangle have vertices at  $(0, 0)$ ,  $(a, 0)$ , and  $(b, c)$ .

Let  $\mathbf{u}$  be the vector joining  $(0, 0)$  and  $(b, c)$ , as indicated in the figure. Then  $\mathbf{v}$ , the vector joining the midpoints, is

$$\begin{aligned} \mathbf{v} &= \left(\frac{a+b}{2} - \frac{a}{2}\right)\mathbf{i} + \frac{c}{2}\mathbf{j} \\ &= \frac{b}{2}\mathbf{i} + \frac{c}{2}\mathbf{j} \\ &= \frac{1}{2}(b\mathbf{i} + c\mathbf{j}) = \frac{1}{2}\mathbf{u}. \end{aligned}$$



97. Let  $\mathbf{u}$  and  $\mathbf{v}$  be the vectors that determine the parallelogram, as indicated in the figure. The two diagonals are  $\mathbf{u} + \mathbf{v}$  and  $\mathbf{v} - \mathbf{u}$ . So,


$$\mathbf{r} = x(\mathbf{u} + \mathbf{v}), \mathbf{s} = 4(\mathbf{v} - \mathbf{u}).$$
 But,

$$\mathbf{u} = \mathbf{r} - \mathbf{s}$$

$$= x(\mathbf{u} + \mathbf{v}) - y(\mathbf{v} - \mathbf{u}) = (x + y)\mathbf{u} + (x - y)\mathbf{v}.$$

So,  $x + y = 1$  and  $x - y = 0$ . Solving you have

$$x = y = \frac{1}{2}.$$



$$\begin{aligned}
98. \quad & \mathbf{w} = \|\mathbf{u}\|\mathbf{v} + \|\mathbf{v}\|\mathbf{u} \\
& = \|\mathbf{u}\|[\|\mathbf{v}\|\cos\theta_v\mathbf{i} + \|\mathbf{v}\|\sin\theta_v\mathbf{j}] + \|\mathbf{v}\|[\|\mathbf{u}\|\cos\theta_u\mathbf{i} + \|\mathbf{u}\|\sin\theta_u\mathbf{j}] \\
& = \|\mathbf{u}\|\|\mathbf{v}\|[(\cos\theta_u + \cos\theta_v)\mathbf{i} + (\sin\theta_u + \sin\theta_v)\mathbf{j}] \\
& = 2\|\mathbf{u}\|\|\mathbf{v}\|\left[\cos\left(\frac{\theta_u + \theta_v}{2}\right)\cos\left(\frac{\theta_u - \theta_v}{2}\right)\mathbf{i} + \sin\left(\frac{\theta_u + \theta_v}{2}\right)\cos\left(\frac{\theta_u - \theta_v}{2}\right)\mathbf{j}\right] \\
& \tan\theta_w = \frac{\sin\left(\frac{\theta_u + \theta_v}{2}\right)\cos\left(\frac{\theta_u - \theta_v}{2}\right)}{\cos\left(\frac{\theta_u + \theta_v}{2}\right)\cos\left(\frac{\theta_u - \theta_v}{2}\right)} = \tan\left(\frac{\theta_u + \theta_v}{2}\right)
\end{aligned}$$

So,  $\theta_w = (\theta_u + \theta_v)/2$  and  $\mathbf{w}$  bisects the angle between  $\mathbf{u}$  and  $\mathbf{v}$ .

99. The set is a circle of radius 5, centered at the origin.

$$\|\mathbf{u}\| = \|\langle x, y \rangle\| = \sqrt{x^2 + y^2} = 5 \Rightarrow x^2 + y^2 = 25$$

100. Let  $x = v_0 t \cos \alpha$  and  $y = v_0 t \sin \alpha - \frac{1}{2}gt^2$ .


$$\begin{aligned}
t &= \frac{x}{v_0 \cos \alpha} \Rightarrow y = v_0 \sin \alpha \left( \frac{x}{v_0 \cos \alpha} \right) - \frac{1}{2}g \left( \frac{x}{v_0 \cos \alpha} \right)^2 \\
&= x \tan \alpha - \frac{g}{2v_0^2} x^2 \sec^2 \alpha \\
&= x \tan \alpha - \frac{gx^2}{2v_0^2} (1 + \tan^2 \alpha) \\
&= \frac{v_0^2}{2g} - \frac{gx^2}{2v_0^2} - \frac{gx^2}{2v_0^2} \tan^2 \alpha + x \tan \alpha - \frac{v_0^2}{2g} \\
&= \frac{v_0^2}{2g} - \frac{gx^2}{2v_0^2} - \frac{gx^2}{2v_0^2} \left[ \tan^2 \alpha - 2 \tan \alpha \left( \frac{v_0^2}{gx} \right) + \frac{v_0^4}{g^2 x^2} \right] \\
&= \frac{v_0^2}{2g} - \frac{gx^2}{2v_0^2} - \frac{gx^2}{2v_0^2} \left( \tan \alpha - \frac{v_0^2}{gx} \right)^2
\end{aligned}$$

If  $y \leq \frac{v_0^2}{2g} - \frac{gx^2}{2v_0^2}$ , then  $\alpha$  can be chosen to hit the point  $(x, y)$ . To hit  $(0, y)$ : Let  $\alpha = 90^\circ$ . Then

$$y = v_0 t - \frac{1}{2}gt^2 = \frac{v_0^2}{2g} - \frac{v_0^2}{2g} \left( \frac{g}{v_0} t - 1 \right)^2, \text{ and you need } y \leq \frac{v_0^2}{2g}$$

The set H is given by  $0 \leq x$ ,  $0 < y$  and  $y \leq \frac{v_0^2}{2g} - \frac{gx^2}{2v_0^2}$


**Note:** The parabola  $y = \frac{v_0^2}{2g} - \frac{gx^2}{2v_0^2}$  is called the “parabola of safety.”




## Section 11.2 Space Coordinates and Vectors in Space

1.  $x_0$  is directed distance to  $yz$ -plane.
2. The  $y$ -coordinate of any point in the  $xz$ -plane is 0.
3. (a)  $x = 4$  is a point on the number line.  
 (b)  $x = 4$  is a vertical line in the plane.  
 (c)  $x = 4$  is a plane in space.
4. The nonzero vectors  $\mathbf{u}$  and  $\mathbf{v}$  are parallel if there exists a scalar such that  $\mathbf{u} = c\mathbf{v}$ .


5.




6.



7.



8.



9.  $x = -3, y = 4, z = 5$ :  $(-3, 4, 5)$

10.  $x = 7, y = -2, z = -1$ :  $(7, -2, -1)$

11.  $y = z = 0, x = 12$ :  $(12, 0, 0)$

12.  $x = 0, y = 3, z = 2$ :  $(0, 3, 2)$

13. The point is 1 unit above the  $xy$ -plane.

14. The point is 6 units in front of the  $xz$ -plane.

15. The point is on the plane parallel to the  $yz$ -plane that passes through  $x = -3$ .

16. The point is 5 units below the  $xy$ -plane.

17. The point is to the left of the  $xz$ -plane.

18. The point more than 4 units away from the  $yz$ -plane.

19. The point is on or between the planes  $y = 3$  and  $y = -3$ .

20. The point is in front of the plane  $x = 4$ .

21. The point  $(x, y, z)$  is 3 units below the  $xy$ -plane, and below either quadrant I or III.

22. The point  $(x, y, z)$  is 4 units above the  $xy$ -plane, and above either quadrant II or IV.

23. The point could be above the  $xy$ -plane and so above quadrants II or IV, or below the  $xy$ -plane, and so below quadrants I or III.

24. The point could be above the  $xy$ -plane, and so above quadrants I and III, or below the  $xy$ -plane, and so below quadrants II or IV.

$$\begin{aligned} 25. d &= \sqrt{(8-4)^2 + (2-1)^2 + (6-5)^2} \\ &= \sqrt{16+1+1} \\ &= \sqrt{18} = 3\sqrt{2} \end{aligned}$$

$$\begin{aligned} 26. d &= \sqrt{(-3-(-1))^2 + (5-1)^2 + (-3-1)^2} \\ &= \sqrt{4+16+16} \\ &= \sqrt{36} = 6 \end{aligned}$$

$$\begin{aligned} 27. d &= \sqrt{(3-0)^2 + (2-2)^2 + (8-4)^2} \\ &= \sqrt{9+0+16} \\ &= \sqrt{25} = 5 \end{aligned}$$

$$\begin{aligned} 28. d &= \sqrt{(-5-(-3))^2 + (8-7)^2 + (-4-1)^2} \\ &= \sqrt{4+1+25} \\ &= \sqrt{30} \end{aligned}$$

29.  $A(0, 0, 4), B(2, 6, 7), C(6, 4, -8)$

$$|AB| = \sqrt{2^2 + 6^2 + 3^2} = \sqrt{49} = 7$$

$$|AC| = \sqrt{6^2 + 4^2 + (-12)^2} = \sqrt{196} = 14$$

$$|BC| = \sqrt{4^2 + (-2)^2 + (-15)^2} = \sqrt{245} = 7\sqrt{5}$$

$$|BC|^2 = 245 = 49 + 196 = |AB|^2 + |AC|^2$$

Right triangle

30.  $A(3, 4, 1), B(0, 6, 2), C(3, 5, 6)$

$$|AB| = \sqrt{9+4+1} = \sqrt{14}$$

$$|AC| = \sqrt{0+1+25} = \sqrt{26}$$

$$|BC| = \sqrt{9+1+16} = \sqrt{26}$$

Because  $|AC| = |BC|$ , the triangle is isosceles.

31.  $A(-1, 0, -2), B(-1, 5, 2), C(-3, -1, 1)$

$$|AB| = \sqrt{0 + 25 + 16} = \sqrt{41}$$

$$|AC| = \sqrt{4 + 1 + 9} = \sqrt{14}$$

$$|BC| = \sqrt{4 + 36 + 1} = \sqrt{41}$$

Because  $|AB| = |BC|$ , the triangle is isosceles.

32.  $A(4, -1, -1), B(2, 0, -4), C(3, 5, -1)$

$$|AB| = \sqrt{4 + 1 + 9} = \sqrt{14}$$

$$|AC| = \sqrt{1 + 36 + 0} = \sqrt{37}$$

$$|BC| = \sqrt{1 + 25 + 9} = \sqrt{35}$$

Neither

33.  $\left(\frac{4+8}{2}, \frac{0+8}{2}, \frac{-6+20}{2}\right) = (6, 4, 7)$

39. Center is midpoint of diameter:

$$\left(\frac{2+1}{2}, \frac{1+3}{2}, \frac{3-1}{2}\right) = \left(\frac{3}{2}, 2, 1\right)$$

Radius is distance from center to endpoint:

$$d = \sqrt{\left(\frac{3}{2} - 1\right)^2 + (2 - 3)^2 + (1 + 1)^2} = \sqrt{\frac{1}{4} + 1 + 4} = \frac{\sqrt{21}}{2}$$

$$\left(x - \frac{3}{2}\right)^2 + (y - 2)^2 + (z - 1)^2 = \frac{21}{4}$$

40. Center is midpoint of diameter:

$$\left(\frac{-2-4}{2}, \frac{4+0}{2}, \frac{-5+3}{2}\right) = (-3, 2, -1)$$

Radius is distance from center to endpoint:

$$d = \sqrt{(-4 - (-3))^2 + (0 - 2)^2 + (3 - (-1))^2} = \sqrt{1 + 4 + 16} = \sqrt{21}$$

$$(x + 3)^2 + (y - 2)^2 + (z + 1)^2 = 21$$

41. Center:  $(-7, 7, 6)$

Tangent to  $xy$ -plane

Radius is  $z$ -coordinate, 6.

$$(x + 7)^2 + (y - 7)^2 + (z - 6)^2 = 36$$

42. Center:  $(-4, 0, 0)$

Tangent to  $yz$ -plane

Radius is distance to  $yz$ -plane, 4.

$$(x + 4)^2 + y^2 + z^2 = 16$$

43.  $x^2 + y^2 + z^2 - 2x + 6y + 8z + 1 = 0$

$$(x^2 - 2x + 1) + (y^2 + 6y + 9) + (z^2 + 8z + 16) = -1 + 1 + 9 + 16$$

$$(x - 1)^2 + (y + 3)^2 + (z + 4)^2 = 25$$

Center:  $(1, -3, -4)$

Radius: 5

34.  $\left(\frac{7-5}{2}, \frac{2-2}{2}, \frac{2-3}{2}\right) = \left(1, 0, -\frac{1}{2}\right)$

35.  $\left(\frac{3+1}{2}, \frac{4+8}{2}, \frac{6+0}{2}\right) = (2, 6, 3)$

36.  $\left(\frac{5+(-2)}{2}, \frac{-9+3}{2}, \frac{7+3}{2}\right) = \left(\frac{3}{2}, -3, 5\right)$

37. Center:  $(7, 1, -2)$

Radius: 1

$$(x - 7)^2 + (y - 1)^2 + (z + 2)^2 = 1$$

38. Center:  $(-1, -5, 8)$

Radius: 5

$$(x + 1)^2 + (y + 5)^2 + (z - 8)^2 = 25$$

44.  $x^2 + y^2 + z^2 + 9x - 2y + 10z + 19 = 0$

$$\left(x^2 + 9x + \frac{81}{4}\right) + (y^2 - 2y + 1) + (z^2 + 10z + 25) = -19 + \frac{81}{4} + 1 + 25$$

$$\left(x + \frac{9}{2}\right)^2 + (y - 1)^2 + (z + 5)^2 = \frac{109}{4}$$

Center:  $\left(-\frac{9}{2}, 1, -5\right)$

Radius:  $\frac{\sqrt{109}}{2}$

45.  $9x^2 + 9y^2 + 9z^2 - 6x + 18y + 1 = 0$

$$x^2 + y^2 + z^2 - \frac{2}{3}x + 2y + \frac{1}{9} = 0$$

$$(x^2 - \frac{2}{3}x + \frac{1}{9}) + (y^2 + 2y + 1) + z^2 = -\frac{1}{9} + \frac{1}{9} + 1$$

$$\left(x - \frac{1}{3}\right)^2 + (y + 1)^2 + (z - 0)^2 = 1$$

Center:  $\left(\frac{1}{3}, -1, 0\right)$

Radius: 1

46.  $4x^2 + 4y^2 + 4z^2 - 24x - 4y + 8z - 23 = 0$

$$(x^2 - 6x + 9) + (y^2 - y + \frac{1}{4}) + (z^2 + 2z + 1) = \frac{23}{4} + 9 + \frac{1}{4} + 1$$

$$(x - 3)^2 + \left(y - \frac{1}{2}\right)^2 + (z + 1)^2 = 16$$


Center:  $\left(3, \frac{1}{2}, -1\right)$

Radius: 4

47. (a)  $\mathbf{v} = \langle 2 - 4, 4 - 2, 3 - 1 \rangle = \langle -2, 2, 2 \rangle$

(b)  $\mathbf{v} = -2\mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$


(c)



49. (b)  $\mathbf{v} = \langle 3 - (-1), 3 - 2, 4 - 3 \rangle = \langle 4, 1, 1 \rangle$

(c)  $\mathbf{v} = 4\mathbf{i} + \mathbf{j} + \mathbf{k}$


(a), (d)



48. (a)  $\mathbf{v} = \langle 4 - 0, 0 - 5, 3 - 1 \rangle = \langle 4, -5, 2 \rangle$

(b)  $\mathbf{v} = 4\mathbf{i} - 5\mathbf{j} + 2\mathbf{k}$


(c)



50. (b)  $\mathbf{v} = \langle -4 - 2, 3 - (-1), 7 - (-2) \rangle = \langle -6, 4, 9 \rangle$

(c)  $\mathbf{v} = 6\mathbf{i} + 4\mathbf{j} + 9\mathbf{k}$

(a), (d)



51.  $\mathbf{v} = \langle 4 - 3, 1 - 2, 6 - 0 \rangle = \langle 1, -1, 6 \rangle$

$$\|\mathbf{v}\| = \sqrt{1 + 1 + 36} = \sqrt{38}$$

Unit vector:  $\frac{\langle 1, -1, 6 \rangle}{\sqrt{38}} = \left\langle \frac{1}{\sqrt{38}}, \frac{-1}{\sqrt{38}}, \frac{6}{\sqrt{38}} \right\rangle$

52.  $\mathbf{v} = \langle 2 - 1, 4 - (-2), -2 - 4 \rangle = \langle 1, 6, -6 \rangle$

$$\|\mathbf{v}\| = \sqrt{1 + 36 + 36} = \sqrt{73}$$

Unit vector:  $\frac{\langle 1, 6, -6 \rangle}{\sqrt{73}} = \left\langle \frac{1}{\sqrt{73}}, \frac{6}{\sqrt{73}}, \frac{-6}{\sqrt{73}} \right\rangle$

53.  $\mathbf{v} = \langle 0 - 4, 5 - 2, 2 - 0 \rangle = \langle -4, 3, 2 \rangle$

$$\|\mathbf{v}\| = \sqrt{(-4)^2 + 3^2 + 2^2} = \sqrt{16 + 9 + 4} = \sqrt{29}$$

Unit vector:

$$\frac{1}{\sqrt{29}} \langle -4, 3, 2 \rangle = \left\langle -\frac{4}{\sqrt{29}}, \frac{3}{\sqrt{29}}, \frac{2}{\sqrt{29}} \right\rangle$$

54.  $\mathbf{v} = \langle 1 - 1, -2 - (-2), -3 - 0 \rangle = \langle 0, 0, -3 \rangle$

$$\|\mathbf{v}\| = \sqrt{0^2 + 0^2 + (-3)^2} = 3$$

Unit vector:

$$\frac{1}{3} \langle 0, 0, -3 \rangle = \langle 0, 0, -1 \rangle$$


55.  $(q_1, q_2, q_3) - (0, 6, 2) = (3, -5, 6)$

$$Q = (3, 1, 8)$$


56.  $(q_1, q_2, q_3) - (0, 2, \frac{5}{2}) = (1, -\frac{2}{3}, \frac{1}{2})$

$$Q = \left(1, -\frac{4}{3}, 3\right)$$


57. (a)  $2\mathbf{v} = \langle 2, 4, 4 \rangle$




(b)  $-\mathbf{v} = \langle -1, -2, -2 \rangle$




(c)  $\frac{3}{2}\mathbf{v} = \left\langle \frac{3}{2}, 3, 3 \right\rangle$




(d)  $0\mathbf{v} = \langle 0, 0, 0 \rangle$




58. (a)  $-\mathbf{v} = \langle -2, 2, -1 \rangle$




(b)  $2\mathbf{v} = \langle 4, -4, 2 \rangle$



(c)  $\frac{1}{2}\mathbf{v} = \langle 1, -1, \frac{1}{2} \rangle$



(d)  $\frac{5}{2}\mathbf{v} = \langle 5, -5, \frac{5}{2} \rangle$



59.  $\mathbf{z} = \mathbf{u} - \mathbf{v} + \mathbf{w}$

$$\begin{aligned} &= \langle 1, 2, 3 \rangle - \langle 2, 2, -1 \rangle + \langle 4, 0, -4 \rangle \\ &= \langle 3, 0, 0 \rangle \end{aligned}$$

60.  $\mathbf{z} = 5\mathbf{u} - 3\mathbf{v} - \frac{1}{2}\mathbf{w}$

$$\begin{aligned} &= \langle 5, 10, 15 \rangle - \langle 6, 6, -3 \rangle - \langle 2, 0, -2 \rangle \\ &= \langle -3, 4, 20 \rangle \end{aligned}$$

61.  $\frac{1}{3}\mathbf{z} - 3\mathbf{u} = \mathbf{w}$

$$\frac{1}{3}\mathbf{z} = 3\mathbf{u} + \mathbf{w}$$

$$\mathbf{z} = 9\mathbf{u} + 3\mathbf{w}$$

$$= 9\langle 1, 2, 3 \rangle + 3\langle 4, 0, -4 \rangle$$

$$= \langle 9, 18, 27 \rangle + \langle 12, 0, -12 \rangle$$

$$= \langle 21, 18, 15 \rangle$$

62.  $2\mathbf{u} + \mathbf{v} - \mathbf{w} + 3\mathbf{z} = 2\langle 1, 2, 3 \rangle + \langle 2, 2, -1 \rangle - \langle 4, 0, -4 \rangle + 3\langle z_1, z_2, z_3 \rangle = \langle 0, 0, 0 \rangle$

$$\langle 0, 6, 9 \rangle + \langle 3z_1, 3z_2, 3z_3 \rangle = \langle 0, 0, 0 \rangle$$

$$0 + 3z_1 = 0 \Rightarrow z_1 = 0$$

$$6 + 3z_2 = 0 \Rightarrow z_2 = -2$$

$$9 + 3z_3 = 0 \Rightarrow z_3 = -3$$

$$\mathbf{z} = \langle 0, -2, -3 \rangle$$

63. (a) and (b) are parallel because

$$\langle -6, -4, 10 \rangle = -2\langle 3, 2, -5 \rangle \text{ and}$$

$$\langle 2, \frac{4}{3}, -\frac{10}{3} \rangle = \frac{2}{3}\langle 3, 2, -5 \rangle.$$

64. (b) and (d) are parallel because

$$-\mathbf{i} + \frac{4}{3}\mathbf{j} - \frac{3}{2}\mathbf{k} = -2\left(\frac{1}{2}\mathbf{i} - \frac{2}{3}\mathbf{j} + \frac{3}{4}\mathbf{k}\right) \text{ and}$$

$$\frac{3}{4}\mathbf{i} - \mathbf{j} + \frac{9}{8}\mathbf{k} = \frac{3}{2}\left(\frac{1}{2}\mathbf{i} - \frac{2}{3}\mathbf{j} + \frac{3}{4}\mathbf{k}\right).$$

65.  $\mathbf{z} = -3\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$

(a) is parallel because  $-6\mathbf{i} + 8\mathbf{j} + 4\mathbf{k} = 2\mathbf{z}$ .

66.  $\mathbf{z} = \langle -7, -8, 3 \rangle$

(b) is parallel because  $(-z)\mathbf{z} = \langle 14, 16, -6 \rangle$ .

67.  $P(0, -2, -5), Q(3, 4, 4), R(2, 2, 1)$

$$\overline{PQ} = \langle 3, 6, 9 \rangle$$

$$\overline{PR} = \langle 2, 4, 6 \rangle$$

$$\langle 3, 6, 9 \rangle = \frac{3}{2}\langle 2, 4, 6 \rangle$$

So,  $\overline{PQ}$  and  $\overline{PR}$  are parallel, the points are collinear.

68.  $P(4, -2, 7), Q(-2, 0, 3), R(7, -3, 9)$

$$\overline{PQ} = \langle -6, 2, -4 \rangle$$

$$\overline{PR} = \langle 3, -1, 2 \rangle$$

$$\langle 3, -1, 2 \rangle = -\frac{1}{2}\langle -6, 2, -4 \rangle$$

So,  $\overline{PQ}$  and  $\overline{PR}$  are parallel. The points are collinear.

69.  $P(1, 2, 4), Q(2, 5, 0), R(0, 1, 5)$

$$\overline{PQ} = \langle 1, 3, -4 \rangle$$

$$\overline{PR} = \langle -1, -1, 1 \rangle$$

Because  $\overline{PQ}$  and  $\overline{PR}$  are not parallel, the points are not collinear.

70.  $P(0, 0, 0), Q(1, 3, -2), R(2, -6, 4)$

$$\overline{PQ} = \langle 1, 3, -2 \rangle$$

$$\overline{PR} = \langle 2, -6, 4 \rangle$$

Because  $\overline{PQ}$  and  $\overline{PR}$  are not parallel, the points are not collinear.

71.  $A(2, 9, 1), B(3, 11, 4), C(0, 10, 2), D(1, 12, 5)$

$$\overline{AB} = \langle 1, 2, 3 \rangle$$

$$\overline{CD} = \langle 1, 2, 3 \rangle$$

$$\overline{AC} = \langle -2, 1, 1 \rangle$$

$$\overline{BD} = \langle -2, 1, 1 \rangle$$

Because  $\overline{AB} = \overline{CD}$  and  $\overline{AC} = \overline{BD}$ , the given points form the vertices of a parallelogram.

72.  $A(1, 1, 3), B(9, -1, -2), C(11, 2, -9), D(3, 4, -4)$

$$\overline{AB} = \langle 8, -2, -5 \rangle$$

$$\overline{DC} = \langle 8, -2, -5 \rangle$$

$$\overline{AD} = \langle 2, 3, -7 \rangle$$

$$\overline{BC} = \langle 2, 3, -7 \rangle$$

Because  $\overline{AB} = \overline{DC}$  and  $\overline{AD} = \overline{BC}$ , the given points form the vertices of a parallelogram.

73.  $\|\mathbf{v}\| = \|\langle -1, 0, 1 \rangle\|$   
 $= \sqrt{(-1)^2 + 0^2 + 1^2}$   
 $= \sqrt{1+1} = \sqrt{2}$

74.  $\|\mathbf{v}\| = \|\langle -5, -3, -4 \rangle\|$   
 $= \sqrt{(-5)^2 + (-3)^2 + (-4)^2}$   
 $= \sqrt{25 + 9 + 16}$   
 $= \sqrt{50}$   
 $= 5\sqrt{2}$

75.  $\mathbf{v} = 3\mathbf{j} - 5\mathbf{k} = \langle 0, 3, -5 \rangle$   
 $\|\mathbf{v}\| = \sqrt{0 + 9 + 25} = \sqrt{34}$

76.  $\mathbf{v} = 2\mathbf{i} + 5\mathbf{j} - \mathbf{k} = \langle 2, 5, -1 \rangle$   
 $\|\mathbf{v}\| = \sqrt{4 + 25 + 1} = \sqrt{30}$

81.  $\mathbf{v} = 4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}$   
 $\|\mathbf{v}\| = \sqrt{16 + 25 + 9} = \sqrt{50} = 5\sqrt{2}$

(a)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{5\sqrt{2}}(4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}) = \frac{2\sqrt{2}}{5}\mathbf{i} - \frac{\sqrt{2}}{2}\mathbf{j} + \frac{3\sqrt{2}}{10}\mathbf{k}$   
(b)  $-\frac{\mathbf{v}}{\|\mathbf{v}\|} = -\frac{1}{5\sqrt{2}}(4\mathbf{i} - 5\mathbf{j} + 3\mathbf{k}) = -\frac{2\sqrt{2}}{5}\mathbf{i} + \frac{\sqrt{2}}{2}\mathbf{j} - \frac{3\sqrt{2}}{10}\mathbf{k}$

82.  $\mathbf{v} = 5\mathbf{i} + 3\mathbf{j} - \mathbf{k}$   
 $\|\mathbf{v}\| = \sqrt{25 + 9 + 1} = \sqrt{35}$

(a)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{\sqrt{35}}(5\mathbf{i} + 3\mathbf{j} - \mathbf{k}) = \frac{\sqrt{35}}{7}\mathbf{i} + \frac{3\sqrt{35}}{35}\mathbf{j} - \frac{\sqrt{35}}{35}\mathbf{k}$   
(b)  $-\frac{\mathbf{v}}{\|\mathbf{v}\|} = -\frac{1}{\sqrt{35}}(5\mathbf{i} + 3\mathbf{j} - \mathbf{k}) = -\frac{\sqrt{35}}{7}\mathbf{i} - \frac{3\sqrt{35}}{35}\mathbf{j} + \frac{\sqrt{35}}{35}\mathbf{k}$

83.  $\mathbf{v} = 10\frac{\mathbf{u}}{\|\mathbf{u}\|} = 10\frac{\langle 0, 3, 3 \rangle}{3\sqrt{2}}$   
 $= 10\left\langle 0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle = \left\langle 0, \frac{10}{\sqrt{2}}, \frac{10}{\sqrt{2}} \right\rangle$

84.  $\mathbf{v} = 3\frac{\mathbf{u}}{\|\mathbf{u}\|} = 3\frac{\langle 1, 1, 1 \rangle}{\sqrt{3}}$   
 $= 3\left\langle \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \right\rangle = \left\langle \frac{3}{\sqrt{3}}, \frac{3}{\sqrt{3}}, \frac{3}{\sqrt{3}} \right\rangle$

77.  $\mathbf{v} = \mathbf{i} - 2\mathbf{j} - 3\mathbf{k} = \langle 1, -2, -3 \rangle$

$$\|\mathbf{v}\| = \sqrt{1 + 4 + 9} = \sqrt{14}$$

78.  $\mathbf{v} = -4\mathbf{i} + 3\mathbf{j} + 7\mathbf{k} = \langle -4, 3, 7 \rangle$   
 $\|\mathbf{v}\| = \sqrt{16 + 9 + 49} = \sqrt{74}$

79.  $\mathbf{v} = \langle 2, -1, 2 \rangle$   
 $\|\mathbf{v}\| = \sqrt{4 + 1 + 4} = 3$   
(a)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{3}\langle 2, -1, 2 \rangle$

(b)  $-\frac{\mathbf{v}}{\|\mathbf{v}\|} = -\frac{1}{3}\langle 2, -1, 2 \rangle$


80.  $\mathbf{v} = \langle 6, 0, 8 \rangle$   
 $\|\mathbf{v}\| = \sqrt{36 + 0 + 64} = 10$   
(a)  $\frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{10}\langle 6, 0, 8 \rangle$   
(b)  $-\frac{\mathbf{v}}{\|\mathbf{v}\|} = -\frac{1}{10}\langle 6, 0, 8 \rangle$

85.  $\mathbf{v} = \frac{3}{2}\frac{\mathbf{u}}{\|\mathbf{u}\|} = \frac{3}{2}\frac{\langle 2, -2, 1 \rangle}{3} = \frac{3}{2}\left\langle \frac{2}{3}, \frac{-2}{3}, \frac{1}{3} \right\rangle = \left\langle 1, -1, \frac{1}{2} \right\rangle$

86.  $\mathbf{v} = 7\frac{\mathbf{u}}{\|\mathbf{u}\|} = 7\frac{\langle -4, 6, 2 \rangle}{2\sqrt{14}} = \left\langle \frac{-14}{\sqrt{14}}, \frac{21}{\sqrt{14}}, \frac{7}{\sqrt{14}} \right\rangle$


87.  $\mathbf{v} = 2[\cos(\pm 30^\circ)\mathbf{j} + \sin(\pm 30^\circ)\mathbf{k}]$

$$= \sqrt{3}\mathbf{j} \pm \mathbf{k} = \langle 0, \sqrt{3}, \pm 1 \rangle$$



88.  $\mathbf{v} = 5(\cos 45^\circ\mathbf{i} + \sin 45^\circ\mathbf{k}) = \frac{5\sqrt{2}}{2}(\mathbf{i} + \mathbf{k})$  or

$$\mathbf{v} = 5(\cos 135^\circ\mathbf{i} + \sin 135^\circ\mathbf{k}) = \frac{5\sqrt{2}}{2}(-\mathbf{i} + \mathbf{k})$$



89.  $\mathbf{v} = \langle -3, -6, 3 \rangle$

$$\frac{2}{3}\mathbf{v} = \langle -2, -4, 2 \rangle$$

$$(4, 3, 0) + (-2, -4, 2) = (2, -1, 2)$$

90.  $\mathbf{v} = \langle 5, 6, -3 \rangle$

$$\frac{2}{3}\mathbf{v} = \left\langle \frac{10}{3}, 4, -2 \right\rangle$$

$$(1, 2, 5) + \left( \frac{10}{3}, 4, -2 \right) = \left( \frac{13}{3}, 6, 3 \right)$$

91. A sphere of radius 4 centered at  $(x_1, y_1, z_1)$ .

$$\begin{aligned} \|\mathbf{v}\| &= \|\langle x - x_1, y - y_1, z - z_1 \rangle\| \\ &= \sqrt{(x - x_1)^2 + (y - y_1)^2 + (z - z_1)^2} = 4 \end{aligned}$$

$$(x - x_1)^2 + (y - y_1)^2 + (z - z_1)^2 = 16$$

92.  $\|\mathbf{r} - \mathbf{r}_0\| = \sqrt{(x - 1)^2 + (y - 1)^2 + (z - 1)^2} = 2$   

$$(x - 1)^2 + (y - 1)^2 + (z - 1)^2 = 4$$

This is a sphere of radius 2 and center  $(1, 1, 1)$ .

93. The set of all points  $(x, y, z)$  such that  $\|\mathbf{r}\| > 1$  represent outside the sphere of radius 1 centered at the origin.

94. (a)  $(x, y, z) = (3, 3, 3)$

$$\mathbf{v} = \langle 3, 3, 3 \rangle - \langle 3, 0, 0 \rangle$$


$$= \langle 3 - 3, 3 - 0, 3 - 0 \rangle = \langle 0, 3, 3 \rangle$$

(b)  $(x, y, z) = (4, 4, 8)$


$$\mathbf{v} = \langle 4, 4, 8 \rangle - \langle 4, 0, 0 \rangle$$

$$= \langle 4 - 4, 4 - 0, 8 - 0 \rangle = \langle 0, 4, 8 \rangle$$

95. The terminal points of the vectors  $t\mathbf{v}$ ,  $\mathbf{u} + t\mathbf{v}$  and  $s\mathbf{u} + t\mathbf{v}$  are collinear.



96. (a)



(b)  $\mathbf{w} = a\mathbf{u} + b\mathbf{v} = a\mathbf{i} + (a + b)\mathbf{j} + b\mathbf{k} = \mathbf{0}$

$$a = 0, a + b = 0, b = 0$$

So,  $a$  and  $b$  are both zero.

(c)  $a\mathbf{i} + (a + b)\mathbf{j} + b\mathbf{k} = \mathbf{i} + 2\mathbf{j} + \mathbf{k}$

$$a = 1, a + b = 2, b = 1$$

$$\mathbf{w} = \mathbf{u} + \mathbf{v}$$

(d)  $a\mathbf{i} + (a + b)\mathbf{j} + b\mathbf{k} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$

$$a = 1, a + b = 2, b = 3$$

Not possible


97. Let  $\alpha$  be the angle between  $\mathbf{v}$  and the coordinate axes.

$$\mathbf{v} = (\cos \alpha)\mathbf{i} + (\cos \alpha)\mathbf{j} + (\cos \alpha)\mathbf{k}$$

$$\|\mathbf{v}\| = \sqrt{3} \cos \alpha = 1$$

$$\cos \alpha = \frac{1}{\sqrt{3}} = \frac{\sqrt{3}}{3}$$

$$\mathbf{v} = \frac{\sqrt{3}}{3}(\mathbf{i} + \mathbf{j} + \mathbf{k}) = \frac{\sqrt{3}}{3}\langle 1, 1, 1 \rangle$$



98.  $2400 = \|c(22.5\mathbf{i} - 15\mathbf{j} - 30\mathbf{k})\|$

$$5,760,000 = 1631.25c^2$$

$$c^2 \approx 3531.034$$

$$c \approx 59.423$$

$$\mathbf{F} \approx 59.423(22.5\mathbf{i} - 15\mathbf{j} - 30\mathbf{k})$$


$$\approx 1337\mathbf{i} - 891\mathbf{j} - 1783\mathbf{k}$$

99. (a) The height of the right triangle is  $h = \sqrt{L^2 - 18^2}$ .

The vector  $\overrightarrow{PQ}$  is given by  $\overrightarrow{PQ} = \langle 0, -18, h \rangle$ .

The tension vector  $\mathbf{T}$  in each wire is  $\mathbf{T} = c\langle 0, -18, h \rangle$  where  $ch = \frac{24}{3} = 8$ .


$$\text{So, } \mathbf{T} = \frac{8}{h}\langle 0, -18, h \rangle \text{ and } T = \|\mathbf{T}\| = \frac{8}{h}\sqrt{18^2 + h^2} = \frac{8}{\sqrt{L^2 - 18^2}}\sqrt{18^2 + (L^2 - 18^2)} = \frac{8L}{\sqrt{L^2 - 18^2}}, L > 18.$$



(b)

$L$	20	25	30	35	40	45	50
$T$	18.4	11.5	10	9.3	9.0	8.7	8.6

(c)



$x = 18$  is a vertical asymptote and  $y = 8$  is a horizontal asymptote.

100. As in Exercise 99(c),  $x = a$  will be a vertical asymptote. So,  $\lim_{r_0 \rightarrow a^-} T = \infty$ .

101.  $\overline{AB} = \langle 0, 70, 115 \rangle, \mathbf{F}_1 = C_1 \langle 0, 70, 115 \rangle$

$$\overline{AC} = \langle -60, 0, 115 \rangle, \mathbf{F}_2 = C_2 \langle -60, 0, 115 \rangle$$

$$\overline{AD} = \langle 45, -65, 115 \rangle, \mathbf{F}_3 = C_3 \langle 45, -65, 115 \rangle$$

$$\mathbf{F} = \mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3 = \langle 0, 0, 500 \rangle$$

So:  $-60C_2 + 45C_3 = 0$

$$70C_1 - 65C_3 = 0$$

$$115(C_1 + C_2 + C_3) = 500$$

Solving this system yields  $C_1 = \frac{104}{69}, C_2 = \frac{28}{23}$ , and  $C_3 = \frac{112}{69}$ . So:

$$\|\mathbf{F}_1\| \approx 202.919 \text{ N}$$

$$\|\mathbf{F}_2\| \approx 157.909 \text{ N}$$

$$\|\mathbf{F}_3\| \approx 226.521 \text{ N}$$

(d)  $\lim_{L \rightarrow 18^+} \frac{8L}{\sqrt{L^2 - 18^2}} = \infty$

$$\lim_{L \rightarrow \infty} \frac{8L}{\sqrt{L^2 - 18^2}} = \lim_{L \rightarrow \infty} \frac{8}{\sqrt{1 - (18/L)^2}} = 8$$

(e) From the table,  $T = 10$  implies  $L = 30$  centimeters.

102. Let  $A$  lie on the  $y$ -axis and the base of the wall on the  $x$ -axis. Then  $A = (0, 4, 0)$ ,  $B = (3, 0, 2)$ ,  $C = (-4, 0, 2)$  and  $\overline{AB} = \langle 3, -4, 2 \rangle$ ,  $\overline{AC} = \langle -4, -4, 2 \rangle$ .

$$\|AB\| = \sqrt{29}, \|AC\| = 6$$

$$\text{Thus, } \mathbf{F}_1 = 1650 \frac{\overline{AB}}{\|AB\|}, \mathbf{F}_2 = 2020 \frac{\overline{AC}}{\|AC\|}$$

$$\mathbf{F} = \mathbf{F}_1 + \mathbf{F}_2 \approx \langle 919.2, -1225.6, 612.8 \rangle + \langle -1346.7, -1346.7, 673.3 \rangle = \langle -427.5, -2572.3, 1286.1 \rangle$$

$$\|\mathbf{F}\| \approx 2907.5 \text{ N}$$

103.  $d(AP) = 2d(BP)$

$$\begin{aligned} \sqrt{x^2 + (y+1)^2 + (z-1)^2} &= 2\sqrt{(x-1)^2 + (y-2)^2 + z^2} \\ x^2 + y^2 + z^2 + 2y - 2z + 2 &= 4(x^2 + y^2 + z^2 - 2x - 4y + 5) \\ 0 &= 3x^2 + 3y^2 + 3z^2 - 8x - 18y + 2z + 18 \\ -6 + \frac{16}{9} + 9 + \frac{1}{9} &= \left( x^2 - \frac{8}{3}x + \frac{16}{9} \right) + (y^2 - 6y + 9) + \left( z^2 + \frac{2}{3}z + \frac{1}{9} \right) \\ \frac{44}{9} &= \left( x - \frac{4}{3} \right)^2 + (y-3)^2 + \left( z + \frac{1}{3} \right)^2 \end{aligned}$$

Sphere; center:  $\left(\frac{4}{3}, 3, -\frac{1}{3}\right)$ , radius:  $\frac{2\sqrt{11}}{3}$

### Section 11.3 The Dot Product of Two Vectors

1. The vectors are orthogonal (perpendicular) if the dot product of the vectors is zero.
  2. If  $\arccos \frac{v_2}{\|v\|} = 30^\circ$ , then  $\cos 30^\circ = \frac{v_2}{\|v\|}$ .  
So, the angle between  $v$  and  $j$  is  $30^\circ$ .
  3.  $\mathbf{u} = \langle 3, 4 \rangle$ ,  $\mathbf{v} = \langle -1, 5 \rangle$ 
    - (a)  $\mathbf{u} \cdot \mathbf{v} = 3(-1) + 4(5) = 17$
    - (b)  $\mathbf{u} \cdot \mathbf{u} = 3(3) + 4(4) = 25$
    - (c)  $\|\mathbf{v}\|^2 = (-1)^2 + 5^2 = 26$
    - (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = 17\langle -1, 5 \rangle = \langle -17, 85 \rangle$
    - (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(17) = 51$
  4.  $\mathbf{u} = \langle 4, 10 \rangle$ ,  $\mathbf{v} = \langle -2, 3 \rangle$ 
    - (a)  $\mathbf{u} \cdot \mathbf{v} = 4(-2) + 10(3) = 22$
    - (b)  $\mathbf{u} \cdot \mathbf{u} = 4(4) + 10(10) = 116$
    - (c)  $\|\mathbf{v}\|^2 = (-2)^2 + 3^2 = 13$
    - (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = 22\langle -2, 3 \rangle = \langle -44, 66 \rangle$
    - (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(22) = 66$
- 5.  $\mathbf{u} = \langle 6, -4 \rangle$ ,  $\mathbf{v} = \langle -3, 2 \rangle$**
- (a)  $\mathbf{u} \cdot \mathbf{v} = 6(-3) + (-4)(2) = -26$
  - (b)  $\mathbf{u} \cdot \mathbf{u} = 6(6) + (-4)(-4) = 52$
  - (c)  $\|\mathbf{v}\|^2 = (-3)^2 + 2^2 = 13$
  - (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = -26\langle -3, 2 \rangle = \langle 78, -52 \rangle$
  - (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(-26) = -78$
- 6.  $\mathbf{u} = \langle -7, -1 \rangle$ ,  $\mathbf{v} = \langle -4, -1 \rangle$**
- (a)  $\mathbf{u} \cdot \mathbf{v} = -7(-4) + -1(-1) = 29$
  - (b)  $\mathbf{u} \cdot \mathbf{u} = -7(-7) + -1(-1) = 50$
  - (c)  $\|\mathbf{v}\|^2 = (-4)^2 + (-1)^2 = 17$
  - (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = 29\langle -4, -1 \rangle = \langle -116, -29 \rangle$
  - (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(29) = 87$

7.  $\mathbf{u} = \langle 2, -3, 4 \rangle, \mathbf{v} = \langle 0, 6, 5 \rangle$

- (a)  $\mathbf{u} \cdot \mathbf{v} = 2(0) + (-3)(6) + (4)(5) = 2$
- (b)  $\mathbf{u} \cdot \mathbf{u} = 2(2) + (-3)(-3) + 4(4) = 29$
- (c)  $\|\mathbf{v}\|^2 = 0^2 + 6^2 + 5^2 = 61$
- (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = 2\langle 0, 6, 5 \rangle = \langle 0, 12, 10 \rangle$
- (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(2) = 6$

8.  $\mathbf{u} = \langle -5, 0, 5 \rangle, \mathbf{v} = \langle -1, 2, 1 \rangle$

- (a)  $\mathbf{u} \cdot \mathbf{v} = -5(-1) + 0(2) + 5(1) = 10$
- (b)  $\mathbf{u} \cdot \mathbf{u} = (-5)(-5) + (0)(0) + 5(5) = 50$
- (c)  $\|\mathbf{v}\|^2 = (-1)^2 + 2^2 + 1^2 = 6$
- (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = 10\langle -1, 2, 1 \rangle = \langle -10, 20, 10 \rangle$
- (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(10) = 30$

9.  $\mathbf{u} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}, \mathbf{v} = \mathbf{i} - \mathbf{k}$

- (a)  $\mathbf{u} \cdot \mathbf{v} = 2(1) + (-1)(0) + 1(-1) = 1$
- (b)  $\mathbf{u} \cdot \mathbf{u} = 2(2) + (-1)(-1) + (1)(1) = 6$
- (c)  $\|\mathbf{v}\|^2 = 1^2 + (-1)^2 = 2$
- (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = \mathbf{v} = \mathbf{i} - \mathbf{k}$
- (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(1) = 3$

10.  $\mathbf{u} = 2\mathbf{i} + \mathbf{j} - 2\mathbf{k}, \mathbf{v} = \mathbf{i} - 3\mathbf{j} + 2\mathbf{k}$

- (a)  $\mathbf{u} \cdot \mathbf{v} = 2(1) + 1(-3) + (-2)(2) = -5$
- (b)  $\mathbf{u} \cdot \mathbf{u} = 2(2) + 1(1) + (-2)(-2) = 9$
- (c)  $\|\mathbf{v}\|^2 = 1^2 + (-3)^2 + 2^2 = 14$
- (d)  $(\mathbf{u} \cdot \mathbf{v})\mathbf{v} = -5(\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}) = -5\mathbf{i} + 15\mathbf{j} - 10\mathbf{k}$
- (e)  $\mathbf{u} \cdot (3\mathbf{v}) = 3(\mathbf{u} \cdot \mathbf{v}) = 3(-5) = -15$

11.  $\mathbf{u} = \langle 1, 1 \rangle, \mathbf{v} = \langle 2, -2 \rangle$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{0}{\sqrt{2}\sqrt{8}} = 0$$

(a)  $\theta = \frac{\pi}{2}$       (b)  $\theta = 90^\circ$

12.  $\mathbf{u} = \langle 3, 1 \rangle, \mathbf{v} = \langle 2, -1 \rangle$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{5}{\sqrt{10}\sqrt{5}} = \frac{1}{\sqrt{2}}$$

(a)  $\theta = \frac{\pi}{4}$       (b)  $\theta = 45^\circ$

13.  $\mathbf{u} = 3\mathbf{i} + \mathbf{j}, \mathbf{v} = -2\mathbf{i} + 4\mathbf{j}$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{-2}{\sqrt{10}\sqrt{20}} = \frac{-1}{5\sqrt{2}}$$

(a)  $\theta = \arccos\left(-\frac{1}{5\sqrt{2}}\right) \approx 1.713$

(b)  $\theta \approx 98.1^\circ$

14.  $\mathbf{u} = \cos\left(\frac{\pi}{6}\right)\mathbf{i} + \sin\left(\frac{\pi}{6}\right)\mathbf{j} = \frac{\sqrt{3}}{2}\mathbf{i} + \frac{1}{2}\mathbf{j}$

$$\mathbf{v} = \cos\left(\frac{3\pi}{4}\right)\mathbf{i} + \sin\left(\frac{3\pi}{4}\right)\mathbf{j} = -\frac{\sqrt{2}}{2}\mathbf{i} + \frac{\sqrt{2}}{2}\mathbf{j}$$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}$$

$$= \frac{\sqrt{3}}{2}\left(-\frac{\sqrt{2}}{2}\right) + \frac{1}{2}\left(\frac{\sqrt{2}}{2}\right) = \frac{\sqrt{2}}{4}(1 - \sqrt{3})$$

(a)  $\theta = \arccos\left[\frac{\sqrt{2}}{4}(1 - \sqrt{3})\right] = \frac{7\pi}{2}$

(b)  $\theta = 105^\circ$

15.  $\mathbf{u} = \langle 1, 1, 1 \rangle, \mathbf{v} = \langle 2, 1, -1 \rangle$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{2}{\sqrt{3}\sqrt{6}} = \frac{\sqrt{2}}{3}$$

(a)  $\theta = \arccos\frac{\sqrt{2}}{3} \approx 1.080$

(b)  $\theta \approx 61.9^\circ$

16.  $\mathbf{u} = 3\mathbf{i} + 2\mathbf{j} + \mathbf{k}, \mathbf{v} = 2\mathbf{i} - 3\mathbf{j}$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{3(2) + 2(-3) + 0}{\|\mathbf{u}\| \|\mathbf{v}\|} = 0$$

(a)  $\theta = \frac{\pi}{2}$

(b)  $\theta = 90^\circ$

17.  $\mathbf{u} = 3\mathbf{i} + 4\mathbf{j}, \mathbf{v} = -2\mathbf{j} + 3\mathbf{k}$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{-8}{5\sqrt{13}} = \frac{-8\sqrt{13}}{65}$$

(a)  $\theta = \arccos\left(-\frac{8\sqrt{13}}{65}\right) \approx 2.031$

(b)  $\theta \approx 116.3^\circ$

18.  $\mathbf{u} = 2\mathbf{i} - 3\mathbf{j} + \mathbf{k}$ ,  $\mathbf{v} = \mathbf{i} - 2\mathbf{j} + \mathbf{k}$

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{9}{\sqrt{14}\sqrt{6}} = \frac{9}{2\sqrt{21}} = \frac{3\sqrt{21}}{14}$$

(a)  $\theta = \arccos\left(\frac{3\sqrt{21}}{14}\right) \approx 0.190$

(b)  $\theta \approx 10.9^\circ$

19.  $\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \cos \theta$

$$\mathbf{u} \cdot \mathbf{v} = (8)(5) \cos \frac{\pi}{3} = 20$$

20.  $\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \cos \theta$

$$\mathbf{u} \cdot \mathbf{v} = (40)(25) \cos \frac{5\pi}{6} = -500\sqrt{3}$$

21.  $\mathbf{u} = \langle 4, 3 \rangle$ ,  $\mathbf{v} = \langle \frac{1}{2}, -\frac{2}{3} \rangle$

$\mathbf{u} \neq c\mathbf{v} \Rightarrow$  not parallel

$\mathbf{u} \cdot \mathbf{v} = 0 \Rightarrow$  orthogonal

22.  $\mathbf{u} = -\frac{1}{3}(\mathbf{i} - 2\mathbf{j})$ ,  $\mathbf{v} = 2\mathbf{i} - 4\mathbf{j}$

$\mathbf{u} = -\frac{1}{6}\mathbf{v} \Rightarrow$  parallel

23.  $\mathbf{u} = \mathbf{j} + 6\mathbf{k}$ ,  $\mathbf{v} = \mathbf{i} - 2\mathbf{j} - \mathbf{k}$

$\mathbf{u} \neq c\mathbf{v} \Rightarrow$  not parallel

$\mathbf{u} \cdot \mathbf{v} = -8 \neq 0 \Rightarrow$  not orthogonal

Neither

24.  $\mathbf{u} = -2\mathbf{i} + 3\mathbf{j} - \mathbf{k}$ ,  $\mathbf{v} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}$

$\mathbf{u} \neq c\mathbf{v} \Rightarrow$  not parallel

$\mathbf{u} \cdot \mathbf{v} = 0 \Rightarrow$  orthogonal

25.  $\mathbf{u} = \langle 2, -3, 1 \rangle$ ,  $\mathbf{v} = \langle -1, -1, -1 \rangle$

$\mathbf{u} \neq c\mathbf{v} \Rightarrow$  not parallel

$\mathbf{u} \cdot \mathbf{v} = 0 \Rightarrow$  orthogonal

26.  $\mathbf{u} = \langle \cos \theta, \sin \theta, -1 \rangle$ ,

$$\mathbf{v} = \langle \sin \theta, -\cos \theta, 0 \rangle$$

$\mathbf{u} \neq c\mathbf{v} \Rightarrow$  not parallel

$\mathbf{u} \cdot \mathbf{v} = 0 \Rightarrow$  orthogonal

27. The vector  $\langle 1, 2, 0 \rangle$  joining  $(1, 2, 0)$  and  $(0, 0, 0)$  is

perpendicular to the vector  $\langle -2, 1, 0 \rangle$  joining  $(-2, 1, 0)$  and  $(0, 0, 0)$ :  $\langle 1, 2, 0 \rangle \cdot \langle -2, 1, 0 \rangle = 0$

The triangle has a right angle, so it is a right triangle.

28. Consider the vector  $\langle -3, 0, 0 \rangle$  joining  $(0, 0, 0)$  and  $(-3, 0, 0)$ , and the vector  $\langle 1, 2, 3 \rangle$  joining  $(0, 0, 0)$  and  $(1, 2, 3)$ :  $\langle -3, 0, 0 \rangle \cdot \langle 1, 2, 3 \rangle = -3 < 0$

The triangle has an obtuse angle, so it is an obtuse triangle.

29.  $A(2, 0, 1)$ ,  $B(0, 1, 2)$ ,  $C(-\frac{1}{2}, \frac{3}{2}, 0)$

$$\overrightarrow{AB} = \langle -2, 1, 1 \rangle \quad \overrightarrow{BA} = \langle 2, -1, -1 \rangle$$

$$\overrightarrow{AC} = \left\langle -\frac{5}{2}, \frac{3}{2}, -1 \right\rangle \quad \overrightarrow{CA} = \left\langle \frac{5}{2}, -\frac{3}{2}, 1 \right\rangle$$

$$\overrightarrow{BC} = \left\langle -\frac{1}{2}, \frac{1}{2}, -2 \right\rangle \quad \overrightarrow{CB} = \left\langle \frac{1}{2}, -\frac{1}{2}, 2 \right\rangle$$

$$\overrightarrow{AB} \cdot \overrightarrow{AC} = 5 + \frac{3}{2} - 1 > 0$$

$$\overrightarrow{BA} \cdot \overrightarrow{BC} = -1 - \frac{1}{2} + 2 > 0$$

$$\overrightarrow{CA} \cdot \overrightarrow{CB} = \frac{5}{4} + \frac{3}{4} + 2 > 0$$

The triangle has three acute angles, so it is an acute triangle.

30.  $A(2, -7, 3)$ ,  $B(-1, 5, 8)$ ,  $C(4, 6, -1)$

$$\overrightarrow{AB} = \langle -3, 12, 5 \rangle \quad \overrightarrow{BA} = \langle 3, -12, -5 \rangle$$

$$\overrightarrow{AC} = \langle 2, 13, -4 \rangle \quad \overrightarrow{CA} = \langle -2, -13, 4 \rangle$$

$$\overrightarrow{BC} = \langle 5, 1, -9 \rangle \quad \overrightarrow{CB} = \langle -5, -1, 9 \rangle$$

$$\overrightarrow{AB} \cdot \overrightarrow{AC} = -6 + 156 - 20 > 0$$

$$\overrightarrow{BA} \cdot \overrightarrow{BC} = 15 - 12 + 45 > 0$$

$$\overrightarrow{CA} \cdot \overrightarrow{CB} = 10 + 13 + 36 > 0$$

The triangle has three acute angles, so it is an acute triangle.

31.  $\mathbf{u} = \mathbf{i} + 2\mathbf{j} + 2\mathbf{k}$ ,  $\|\mathbf{u}\| = \sqrt{1 + 4 + 4} = 3$

$$\cos \alpha = \frac{1}{3} \Rightarrow \alpha \approx 1.2310 \text{ or } 70.5^\circ$$

$$\cos \beta = \frac{2}{3} \Rightarrow \beta \approx 0.8411 \text{ or } 48.2^\circ$$

$$\cos \gamma = \frac{2}{3} \Rightarrow \gamma \approx 0.8411 \text{ or } 48.2^\circ$$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{1}{9} + \frac{4}{9} + \frac{4}{9} = 1$$

32.  $\mathbf{u} = 5\mathbf{i} + 3\mathbf{j} - \mathbf{k}$ ,  $\|\mathbf{u}\| = \sqrt{25 + 9 + 1} = \sqrt{35}$

$$\cos \alpha = \frac{5}{\sqrt{35}} \Rightarrow \alpha \approx 0.5639 \text{ or } 32.3^\circ$$

$$\cos \beta = \frac{3}{\sqrt{35}} \Rightarrow \beta \approx 1.0390 \text{ or } 59.5^\circ$$

$$\cos \gamma = \frac{-1}{\sqrt{35}} \Rightarrow \gamma \approx 1.7406 \text{ or } 99.7^\circ$$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{25}{35} + \frac{9}{35} + \frac{1}{35} = 1$$

33.  $\mathbf{u} = 7\mathbf{i} + \mathbf{j} - \mathbf{k}$ ,  $\|\mathbf{u}\| = \sqrt{49 + 1 + 1} = \sqrt{51}$

$$\cos \alpha = \frac{7}{\sqrt{51}} \Rightarrow \alpha \approx 11.4^\circ$$

$$\cos \beta = \frac{1}{\sqrt{51}} \Rightarrow \beta \approx 82.0^\circ$$

$$\cos \gamma = -\frac{1}{\sqrt{51}} \Rightarrow \gamma \approx 98.0^\circ$$

34.  $\mathbf{u} = -4\mathbf{i} + 3\mathbf{j} + 5\mathbf{k}$ ,  $\|\mathbf{u}\| = \sqrt{16 + 9 + 25} = \sqrt{50} = 5\sqrt{2}$

$$\cos \alpha = \frac{-4}{5\sqrt{2}} \Rightarrow \alpha \approx 2.1721 \text{ or } 124.4^\circ$$

$$\cos \beta = \frac{3}{5\sqrt{2}} \Rightarrow \beta \approx 1.1326 \text{ or } 64.9^\circ$$

$$\cos \gamma = \frac{5}{5\sqrt{2}} = \frac{1}{\sqrt{2}} \Rightarrow \gamma \approx \frac{\pi}{4} \text{ or } 45^\circ$$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{16}{50} + \frac{9}{50} + \frac{25}{50} = 1$$

35.  $\mathbf{u} = \langle 0, 6, -4 \rangle$ ,  $\|\mathbf{u}\| = \sqrt{0 + 36 + 16} = \sqrt{52} = 2\sqrt{13}$

$$\cos \alpha = 0 \Rightarrow \alpha = \frac{\pi}{2} \text{ or } 90^\circ$$

$$\cos \beta = \frac{3}{\sqrt{13}} \Rightarrow \beta \approx 0.5880 \text{ or } 33.7^\circ$$

$$\cos \gamma = -\frac{2}{\sqrt{13}} \Rightarrow \gamma \approx 2.1588 \text{ or } 123.7^\circ$$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 0 + \frac{9}{13} + \frac{4}{13} = 1$$

36.  $\mathbf{u} = \langle -1, 5, 2 \rangle$ ,  $\|\mathbf{u}\| = \sqrt{1 + 25 + 4} = \sqrt{30}$

$$\cos \alpha = \frac{-1}{\sqrt{30}} \Rightarrow \alpha \approx 1.7544 \text{ or } 100.5^\circ$$

$$\cos \beta = \frac{5}{\sqrt{30}} \Rightarrow \beta \approx 0.4205 \text{ or } 24.1^\circ$$

$$\cos \gamma = \frac{2}{\sqrt{30}} \Rightarrow \gamma \approx 1.1970 \text{ or } 68.6^\circ$$

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \frac{1}{30} + \frac{25}{30} + \frac{4}{30} = 1$$

37.  $\mathbf{u} = \langle 6, 7 \rangle$ ,  $\mathbf{v} = \langle 1, 4 \rangle$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \frac{6(1) + 7(4)}{1^2 + 4^2} \langle 1, 4 \rangle \\ &= \frac{34}{17} \langle 1, 4 \rangle = \langle 2, 8 \rangle \end{aligned}$$

$$\text{(b)} \quad \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 6, 7 \rangle - \langle 2, 8 \rangle = \langle 4, -1 \rangle$$

38.  $\mathbf{u} = \langle 9, 7 \rangle$ ,  $\mathbf{v} = \langle 1, 3 \rangle$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \frac{9(1) + 7(3)}{1 + 3^2} \langle 1, 3 \rangle \\ &= \frac{30}{10} \langle 1, 3 \rangle = \langle 3, 9 \rangle \end{aligned}$$

$$\text{(b)} \quad \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 9, 7 \rangle - \langle 3, 9 \rangle = \langle 6, -2 \rangle$$

39.  $\mathbf{u} = 2\mathbf{i} + 3\mathbf{j} = \langle 2, 3 \rangle$ ,  $\mathbf{v} = 5\mathbf{i} + \mathbf{j} = \langle 5, 1 \rangle$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \frac{2(5) + 3(1)}{5^2 + 1} \langle 5, 1 \rangle \\ &= \frac{13}{26} \langle 5, 1 \rangle = \left\langle \frac{5}{2}, \frac{1}{2} \right\rangle \end{aligned}$$

$$\text{(b)} \quad \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 2, 3 \rangle - \left\langle \frac{5}{2}, \frac{1}{2} \right\rangle = \left\langle -\frac{1}{2}, \frac{5}{2} \right\rangle$$

40.  $\mathbf{u} = 2\mathbf{i} - 3\mathbf{j} = \langle 2, -3 \rangle$ ,  $\mathbf{v} = 3\mathbf{i} + 2\mathbf{j} = \langle 3, 2 \rangle$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \frac{2(3) + (-3)(2)}{3^2 + 2^2} \langle 3, 2 \rangle \\ &= 0 \langle 3, 2 \rangle = \langle 0, 0 \rangle \end{aligned}$$

$$\text{(b)} \quad \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 2, -3 \rangle$$

41.  $\mathbf{u} = \langle 0, 3, 3 \rangle, \mathbf{v} = \langle -1, 1, 1 \rangle$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \frac{0(-1) + 3(1) + 3(1)}{1+1+1} \langle -1, 1, 1 \rangle \\ &= \frac{6}{3} \langle -1, 1, 1 \rangle = \langle -2, 2, 2 \rangle \end{aligned}$$

$$\text{(b)} \quad \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 0, 3, 3 \rangle - \langle -2, 2, 2 \rangle = \langle 2, 1, 1 \rangle$$

42.  $\mathbf{u} = \langle 8, 2, 0 \rangle, \mathbf{v} = \langle 2, 1, -1 \rangle$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \frac{8(2) + 2(1) + 0(-1)}{2^2 + 1 + 1} \langle 2, 1, -1 \rangle \\ &= \frac{18}{6} \langle 2, 1, -1 \rangle = \langle 6, 3, -3 \rangle \end{aligned}$$

$$\text{(b)} \quad \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 8, 2, 0 \rangle - \langle 6, 3, -3 \rangle = \langle 2, -1, 3 \rangle$$

44.  $\mathbf{u} = 5\mathbf{i} - \mathbf{j} - \mathbf{k}, \mathbf{v} = -\mathbf{i} + 5\mathbf{j} + 8\mathbf{k}$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \left( \frac{5(-1) + (-1)(5) + (-1)(8)}{(-1)^2 + 5^2 + 8^2} \right) \langle -1, 5, 8 \rangle \\ &= \frac{-18}{90} \langle -1, 5, 8 \rangle \\ &= \frac{-1}{5} \langle -1, 5, 8 \rangle \\ &= \left\langle \frac{1}{5}, -1, -\frac{8}{5} \right\rangle \end{aligned}$$

45.  $\mathbf{u}$  is a vector and  $\mathbf{v} \cdot \mathbf{w}$  is a scalar. You cannot add a vector and a scalar.

46.  $\left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} = \mathbf{u} \Rightarrow \mathbf{u} = c\mathbf{v} \Rightarrow \mathbf{u}$  and  $\mathbf{v}$  are parallel.

$$\begin{aligned} \text{47. Yes, } \quad &\left\| \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v} \right\| = \left\| \frac{\mathbf{v} \cdot \mathbf{u}}{\|\mathbf{u}\|^2} \mathbf{u} \right\| \\ &| \mathbf{u} \cdot \mathbf{v} | \frac{\|\mathbf{v}\|}{\|\mathbf{v}\|^2} = | \mathbf{v} \cdot \mathbf{u} | \frac{\|\mathbf{u}\|}{\|\mathbf{u}\|^2} \\ &\frac{1}{\|\mathbf{v}\|} = \frac{1}{\|\mathbf{u}\|} \\ &\|\mathbf{u}\| = \|\mathbf{v}\| \end{aligned}$$

43.  $\mathbf{u} = -9\mathbf{i} - 2\mathbf{j} - 4\mathbf{k}, \mathbf{v} = 4\mathbf{j} + 4\mathbf{k}$

$$\begin{aligned} \text{(a)} \quad \mathbf{w}_1 &= \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v} \\ &= \left( \frac{(-2)(4) + (-4)(4)}{4^2 + 4^2} \right) \langle 0, 4, 4 \rangle \\ &= -\frac{3}{4} \langle 0, 4, 4 \rangle \\ &= \langle 0, -3, -3 \rangle \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \mathbf{w}_2 &= \mathbf{u} - \mathbf{w}_1 \\ &= \langle -9, -2, -4 \rangle - \langle 0, -3, -3 \rangle \\ &= \langle -9, 1, -1 \rangle \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \mathbf{w}_2 &= \mathbf{u} - \mathbf{w}_1 \\ &= \langle 5, -1, -1 \rangle - \left\langle \frac{1}{5}, -1, -\frac{8}{5} \right\rangle \\ &= \left\langle \frac{24}{5}, 0, \frac{3}{5} \right\rangle \end{aligned}$$

48. (a) Orthogonal,  $\theta = \frac{\pi}{2}$

(b) Acute,  $0 < \theta < \frac{\pi}{2}$

(c) Obtuse,  $\frac{\pi}{2} < \theta < \pi$

49.  $\mathbf{u} = \langle 3240, 1450, 2235 \rangle$

$$\mathbf{v} = \langle 2.25, 2.95, 2.65 \rangle$$

$$\begin{aligned} \mathbf{u} \cdot \mathbf{v} &= 3240(2.25) + 1450(2.95) + 2235(2.65) \\ &= \$17,490.25 \end{aligned}$$

This represents the total revenue the restaurant earned on its three products.

50.  $\mathbf{u} = \langle 3240, 1450, 2235 \rangle$

$\mathbf{v} = \langle 2.25, 2.95, 2.65 \rangle$

Decrease prices by 2%:  $0.98\mathbf{v}$

New total revenue:

$$0.98\langle 3240, 1450, 2235 \rangle \cdot \langle 2.25, 2.95, 2.65 \rangle = 0.98(17490.25)$$

$$= \$17,140.45$$

51. Answers will vary. *Sample answer:*

$$\mathbf{u} = -\frac{1}{4}\mathbf{i} + \frac{3}{2}\mathbf{j}. \text{ Want } \mathbf{u} \cdot \mathbf{v} = 0.$$

$\mathbf{v} = 12\mathbf{i} + 2\mathbf{j}$  and  $-\mathbf{v} = -12\mathbf{i} - 2\mathbf{j}$  are orthogonal to  $\mathbf{u}$ .

52. Answers will vary. *Sample answer:*

$$\mathbf{u} = 9\mathbf{i} - 4\mathbf{j}. \text{ Want } \mathbf{u} \cdot \mathbf{v} = 0.$$

$$\mathbf{v} = 4\mathbf{i} + 9\mathbf{j} \text{ and } -\mathbf{v} = -4\mathbf{i} - 9\mathbf{j}$$

are orthogonal to  $\mathbf{u}$ .

53. Answers will vary. *Sample answer:*

$$\mathbf{u} = \langle 3, 1, -2 \rangle. \text{ Want } \mathbf{u} \cdot \mathbf{v} = 0.$$

$$\mathbf{v} = \langle 0, 2, 1 \rangle \text{ and } -\mathbf{v} = \langle 0, -2, -1 \rangle \text{ are orthogonal to } \mathbf{u}.$$

54. Answers will vary. *Sample answer:*

$$\mathbf{u} = \langle 4, -3, 6 \rangle. \text{ Want } \mathbf{u} \cdot \mathbf{v} = 0$$

$$\mathbf{v} = \langle 0, 6, 3 \rangle \text{ and } -\mathbf{v} = \langle 0, -6, -3 \rangle$$

are orthogonal to  $\mathbf{u}$ .


55. Let  $s$  = length of a side.

$$\mathbf{v} = \langle s, s, s \rangle$$

$$\|\mathbf{v}\| = s\sqrt{3}$$

$$\cos \alpha = \cos \beta = \cos \gamma = \frac{s}{s\sqrt{3}} = \frac{1}{\sqrt{3}}$$

$$\alpha = \beta = \gamma = \arccos\left(\frac{1}{\sqrt{3}}\right) \approx 54.7^\circ$$



56.  $\mathbf{v}_1 = \langle s, s, s \rangle$


$$\|\mathbf{v}_1\| = s\sqrt{3}$$

$$\mathbf{v}_2 = \langle s, s, 0 \rangle$$

$$\|\mathbf{v}_2\| = s\sqrt{2}$$

$$\cos \theta = \frac{2\sqrt{2}}{2\sqrt{3}} = \frac{\sqrt{6}}{3}$$

$$\theta = \arccos \frac{\sqrt{6}}{3} \approx 35.26^\circ$$



57. (a) Gravitational Force  $\mathbf{F} = -22\mathbf{j}$

$$\mathbf{v} = \cos 10^\circ \mathbf{i} + \sin 10^\circ \mathbf{j}$$

$$\mathbf{w}_1 = \frac{\mathbf{F} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v} = (\mathbf{F} \cdot \mathbf{v})\mathbf{v}$$

$$= (-22)(\sin 10^\circ)\mathbf{v}$$

$$\approx -3.82(\cos 10^\circ \mathbf{i} + \sin 10^\circ \mathbf{j})$$

$$\|\mathbf{w}_1\| \approx 3.82 \text{ metric tons}$$

(b)  $\mathbf{w}_2 = \mathbf{F} - \mathbf{w}_1$

$$\approx -22\mathbf{j} + 3.82(\cos 10^\circ \mathbf{i} + \sin 10^\circ \mathbf{j})$$

$$\approx 3.76\mathbf{i} - 21.34\mathbf{j}$$

$$\|\mathbf{w}_2\| \approx 21.67 \text{ metric tons}$$

58. (a) Gravitational Force  $\mathbf{F} = -24,000\mathbf{j}$

$$\mathbf{v} = \cos 18^\circ \mathbf{i} + \sin 18^\circ \mathbf{j}$$

$$\mathbf{w}_1 = \frac{\mathbf{F} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v} = (\mathbf{F} \cdot \mathbf{v})\mathbf{v}$$

$$= (-24,000)(\sin 18^\circ)\mathbf{v}$$

$$\approx -7416.4(\cos 18^\circ \mathbf{i} + \sin 18^\circ \mathbf{j})$$

$$\|\mathbf{w}_1\| = 7416.4 \text{ N}$$

(b)  $\mathbf{w}_2 = \mathbf{F} - \mathbf{w}_1$

$$= -24,000\mathbf{j} + 7416.4(\cos 18^\circ \mathbf{i} + \sin 18^\circ \mathbf{j})$$

$$\approx 7053.4\mathbf{i} - 21,708.2\mathbf{j}$$

$$\|\mathbf{w}_2\| \approx 22,825.3 \text{ N}$$

59.  $\mathbf{F} = 85\left(\frac{1}{2}\mathbf{i} + \frac{\sqrt{3}}{2}\mathbf{j}\right)$

$\mathbf{v} = 10\mathbf{i}$

$W = \mathbf{F} \cdot \mathbf{v} = 425 \text{ J}$

60.  $W = \|\text{proj}_{\overrightarrow{PQ}} \mathbf{F}\| \|\overrightarrow{PQ}\|$

$= \cos 20^\circ \|\mathbf{F}\| |\overrightarrow{PQ}|$

$= (\cos 20^\circ)(65)(50)$

$\approx 3054.0 \text{ J}$

61.  $\mathbf{F} = 1600(\cos 25^\circ \mathbf{i} + \sin 25^\circ \mathbf{j})$

$\mathbf{v} = 2000\mathbf{i}$

$W = \mathbf{F} \cdot \mathbf{v} = 1600(2000)\cos 25^\circ$

$\approx 2,900,184.9 \text{ Newton-meters (joules)}$

$\approx 2900.2 \text{ kJ}$

62.  $W = \|\text{proj}_{\overrightarrow{PQ}} \mathbf{F}\| \|\overrightarrow{PQ}\|$

$= (\cos 60^\circ) \|\mathbf{F}\| \|\overrightarrow{PQ}\|$

$= \frac{1}{2}(400)(40)$

$= 8000 \text{ J}$

63. False.

For example, let  $\mathbf{u} = \langle 1, 1 \rangle$ ,  $\mathbf{v} = \langle 2, 3 \rangle$  and  $\mathbf{w} = \langle 1, 4 \rangle$ .

Then  $\mathbf{u} \cdot \mathbf{v} = 2 + 3 = 5$  and  $\mathbf{u} \cdot \mathbf{w} = 1 + 4 = 5$ .

64. True

$\mathbf{w} \cdot (\mathbf{u} + \mathbf{v}) = \mathbf{w} \cdot \mathbf{u} + \mathbf{w} \cdot \mathbf{v} = 0 + 0 = 0$  so,  $\mathbf{w}$  and  $\mathbf{u} + \mathbf{v}$  are orthogonal.

65. (a) The graphs  $y_1 = x^2$  and  $y_2 = x^{1/3}$  intersect at  $(0, 0)$  and  $(1, 1)$ .

(b)  $y'_1 = 2x$  and  $y'_2 = \frac{1}{3x^{2/3}}$ .

At  $(0, 0)$ ,  $\pm \langle 1, 0 \rangle$  is tangent to  $y_1$  and  $\pm \langle 0, 1 \rangle$  is tangent to  $y_2$ .

At  $(1, 1)$ ,  $y'_1 = 2$  and  $y'_2 = \frac{1}{3}$ .


$\pm \frac{1}{\sqrt{5}} \langle 1, 2 \rangle$  is tangent to  $y_1$ ,  $\pm \frac{1}{\sqrt{10}} \langle 3, 1 \rangle$  is tangent to  $y_2$ .

(c) At  $(0, 0)$ , the vectors are perpendicular ( $90^\circ$ ).

At  $(1, 1)$ ,

$$\cos \theta = \frac{\frac{1}{\sqrt{5}} \langle 1, 2 \rangle \cdot \frac{1}{\sqrt{10}} \langle 3, 1 \rangle}{(1)(1)} = \frac{5}{\sqrt{50}} = \frac{1}{\sqrt{2}}$$

$$\theta = 45^\circ$$



66. (a) The graphs  $y_1 = x^3$  and  $y_2 = x^{1/3}$  intersect at  $(-1, -1)$ ,  $(0, 0)$  and  $(1, 1)$ .

(b)  $y'_1 = 3x^2$  and  $y'_2 = \frac{1}{3x^{2/3}}$ .


At  $(0, 0)$ ,  $\pm \langle 1, 0 \rangle$  is tangent to  $y_1$  and  $\pm \langle 0, 1 \rangle$  is tangent to  $y_2$ .

At  $(1, 1)$ ,  $y'_1 = 3$  and  $y'_2 = \frac{1}{3}$ .

$\pm \frac{1}{\sqrt{10}} \langle 1, 3 \rangle$  is tangent to  $y_1$ ,  $\pm \frac{1}{\sqrt{10}} \langle 3, 1 \rangle$  is tangent to  $y_2$ .

At  $(-1, -1)$ ,  $y'_1 = 3$  and  $y'_2 = \frac{1}{3}$ .

$\pm \frac{1}{\sqrt{10}} \langle 1, 3 \rangle$  is tangent to  $y_1$ ,  $\pm \frac{1}{\sqrt{10}} \langle 3, 1 \rangle$  is tangent to  $y_2$ .



(c) At  $(0, 0)$ , the vectors are perpendicular ( $90^\circ$ ).

At  $(1, 1)$ ,

$$\cos \theta = \frac{\frac{1}{\sqrt{10}} \langle 1, 3 \rangle \cdot \frac{1}{\sqrt{10}} \langle 3, 1 \rangle}{(1)(1)} = \frac{6}{10} = \frac{3}{5}.$$

$\theta \approx 0.9273 \text{ or } 53.13^\circ$

By symmetry, the angle is the same at  $(-1, -1)$ .

67. (a) The graphs of  $y_1 = 1 - x^2$  and  $y^2 = x^2 - 1$  intersect at  $(1, 0)$  and  $(-1, 0)$ .

(b)  $y'_1 = -2x$  and  $y'_2 = 2x$ .

At  $(1, 0)$ ,  $y'_1 = -2$  and  $y'_2 = 2$ .  $\pm \frac{1}{\sqrt{5}}\langle 1, -2 \rangle$  is tangent to  $y_1$ ,  $\pm \frac{1}{\sqrt{5}}\langle 1, 2 \rangle$  is tangent to  $y_2$ .

At  $(-1, 0)$ ,  $y'_1 = 2$  and  $y'_2 = -2$ .  $\pm \frac{1}{\sqrt{5}}\langle 1, 2 \rangle$  is tangent to  $y_1$ ,  $\pm \frac{1}{\sqrt{5}}\langle 1, -2 \rangle$  is tangent to  $y_2$ .

(c) At  $(1, 0)$ ,  $\cos \theta = \frac{1}{\sqrt{5}}\langle 1, -2 \rangle \cdot \frac{-1}{\sqrt{5}}\langle 1, -2 \rangle = \frac{3}{5}$ .


$\theta \approx 0.9273$  or  $53.13^\circ$

By symmetry, the angle is the same at  $(-1, 0)$ .

68. (a) To find the intersection points, rewrite the second equation as  $y + 1 = x^3$ . Substituting into the first equation

$$(y + 1)^2 = x \Rightarrow x^6 = x \Rightarrow x = 0, 1.$$

There are two points of intersection,  $(0, -1)$  and  $(1, 0)$ , as indicated in the figure.



(b) First equation:

$$(y + 1)^2 = x \Rightarrow 2(y + 1)y' = 1 \Rightarrow y' = \frac{1}{2(y + 1)}$$

At  $(1, 0)$ ,  $y' = \frac{1}{2}$ .

Second equation:  $y = x^3 - 1 \Rightarrow y' = 3x^2$ . At  $(1, 0)$ ,  $y' = 3$ .

$\pm \frac{1}{\sqrt{5}}\langle 2, 1 \rangle$  unit tangent vectors to first curve,

$\pm \frac{1}{\sqrt{10}}\langle 1, 3 \rangle$  unit tangent vectors to second curve

At  $(0, 1)$ , the unit tangent vectors to the first curve

are  $\pm \langle 0, 1 \rangle$ , and the unit tangent vectors to the second curve are  $\pm \langle 1, 0 \rangle$ .

(c) At  $(1, 0)$ ,

$$\cos \theta = \frac{1}{\sqrt{5}}\langle 2, 1 \rangle \cdot \frac{1}{\sqrt{10}}\langle 1, 3 \rangle = \frac{5}{\sqrt{50}} = \frac{1}{\sqrt{2}}$$


$\theta \approx \frac{\pi}{4}$  or  $45^\circ$

At  $(0, -1)$  the vectors are perpendicular,  $\theta = 90^\circ$ .

69. In a rhombus,  $\|\mathbf{u}\| = \|\mathbf{v}\|$ . The diagonals are  $\mathbf{u} + \mathbf{v}$  and  $\mathbf{u} - \mathbf{v}$ .

$$\begin{aligned} (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v}) &= (\mathbf{u} + \mathbf{v}) \cdot \mathbf{u} - (\mathbf{u} + \mathbf{v}) \cdot \mathbf{v} \\ &= \mathbf{u} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{u} - \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{v} \\ &= \|\mathbf{u}\|^2 - \|\mathbf{v}\|^2 = 0 \end{aligned}$$

So, the diagonals are orthogonal.



70. If  $\mathbf{u}$  and  $\mathbf{v}$  are the sides of the parallelogram, then the diagonals are  $\mathbf{u} + \mathbf{v}$  and  $\mathbf{u} - \mathbf{v}$ , as indicated in the figure.

the parallelogram is a rectangle.


$$\Leftrightarrow \mathbf{u} \cdot \mathbf{v} = 0$$

$$\Leftrightarrow 2\mathbf{u} \cdot \mathbf{v} = -2\mathbf{u} \cdot \mathbf{v}$$


$$\Leftrightarrow (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v}) = (\mathbf{u} - \mathbf{v}) \cdot (\mathbf{u} - \mathbf{v})$$

$$\Leftrightarrow \|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u} - \mathbf{v}\|^2$$

$\Leftrightarrow$  The diagonals are equal in length.



71. (a)



(b) Length of each edge:  $\sqrt{k^2 + k^2 + 0^2} = k\sqrt{2}$

(c)  $\cos \theta = \frac{k^2}{(k\sqrt{2})(k\sqrt{2})} = \frac{1}{2}$

$\theta = \arccos\left(\frac{1}{2}\right) = 60^\circ$

(d)  $\vec{r}_1 = \langle k, k, 0 \rangle - \left\langle \frac{k}{2}, \frac{k}{2}, \frac{k}{2} \right\rangle = \left\langle \frac{k}{2}, \frac{k}{2}, -\frac{k}{2} \right\rangle$

$\vec{r}_2 = \langle 0, 0, 0 \rangle - \left\langle \frac{k}{2}, \frac{k}{2}, \frac{k}{2} \right\rangle = \left\langle -\frac{k}{2}, -\frac{k}{2}, -\frac{k}{2} \right\rangle$

$\cos \theta = \frac{-\frac{k^2}{4}}{\left(\frac{k}{2}\right)^2 \cdot 3} = -\frac{1}{3}$

$\theta = 109.5^\circ$

72.  $\mathbf{u} = \langle \cos \alpha, \sin \alpha, 0 \rangle, \mathbf{v} = \langle \cos \beta, \sin \beta, 0 \rangle$

The angle between  $\mathbf{u}$  and  $\mathbf{v}$  is  $\alpha - \beta$ . (Assuming that  $\alpha > \beta$ ). Also,


$$\begin{aligned}\cos(\alpha - \beta) &= \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} \\ &= \frac{\cos \alpha \cos \beta + \sin \alpha \sin \beta}{(1)(1)} \\ &= \cos \alpha \cos \beta + \sin \alpha \sin \beta.\end{aligned}$$

## Section 11.4 The Cross Product of Two Vectors in Space

1.  $\mathbf{u} \times \mathbf{v}$  is a vector that is perpendicular (orthogonal) to both  $\mathbf{u}$  and  $\mathbf{v}$ .

2. If  $\mathbf{u}$  and  $\mathbf{v}$  are the adjacent sides of a parallelogram, then  $A = \|\mathbf{u} \times \mathbf{v}\|$ .

3.  $\mathbf{j} \times \mathbf{i} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{vmatrix} = -\mathbf{k}$



73.  $\|\mathbf{u} + \mathbf{v}\|^2 = (\mathbf{u} + \mathbf{v}) \cdot (\mathbf{u} + \mathbf{v})$

$= (\mathbf{u} + \mathbf{v}) \cdot \mathbf{u} + (\mathbf{u} + \mathbf{v}) \cdot \mathbf{v}$

$= \mathbf{u} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{v}$

$= \|\mathbf{u}\|^2 + 2\mathbf{u} \cdot \mathbf{v} + \|\mathbf{v}\|^2$

$\leq \|\mathbf{u}\|^2 + 2\|\mathbf{u}\|\|\mathbf{v}\| + \|\mathbf{v}\|^2 \leq (\|\mathbf{u}\| + \|\mathbf{v}\|)^2$

So,  $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ .

74. Let  $\mathbf{w}_1 = \text{proj}_{\mathbf{v}} \mathbf{u}$ , as indicated in the figure. Because  $\mathbf{w}_1$  is a scalar multiple of  $\mathbf{v}$ , you can write

$\mathbf{u} = \mathbf{w}_1 + \mathbf{w}_2 = c\mathbf{v} + \mathbf{w}_2.$


Taking the dot product of both sides with  $\mathbf{v}$  produces

$\mathbf{u} \cdot \mathbf{v} = (c\mathbf{v} + \mathbf{w}_2) \cdot \mathbf{v} = c\mathbf{v} \cdot \mathbf{v} + \mathbf{w}_2 \cdot \mathbf{v}$

$= c\|\mathbf{v}\|^2, \text{ because } \mathbf{w}_2 \text{ and } \mathbf{v} \text{ are orthogonal.}$

So,  $\mathbf{u} \cdot \mathbf{v} = c\|\mathbf{v}\|^2 \Rightarrow c = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}$  and

$\mathbf{w}_1 = \text{proj}_{\mathbf{v}} \mathbf{u} = c\mathbf{v} = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \mathbf{v}.$




75.  $\mathbf{u} \cdot \mathbf{v} = \|\mathbf{u}\| \|\mathbf{v}\| \cos \theta$

$|\mathbf{u} \cdot \mathbf{v}| = |\|\mathbf{u}\| \|\mathbf{v}\| \cos \theta|$


$= \|\mathbf{u}\| \|\mathbf{v}\| |\cos \theta|$

$\leq \|\mathbf{u}\| \|\mathbf{v}\| \text{ because } |\cos \theta| \leq 1.$


4.  $\mathbf{j} \times \mathbf{k} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = \mathbf{i}$



5.  $\mathbf{i} \times \mathbf{k} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{vmatrix} = -\mathbf{j}$



6.  $\mathbf{k} \times \mathbf{i} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{vmatrix} = \mathbf{j}$



11.  $\mathbf{u} = \langle 4, -1, 0 \rangle, \mathbf{v} = \langle -6, 3, 0 \rangle$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & -1 & 0 \\ -6 & 3 & 0 \end{vmatrix} = 6\mathbf{k} = \langle 0, 0, 6 \rangle$$

$$\mathbf{u} \cdot (\mathbf{u} \times \mathbf{v}) = 4(0) + (-1)(0) + 0(6) = 0 \Rightarrow \mathbf{u} \perp \mathbf{u} \times \mathbf{v}$$

$$\mathbf{v} \cdot (\mathbf{u} \times \mathbf{v}) = -6(0) + 3(0) + 0(6) = 0 \Rightarrow \mathbf{v} \perp \mathbf{u} \times \mathbf{v}$$

12.  $\mathbf{u} = \langle -5, 2, 2 \rangle, \mathbf{v} = \langle 0, 1, 8 \rangle$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -5 & 2 & 2 \\ 0 & 1 & 8 \end{vmatrix} = 14\mathbf{i} + 40\mathbf{j} - 5\mathbf{k} = \langle 14, 40, -5 \rangle$$

$$\mathbf{u} \cdot (\mathbf{u} \times \mathbf{v}) = (-5)(14) + 2(40) + 2(-5) = 0 \Rightarrow \mathbf{u} \perp \mathbf{u} \times \mathbf{v}$$

$$\mathbf{v} \cdot (\mathbf{u} \times \mathbf{v}) = (0)(14) + 1(40) + 8(-5) = 0 \Rightarrow \mathbf{v} \perp \mathbf{u} \times \mathbf{v}$$

7. (a)  $\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 4 & 0 \\ 3 & 2 & 5 \end{vmatrix} = 20\mathbf{i} + 10\mathbf{j} - 16\mathbf{k}$

(b)  $\mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = -20\mathbf{i} - 10\mathbf{j} + 16\mathbf{k}$

(c)  $\mathbf{v} \times \mathbf{v} = \mathbf{0}$

8. (a)  $\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 0 & 5 \\ 2 & 3 & -2 \end{vmatrix} = -15\mathbf{i} + 16\mathbf{j} + 9\mathbf{k}$

(b)  $\mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = 15\mathbf{i} - 16\mathbf{j} - 9\mathbf{k}$

(c)  $\mathbf{v} \times \mathbf{v} = \mathbf{0}$

9. (a)  $\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 7 & 3 & 2 \\ 1 & -1 & 5 \end{vmatrix} = 17\mathbf{i} - 33\mathbf{j} - 10\mathbf{k}$

(b)  $\mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = -17\mathbf{i} + 33\mathbf{j} + 10\mathbf{k}$

(c)  $\mathbf{v} \times \mathbf{v} = \mathbf{0}$

10. (a)  $\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & -9 \\ -6 & -2 & -1 \end{vmatrix} = -19\mathbf{i} + 56\mathbf{j} + 2\mathbf{k}$

(b)  $\mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = 19\mathbf{i} - 56\mathbf{j} - 2\mathbf{k}$

(c)  $\mathbf{v} \times \mathbf{v} = \mathbf{0}$

13.  $\mathbf{u} = \mathbf{i} + \mathbf{j} + \mathbf{k}$ ,  $\mathbf{v} = 2\mathbf{i} + \mathbf{j} - \mathbf{k}$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 2 & 1 & -1 \end{vmatrix} = -2\mathbf{i} + 3\mathbf{j} - \mathbf{k} = \langle -2, 3, -1 \rangle$$

$$\mathbf{u} \cdot (\mathbf{u} \times \mathbf{v}) = 1(-2) + 1(3) + 1(-1)$$

$$= 0 \Rightarrow \mathbf{u} \perp \mathbf{u} \times \mathbf{v}$$

$$\mathbf{v} \cdot (\mathbf{u} \times \mathbf{v}) = 2(-2) + 1(3) + (-1)(-1)$$

$$= 0 \Rightarrow \mathbf{v} \perp \mathbf{u} \times \mathbf{v}$$

14.  $\mathbf{u} = \mathbf{i} + 6\mathbf{j}$ ,  $\mathbf{v} = -2\mathbf{i} + \mathbf{j} + \mathbf{k}$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 6 & 0 \\ -2 & 1 & 1 \end{vmatrix} = 6\mathbf{i} - \mathbf{j} + 13\mathbf{k}$$

$$\mathbf{u} \cdot (\mathbf{u} \times \mathbf{v}) = 1(6) + 6(-1) = 0 \Rightarrow \mathbf{u} \perp (\mathbf{u} \times \mathbf{v})$$

$$\mathbf{v} \cdot (\mathbf{u} \times \mathbf{v}) = -2(6) + 1(-1) + 1(13) = 0 \Rightarrow \mathbf{v} \perp (\mathbf{u} \times \mathbf{v})$$

15.  $\mathbf{u} = \langle 4, -3, 1 \rangle$

$$\mathbf{v} = \langle 2, 5, 3 \rangle$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & -3 & 1 \\ 2 & 5 & 3 \end{vmatrix} = -14\mathbf{i} - 10\mathbf{j} + 26\mathbf{k}$$

$$\frac{\mathbf{u} \times \mathbf{v}}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{1}{\sqrt{972}} \langle -14, -10, 26 \rangle$$

$$= \frac{1}{18\sqrt{3}} \langle -14, -10, 26 \rangle$$

$$= \left\langle -\frac{7}{9\sqrt{3}}, -\frac{5}{9\sqrt{3}}, \frac{13}{9\sqrt{3}} \right\rangle$$

16.  $\mathbf{u} = \langle -8, -6, 4 \rangle$

$$\mathbf{v} = \langle 10, -12, -2 \rangle$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -8 & -6 & 4 \\ 10 & -12 & -2 \end{vmatrix} = 60\mathbf{i} + 24\mathbf{j} + 156\mathbf{k}$$

$$\frac{\mathbf{u} \times \mathbf{v}}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{1}{36\sqrt{22}} \langle 60, 24, 156 \rangle$$

$$= \left\langle \frac{5}{3\sqrt{22}}, \frac{2}{3\sqrt{22}}, \frac{13}{3\sqrt{22}} \right\rangle$$

17.  $\mathbf{u} = -3\mathbf{i} + 2\mathbf{j} - 5\mathbf{k}$

$$\mathbf{v} = \mathbf{i} - \mathbf{j} + 4\mathbf{k}$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & 2 & -5 \\ 1 & -1 & -4 \end{vmatrix} = 3\mathbf{i} + 7\mathbf{j} + \mathbf{k}$$

$$\frac{\mathbf{u} \times \mathbf{v}}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{1}{\sqrt{59}} \langle 3, 7, 1 \rangle$$

$$= \left\langle \frac{3}{\sqrt{59}}, \frac{7}{\sqrt{59}}, \frac{1}{\sqrt{59}} \right\rangle$$

18.  $\mathbf{u} = 2\mathbf{k}$

$$\mathbf{v} = 4\mathbf{i} + 6\mathbf{k}$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 2 \\ 4 & 0 & 6 \end{vmatrix} = 8\mathbf{j}$$

$$\frac{\mathbf{u} \times \mathbf{v}}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{1}{8} (8\mathbf{j}) = \mathbf{j} = \langle 0, 1, 0 \rangle$$

19.  $\mathbf{u} = \mathbf{j}$

$$\mathbf{v} = \mathbf{j} + \mathbf{k}$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{vmatrix} = \mathbf{i}$$

$$A = \|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{i}\| = 1$$

20.  $\mathbf{u} = \mathbf{i} + \mathbf{j} + \mathbf{k}$

$$\mathbf{v} = \mathbf{j} + \mathbf{k}$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 0 & 1 & 1 \end{vmatrix} = -\mathbf{j} + \mathbf{k}$$

$$A = \|\mathbf{u} \times \mathbf{v}\| = \|-\mathbf{j} + \mathbf{k}\| = \sqrt{2}$$

21.  $\mathbf{u} = \langle 3, 2, -1 \rangle$

$$\mathbf{v} = \langle 1, 2, 3 \rangle$$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 2 & -1 \\ 1 & 2 & 3 \end{vmatrix} = \langle 8, -10, 4 \rangle$$

$$A = \|\mathbf{u} \times \mathbf{v}\| = \|\langle 8, -10, 4 \rangle\| = \sqrt{180} = 6\sqrt{5}$$

22.  $\mathbf{u} = \langle 2, -1, 0 \rangle$

$\mathbf{v} = \langle -1, 2, 0 \rangle$

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 0 \\ -1 & 2 & 0 \end{vmatrix} = \langle 0, 0, 3 \rangle$$

$A = \|\mathbf{u} \times \mathbf{v}\| = \|\langle 0, 0, 3 \rangle\| = 3$

23.  $A(0, 3, 2), B(1, 5, 5), C(6, 9, 5), D(5, 7, 2)$

$\overline{AB} = \langle 1, 2, 3 \rangle$

$\overline{DC} = \langle 1, 2, 3 \rangle$

$\overline{BC} = \langle 5, 4, 0 \rangle$

$\overline{AD} = \langle 5, 4, 0 \rangle$

Because  $\overline{AB} = \overline{DC}$  and  $\overline{BC} = \overline{AD}$ , the figure  $ABCD$  is a parallelogram.

$\overline{AB}$  and  $\overline{AD}$  are adjacent sides

$$\overline{AB} \times \overline{AD} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 3 \\ 5 & 4 & 0 \end{vmatrix} = \langle -12, 15, -6 \rangle$$

$A = \|\overline{AB} \times \overline{AD}\| = \sqrt{144 + 225 + 36} = 9\sqrt{5}$

24.  $A(2, -3, 1), B(6, 5, -1), C(7, 2, 2), D(3, -6, 4)$

$\overline{AB} = \langle 4, 8, -2 \rangle$

$\overline{DC} = \langle 4, 8, -2 \rangle$

$\overline{BC} = \langle 1, -3, 3 \rangle$

$\overline{AD} = \langle 1, -3, 3 \rangle$

Because  $\overline{AB} = \overline{DC}$  and  $\overline{BC} = \overline{AD}$ , the figure  $ABCD$  is a parallelogram.

$\overline{AB}$  and  $\overline{AD}$  are adjacent sides

$$\overline{AB} \times \overline{AD} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & 8 & -2 \\ 1 & -3 & 3 \end{vmatrix} = \langle 18, -14, -20 \rangle$$

$A = \|\overline{AB} \times \overline{AD}\| = \sqrt{324 + 196 + 400} = 2\sqrt{230}$

25.  $A(0, 0, 0), B(1, 0, 3), C(-3, 2, 0)$

$\overline{AB} = \langle 1, 0, 3 \rangle, \overline{AC} = \langle -3, 2, 0 \rangle$

$$\overline{AB} \times \overline{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 3 \\ -3 & 2 & 0 \end{vmatrix} = \langle -6, -9, 2 \rangle$$

$A = \frac{1}{2} \|\overline{AB} \times \overline{AC}\| = \frac{1}{2} \sqrt{36 + 81 + 4} = \frac{11}{2}$

26.  $A(2, -3, 4), B(0, 1, 2), C(-1, 2, 0)$

$\overline{AB} = \langle -2, 4, -2 \rangle, \overline{AC} = \langle -3, 5, -4 \rangle$

$$\overline{AB} \times \overline{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 4 & -2 \\ -3 & 5 & -4 \end{vmatrix} = -6\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$$


$A = \frac{1}{2} \|\overline{AB} \times \overline{AC}\| = \frac{1}{2} \sqrt{44} = \sqrt{11}$

27.  $\mathbf{F} = -100\mathbf{k}$

$\overline{PQ} = 0.17(\cos 40^\circ \mathbf{j} + \sin 40^\circ \mathbf{k})$

$$\begin{aligned} \overline{PQ} \times \mathbf{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0.17 \cos 40^\circ & 0.17 \sin 40^\circ \\ 0 & 0 & -100 \end{vmatrix} \\ &= -17 \cos 40^\circ \mathbf{i} \end{aligned}$$

$\|\overline{PQ} \times \mathbf{F}\| = 17 \cos 40^\circ \approx 13.02 \text{ N-m}$




28.  $\mathbf{F} = -2000(\cos 30^\circ \mathbf{j} + \sin 30^\circ \mathbf{k}) = -1000\sqrt{3}\mathbf{j} - 1000\mathbf{k}$

$\overline{PQ} = 0.05\mathbf{k}$

$$\begin{aligned} \overline{PQ} \times \mathbf{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 0.05 \\ 0 & -1000\sqrt{3} & -1000 \end{vmatrix} \\ &= 50\sqrt{3}\mathbf{i} \end{aligned}$$

$\|\overline{PQ} \times \mathbf{F}\| = 50\sqrt{3} \text{ N-m}$




29. (a)  $AC = 40 \text{ cm} = 0.4 \text{ m}$

$BC = 32 \text{ cm} = 0.32 \text{ m}$

$\overline{AB} = -0.4\mathbf{j} + 0.32\mathbf{k}$

$\mathbf{F} = -180(\cos \theta \mathbf{j} + \sin \theta \mathbf{k})$




(b) 
$$\begin{aligned}\overline{AB} \times \mathbf{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & -0.4 & 0.32 \\ 0 & -180 \cos \theta & -180 \sin \theta \end{vmatrix} \\ &= (72 \sin \theta + 57.6 \cos \theta)\mathbf{i}\end{aligned}$$

$\|\overline{AB} \times \mathbf{F}\| = |72 \sin \theta + 57.6 \cos \theta|$

(c) When  $\theta = 30^\circ$ ,  $\|\overline{AB} \times \mathbf{F}\| = 72\left(\frac{1}{2}\right) + 57.6\left(\frac{\sqrt{3}}{2}\right) \approx 85.88$

(d) If  $T = |72 \sin \theta + 57.6 \cos \theta|$ ,  $T = 0$  for  $72 \sin \theta = -57.6 \cos \theta \Rightarrow \tan \theta = -\frac{57.6}{72} \Rightarrow \theta \approx 141.34^\circ$ .

For  $0^\circ < \theta < 141.34^\circ$ ,  $T'(\theta) = 72 \cos \theta - 57.6 \sin \theta = 0 \Rightarrow \tan \theta = \frac{72}{57.6} \Rightarrow \theta \approx 51.34^\circ$ .  $\overline{AB}$  and  $\mathbf{F}$  are perpendicular.



From part (d), the zero is  $\theta \approx 141.34^\circ$ , which is when the vectors are parallel.


30. (a) Place the wrench in the  $xy$ -plane, as indicated in the figure.

The angle from  $\overline{AB}$  to  $\mathbf{F}$  is  $30^\circ + 180^\circ + \theta = 210^\circ + \theta$


$\|\overline{OA}\| = 50 \text{ cm} = 0.5 \text{ m}$

$$\overline{OA} = 0.5[\cos(30^\circ)\mathbf{i} + \sin(30^\circ)\mathbf{j}] = \frac{\sqrt{3}}{4}\mathbf{i} + \frac{1}{4}\mathbf{j}$$

$$\mathbf{F} = 56[\cos(210^\circ + \theta)\mathbf{i} + \sin(210^\circ + \theta)\mathbf{j}]$$



$$\begin{aligned}\overline{OA} \times \mathbf{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\sqrt{3}}{4} & \frac{1}{4} & 0 \\ 56 \cos(210^\circ + \theta) & 56 \sin(210^\circ + \theta) & 0 \end{vmatrix} \\ &= [14\sqrt{3} \sin(210^\circ + \theta) - 14 \cos(210^\circ + \theta)]\mathbf{k} \\ &= [14\sqrt{3}(\sin 210^\circ \cos \theta + \cos 210^\circ \sin \theta) - 14(\cos 210^\circ \cos \theta - \sin 210^\circ \sin \theta)]\mathbf{k} \\ &= [14\sqrt{3}\left(-\frac{1}{2} \cos \theta - \frac{\sqrt{3}}{2} \sin \theta\right) - 14\left(-\frac{\sqrt{3}}{2} \cos \theta + \frac{1}{2} \sin \theta\right)]\mathbf{k} = (-28 \sin \theta)\mathbf{k}\end{aligned}$$



$\|\overline{OA} \times \mathbf{F}\| = 28 \sin \theta, 0^\circ \leq \theta \leq 180^\circ$

(b) When  $\theta = 45^\circ$ ,  $\|\overline{OA} \times \mathbf{F}\| = 28 \frac{\sqrt{2}}{2} = 14\sqrt{2} \approx 19.80$

- (c) Let  $T = 28 \sin \theta$

$$\frac{dT}{d\theta} = 28 \cos \theta = 0 \text{ when } \theta = 90^\circ.$$

This is reasonable. When  $\theta = 90^\circ$ , the force is perpendicular to the wrench.

31.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 1$

32.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 1 & 1 & 1 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix} = -1$

33.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 2 & 0 & 1 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{vmatrix} = 6$

34.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 2 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 2 & 2 \end{vmatrix} = 0$

35.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{vmatrix} = 2$

$V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})| = 2$

36.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 1 & 3 & 1 \\ 0 & 6 & 6 \\ -4 & 0 & -4 \end{vmatrix} = -72$

$V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})| = 72$

37.  $\mathbf{u} = \langle 3, 0, 0 \rangle$

$\mathbf{v} = \langle 0, 5, 1 \rangle$

$\mathbf{w} = \langle 2, 0, 5 \rangle$

$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 3 & 0 & 0 \\ 0 & 5 & 1 \\ 2 & 0 & 5 \end{vmatrix} = 75$

$V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})| = 75$

47.  $\mathbf{u} = \langle u_1, u_2, u_3 \rangle, \mathbf{v} = \langle v_1, v_2, v_3 \rangle, \mathbf{w} = \langle w_1, w_2, w_3 \rangle$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v} + \mathbf{w}) &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 + w_1 & v_2 + w_2 & v_3 + w_3 \end{vmatrix} \\ &= [u_2(v_3 + w_3) - u_3(v_2 + w_2)]\mathbf{i} - [u_1(v_3 + w_3) - u_3(v_1 + w_1)]\mathbf{j} + [u_1(v_2 + w_2) - u_2(v_1 + w_1)]\mathbf{k} \\ &= (u_2v_3 - u_3v_2)\mathbf{i} - (u_1v_3 - u_3v_1)\mathbf{j} + (u_1v_2 - u_2v_1)\mathbf{k} + (u_2w_3 - u_3w_2)\mathbf{i} - (u_1w_3 - u_3w_1)\mathbf{j} + (u_1w_2 - u_2w_1)\mathbf{k} \\ &= (\mathbf{u} \times \mathbf{v}) + (\mathbf{u} \times \mathbf{w}) \end{aligned}$$

38.  $\mathbf{u} = \langle 0, 4, 0 \rangle$

$\mathbf{v} = \langle -3, 0, 0 \rangle$

$\mathbf{w} = \langle -1, 1, 5 \rangle$

$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} 0 & 4 & 0 \\ -3 & 0 & 0 \\ -1 & 1 & 5 \end{vmatrix} = -4(-15) = 60$

$V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})| = 60$

$$\begin{aligned} 39. \text{(a)} \quad \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) &= (\mathbf{v} \times \mathbf{w}) \cdot \mathbf{u} \quad \text{(b)} \\ &= \mathbf{w} \cdot (\mathbf{u} \times \mathbf{v}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} \quad \text{(c)} \\ &= \mathbf{v} \cdot (\mathbf{w} \times \mathbf{u}) = (\mathbf{u} \times \mathbf{w}) \cdot \mathbf{v} \quad \text{(d)} \\ &= \mathbf{v} \cdot (\mathbf{w} \times \mathbf{u}) = (\mathbf{w} \times \mathbf{u}) \cdot \mathbf{v} \quad \text{(h)} \\ \text{(e)} \quad \mathbf{u} \cdot (\mathbf{w} \times \mathbf{v}) &= \mathbf{w} \cdot (\mathbf{v} \times \mathbf{u}) \quad \text{(f)} \\ &= \mathbf{w} \cdot (\mathbf{v} \times \mathbf{u}) = (-\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} \quad \text{(g)} \end{aligned}$$

So,  $a = b = c = d = h$  and  $e = f = g$

40.  $\mathbf{u} \times \mathbf{v} = \mathbf{0} \Rightarrow \mathbf{u}$  and  $\mathbf{v}$  are parallel.

$\mathbf{u} \cdot \mathbf{v} = \mathbf{0} \Rightarrow \mathbf{u}$  and  $\mathbf{v}$  are orthogonal.

So,  $\mathbf{u}$  or  $\mathbf{v}$  (or both) is the zero vector.

41. The cross product is orthogonal to the two vectors, so it is orthogonal to the  $yz$ -plane. It lies on the  $x$ -axis, since it is of the form  $\langle k, 0, 0 \rangle$ .

42. Form the vectors for two sides of the triangle, and compute their cross product.

$$\langle x_2 - x_1, y_2 - y_1, z_2 - z_1 \rangle \times \langle x_3 - x_1, y_3 - y_1, z_3 - z_1 \rangle$$

43. False. If the vectors are ordered pairs, then the cross product does not exist.

44. False. The cross product is zero if the given vectors are parallel.

45. False. Let  $\mathbf{u} = \langle 1, 0, 0 \rangle, \mathbf{v} = \langle 1, 0, 0 \rangle, \mathbf{w} = \langle -1, 0, 0 \rangle$ .

Then,  $\mathbf{u} \times \mathbf{v} = \mathbf{u} \times \mathbf{w} = \mathbf{0}$ , but  $\mathbf{v} \neq \mathbf{w}$ .

46. True

48.  $\mathbf{u} = \langle u_1, u_2, u_3 \rangle$ ,  $\mathbf{v} = \langle v_1, v_2, v_3 \rangle$ ,  $c$  is a scalar:

$$\begin{aligned} (c\mathbf{u}) \times \mathbf{v} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ cu_1 & cu_2 & cu_3 \\ v_1 & v_2 & v_3 \end{vmatrix} \\ &= (cu_2v_3 - cu_3v_2)\mathbf{i} - (cu_1v_3 - cu_3v_1)\mathbf{j} + (cu_1v_2 - cu_2v_1)\mathbf{k} \\ &= c[(u_2v_3 - u_3v_2)\mathbf{i} - (u_1v_3 - u_3v_1)\mathbf{j} + (u_1v_2 - u_2v_1)\mathbf{k}] = c(\mathbf{u} \times \mathbf{v}) \end{aligned}$$

49.  $\mathbf{u} = \langle u_1, u_2, u_3 \rangle$

$$\mathbf{u} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ u_1 & u_2 & u_3 \end{vmatrix} = (u_2u_3 - u_3u_2)\mathbf{i} - (u_1u_3 - u_3u_1)\mathbf{j} + (u_1u_2 - u_2u_1)\mathbf{k} = \mathbf{0}$$

50.  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}$

$$\begin{aligned} (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} &= \mathbf{w} \cdot (\mathbf{u} \times \mathbf{v}) = \begin{vmatrix} w_1 & w_2 & w_3 \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix} \\ &= w_1(u_2v_3 - v_2u_3) - w_2(u_1v_3 - v_1u_3) + w_3(u_1v_2 - v_1u_2) \\ &= u_1(v_2w_3 - w_2v_3) - u_2(v_1w_3 - w_1v_3) + u_3(v_1w_2 - w_1v_2) = \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \end{aligned}$$

51.  $\mathbf{u} \times \mathbf{v} = (u_2v_3 - u_3v_2)\mathbf{i} - (u_1v_3 - u_3v_1)\mathbf{j} + (u_1v_2 - u_2v_1)\mathbf{k}$

$$(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{u} = (u_2v_3 - u_3v_2)u_1 + (u_3v_1 - u_1v_3)u_2 + (u_1v_2 - u_2v_1)u_3 = \mathbf{0}$$

$$(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{v} = (u_2v_3 - u_3v_2)v_1 + (u_3v_1 - u_1v_3)v_2 + (u_1v_2 - u_2v_1)v_3 = \mathbf{0}$$

So,  $\mathbf{u} \times \mathbf{v} \perp \mathbf{u}$  and  $\mathbf{u} \times \mathbf{v} \perp \mathbf{v}$ .

52. If  $\mathbf{u}$  and  $\mathbf{v}$  are scalar multiples of each other,  $\mathbf{u} = cv$  for some scalar  $c$ .

$$\mathbf{u} \times \mathbf{v} = (cv) \times \mathbf{v} = c(\mathbf{v} \times \mathbf{v}) = c(\mathbf{0}) = \mathbf{0}$$

If  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ , then  $\|\mathbf{u}\|\|\mathbf{v}\| \sin \theta = 0$ . (Assume  $\mathbf{u} \neq \mathbf{0}, \mathbf{v} \neq \mathbf{0}$ .) So,  $\sin \theta = 0, \theta = 0$ , and  $\mathbf{u}$  and  $\mathbf{v}$  are parallel. So,

$\mathbf{u} = cv$  for some scalar  $c$ .

53.  $\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\|\|\mathbf{v}\| \sin \theta$

If  $\mathbf{u}$  and  $\mathbf{v}$  are orthogonal,  $\theta = \pi/2$  and  $\sin \theta = 1$ . So,  $\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\|\|\mathbf{v}\|$ .

54.  $\mathbf{u} = \langle a_1, b_1, c_1 \rangle$ ,  $\mathbf{v} = \langle a_2, b_2, c_2 \rangle$ ,  $\mathbf{w} = \langle a_3, b_3, c_3 \rangle$

$$\begin{aligned} \mathbf{v} \times \mathbf{w} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = (b_2c_3 - b_3c_2)\mathbf{i} - (a_2c_3 - a_3c_2)\mathbf{j} + (a_2b_3 - a_3b_2)\mathbf{k} \\ \mathbf{u} \times (\mathbf{v} \times \mathbf{w}) &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & b_1 & c_1 \\ (b_2c_3 - b_3c_2) & (a_3c_2 - a_2c_3) & (a_2b_3 - a_3b_2) \end{vmatrix} \\ \mathbf{u} \times (\mathbf{v} \times \mathbf{w}) &= [b_1(a_2b_3 - a_3b_2) - c_1(a_3c_2 - a_2c_3)]\mathbf{i} - [a_1(a_2b_3 - a_3b_2) - c_1(b_2c_3 - b_3c_2)]\mathbf{j} \\ &\quad + [a_1(a_3c_2 - a_2c_3) - b_1(b_2c_3 - b_3c_2)]\mathbf{k} \\ &= [a_2(a_1a_3 + b_1b_3 + c_1c_3) - a_3(a_1a_2 + b_1b_2 + c_1c_2)]\mathbf{i} + [b_2(a_1a_3 + b_1b_3 + c_1c_3) - b_3(a_1a_2 + b_1b_2 + c_1c_2)]\mathbf{j} \\ &\quad + [c_2(a_1a_3 + b_1b_3 + c_1c_3) - c_3(a_1a_2 + b_1b_2 + c_1c_2)]\mathbf{k} \\ &= (a_1a_3 + b_1b_3 + c_1c_3)\langle a_2, b_2, c_2 \rangle - (a_1a_2 + b_1b_2 + c_1c_2)\langle a_3, b_3, c_3 \rangle = (\mathbf{u} \cdot \mathbf{w})\mathbf{v} - (\mathbf{u} \cdot \mathbf{v})\mathbf{w} \end{aligned}$$

55.  $\mathbf{u} = \mathbf{u}_1\mathbf{i} + \mathbf{u}_2\mathbf{j} + \mathbf{u}_3\mathbf{k}$ ,  $\mathbf{v} = \mathbf{v}_1\mathbf{i} + \mathbf{v}_2\mathbf{j} + \mathbf{v}_3\mathbf{k}$ ,  $\mathbf{w} = \mathbf{w}_1\mathbf{i} + \mathbf{w}_2\mathbf{j} + \mathbf{w}_3\mathbf{k}$

$$\begin{aligned}\mathbf{v} \times \mathbf{w} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix} = \langle v_2 w_3 - w_2 v_3, -(v_1 w_3 - w_1 v_3), v_1 w_2 - w_1 v_2 \rangle \\ \mathbf{u} \times (\mathbf{v} \times \mathbf{w}) &= \langle u_1, u_2, u_3 \rangle \cdot \langle v_2 w_3 - w_2 v_3, -(v_1 w_3 - w_1 v_3), v_1 w_2 - w_1 v_2 \rangle \\ &= u_1 v_2 w_3 - u_1 v_3 w_2 - u_2 v_1 w_3 + u_2 v_3 w_1 + u_3 v_1 w_2 - u_3 v_2 w_1 \\ &= \begin{vmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ w_1 & w_2 & w_3 \end{vmatrix}\end{aligned}$$

## Section 11.5 Lines and Planes in Space

1. The parametric equations of a line L parallel to

$\mathbf{v} = \langle a, b, c \rangle$  and passing through the point  $P(x_1, y_1, z_1)$

are  $x = x_1 + at$ ,  $y = y_1 + bt$ ,  $z = z_1 + ct$ .

The symmetric equations are

$$\frac{x - x_1}{a} = \frac{y - y_1}{b} = \frac{z - z_1}{c}.$$

2. In the equation of the plane

$$2(x - 1) + 4(y - 3) - (z + 5) = 0, \quad a = 2, b = 4, \text{ and}$$

$c = -1$ . Therefore, the normal vector is  $\langle 2, 4, -1 \rangle$ .

3. Answers will vary. Any plane that has a missing  $x$ -variable in its equation is parallel to the  $x$ -axis.

Sample answer:  $3y - z = 5$

4. First choose a point  $Q$  in one plane. Then use Theorem 11.13:

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|}$$

where  $P$  is a point in the other plane and  $\mathbf{n}$  is normal to that plane.

5.  $x = -2 + t, y = 3t, z = 4 + t$

- (a)  $(0, 6, 6)$ : For  $x = 0 = -2 + t$ , you have  $t = 2$ .

Then  $y = 3(2) = 6$  and  $z = 4 + 2 = 6$ . Yes,

$(0, 6, 6)$  lies on the line.

- (b)  $(2, 3, 5)$ : For  $x = 2 = -2 + t$ , you have  $t = 4$ .

Then  $y = 3(4) = 12 \neq 3$ . No,  $(2, 3, 5)$  does not lie on the line.

- (c)  $(-4, -6, 2)$ : For  $x = -4 = -2 + t$ , you have

$$t = -2.$$

Then  $y = 3(-2) = -6$  and  $z = 4 - 2 = 2$ . Yes,

$(-4, -6, 2)$  lies on the line.

6.  $\frac{x - 3}{2} = \frac{y - 7}{8} = z + 2$

(a)  $(7, 23, 0)$ : Substituting, you have

$$\begin{aligned}\frac{7 - 3}{2} &= \frac{23 - 7}{8} = 0 + 2 \\ 2 &= 2 = 2\end{aligned}$$

Yes,  $(7, 23, 0)$  lies on the line.

(b)  $(1, -1, -3)$ : Substituting, you have

$$\begin{aligned}\frac{1 - 3}{2} &= \frac{-1 - 7}{8} = -3 + 2 \\ -1 &= -1 = -1\end{aligned}$$

Yes,  $(1, -1, -3)$  lies on the line.

(c)  $(-7, 47, -7)$ : Substituting, you have

$$\begin{aligned}\frac{-7 - 3}{2} &= \frac{47 - 7}{8} = -7 + 2 \\ -5 &\neq 5 \neq -5\end{aligned}$$

No,  $(-7, 47, -7)$  does not lie on the line.

7. Point:  $(0, 0, 0)$

Direction vector:  $\langle 3, 1, 5 \rangle$

Direction numbers: 3, 1, 5

(a) Parametric:  $x = 3t, y = t, z = 5t$

(b) Symmetric:  $\frac{x}{3} = \frac{y}{1} = \frac{z}{5}$

8. Point:  $(0, 0, 0)$ Direction vector:  $\mathbf{v} = \left\langle -2, \frac{5}{2}, 1 \right\rangle$ Direction numbers:  $-4, 5, 2$ (a) Parametric:  $x = -4t, y = 5t, z = 2t$ (b) Symmetric:  $\frac{x}{-4} = \frac{y}{5} = \frac{z}{2}$ 9. Point:  $(-2, 0, 3)$ Direction vector:  $\mathbf{v} = \langle 2, 4, -2 \rangle$ Direction numbers:  $2, 4, -2$ (a) Parametric:  $x = -2 + 2t, y = 4t, z = 3 - 2t$ (b) Symmetric:  $\frac{x + 2}{2} = \frac{y}{4} = \frac{z - 3}{-2}$ 10. Point:  $(-3, 0, 2)$ Direction vector:  $\mathbf{v} = \langle 0, 6, 3 \rangle$ Direction numbers:  $0, 2, 1$ (a) Parametric:  $x = -3, y = 2t, z = 2 + t$ (b) Symmetric:  $\frac{y}{2} = z - 2, x = -3$ 11. Point:  $(1, 0, 1)$ Direction vector:  $\mathbf{v} = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$ Direction numbers:  $3, -2, 1$ (a) Parametric:  $x = 1 + 3t, y = -2t, z = 1 + t$ (b) Symmetric:  $\frac{x - 1}{3} = \frac{y}{-2} = \frac{z - 1}{1}$ 12. Point:  $(-3, 5, 4)$ Direction numbers:  $3, -2, 1$ (a) Parametric:  $x = -3 + 3t, y = 5 - 2t, z = 4 + t$ (b) Symmetric:  $\frac{x + 3}{3} = \frac{y - 5}{-2} = z - 4$ 13. Points:  $(5, -3, -2), \left( -\frac{2}{3}, \frac{2}{3}, 1 \right)$ Direction vector:  $\mathbf{v} = \frac{17}{3}\mathbf{i} - \frac{11}{3}\mathbf{j} - 3\mathbf{k}$ Direction numbers:  $17, -11, -9$ 

(a) Parametric:

$$x = 5 + 17t, y = -3 - 11t, z = -2 - 9t$$

(b) Symmetric:  $\frac{x - 5}{17} = \frac{y + 3}{-11} = \frac{z + 2}{-9}$ 14. Points:  $(0, 4, 3), (-1, 2, 5)$ Direction vector:  $\langle 1, 2, -2 \rangle$ Direction numbers:  $1, 2, -2$ (a) Parametric:  $x = t, y = 4 + 2t, z = 3 - 2t$ (b) Symmetric:  $x = \frac{y - 4}{2} = \frac{z - 3}{-2}$ 15. Points:  $(7, -2, 6), (-3, 0, 6)$ Direction vector:  $\langle -10, 2, 0 \rangle$ Direction numbers:  $-10, 2, 0$ (a) Parametric:  $x = 7 - 10t, y = -2 + 2t, z = 6$ (b) Symmetric: Not possible because the direction number for  $z$  is 0. But, you could describe the line as  $\frac{x - 7}{10} = \frac{y + 2}{-2}, z = 6$ .16. Points:  $(0, 0, 25), (10, 10, 0)$ Direction vector:  $\langle 10, 10, -25 \rangle$ Direction numbers:  $2, 2, -5$ (a) Parametric:  $x = 2t, y = 2t, z = 25 - 5t$ (b) Symmetric:  $\frac{x}{2} = \frac{y}{2} = \frac{z - 25}{-5}$ 17. Point:  $(2, 3, 4)$ Direction vector:  $\mathbf{v} = \mathbf{k}$ Direction numbers:  $0, 0, 1$ Parametric:  $x = 2, y = 3, z = 4 + t$ 18. Point:  $(-4, 5, 2)$ Direction vector:  $\mathbf{v} = \mathbf{j}$ Direction numbers:  $0, 1, 0$ Parametric:  $x = -4, y = 5 + t, z = 2$ 19. Point:  $(2, 3, 4)$ Direction vector:  $\mathbf{v} = 3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ Direction numbers:  $3, 2, -1$ Parametric:  $x = 2 + 3t, y = 3 + 2t, z = 4 - t$ 20. Point  $(-4, 5, 2)$ Direction vector:  $\mathbf{v} = -\mathbf{i} + 2\mathbf{j} + \mathbf{k}$ Direction numbers:  $-1, 2, 1$ Parametric:  $x = -4 - t, y = 5 + 2t, z = 2 + t$

21. Point:  $(5, -3, -4)$

Direction vector:  $\mathbf{v} = \langle 2, -1, 3 \rangle$

Direction numbers:  $2, -1, 3$

Parametric:  $x = 5 + 2t, y = -3 - t, z = -4 + 3t$

22. Point:  $(-1, 4, -3)$

Direction vector:  $\mathbf{v} = 5\mathbf{i} - \mathbf{j}$

Direction numbers:  $5, -1, 0$

Parametric:  $x = -1 + 5t, y = 4 - t, z = -3$

23. Point:  $(2, 1, 2)$

Direction vector:  $\langle -1, 1, 1 \rangle$

Direction numbers:  $-1, 1, 1$

Parametric:  $x = 2 - t, y = 1 + t, z = 2 + t$

24. Point:  $(-6, 0, 8)$

Direction vector:  $\langle -2, 2, 0 \rangle$

Direction numbers:  $-2, 2, 0$

Parametric:  $x = -6 - 2t, y = 2t, z = 8$

25. Let  $t = 0$ :  $P = (3, -1, -2)$  (other answers possible)

$\mathbf{v} = \langle -1, 2, 0 \rangle$  (any nonzero multiple of  $\mathbf{v}$  is correct)

26. Let  $t = 0$ :  $P = (0, 5, 4)$  (other answers possible)

$\mathbf{v} = \langle 4, -1, 3 \rangle$  (any nonzero multiple of  $\mathbf{v}$  is correct)

27. Let each quantity equal 0:

$P = (7, -6, -2)$  (other answers possible)

$\mathbf{v} = \langle 4, 2, 1 \rangle$  (any nonzero multiple of  $\mathbf{v}$  is correct)

28. Let each quantity equal 0:

$P = (-3, 0, 3)$  (other answers possible)

$\mathbf{v} = \langle 5, 8, 6 \rangle$  (any nonzero multiple of  $\mathbf{v}$  is correct)

29.  $L_1$ :  $\mathbf{v}_1 = \langle -3, 2, 4 \rangle$  and  $P = (6, -2, 5)$  on  $L_1$

$L_2$ :  $\mathbf{v}_2 = \langle 6, -4, -8 \rangle$  and  $P = (6, -2, 5)$  on  $L_2$

The lines are identical.

30.  $L_1$ :  $\mathbf{v}_1 = \langle 2, -1, 3 \rangle$  and  $P = (1, -1, 0)$  on  $L_1$

$L_2$ :  $\mathbf{v}_2 = \langle 2, -1, 3 \rangle$  and  $P$  not on  $L_1$

The lines are parallel.

31.  $L_1$ :  $\mathbf{v}_1 = \langle 4, -2, 3 \rangle$  and  $P = (8, -5, -9)$  on  $L_1$

$L_2$ :  $\mathbf{v}_2 = \langle -8, 4, -6 \rangle$  and  $P = (8, -5, -9)$  on  $L_2$

The lines are identical.

32.  $L_1$ :  $\mathbf{v}_1 = \langle 4, 2, 4 \rangle$  and  $P = (1, 1, -3)$  on  $L_1$

$L_2$ :  $\mathbf{v}_2 = \langle 1, 0.5, 1 \rangle$  and  $P$  not on  $L_2$

The lines are parallel.

33. At the point of intersection, the coordinates for one line equal the corresponding coordinates for the other line. So,

(i)  $4t + 2 = 2s + 2$ , (ii)  $3 = 2s + 3$ , and

(iii)  $-t + 1 = s + 1$ .

From (ii), you find that  $s = 0$  and consequently, from (iii),  $t = 0$ . Letting  $s = t = 0$ , you see that equation (i) is satisfied and so the two lines intersect. Substituting zero for  $s$  or for  $t$ , you obtain the point  $(2, 3, 1)$ .

$$\mathbf{u} = 4\mathbf{i} - \mathbf{k} \quad (\text{First line})$$

$$\mathbf{v} = 2\mathbf{i} + 2\mathbf{j} + \mathbf{k} \quad (\text{Second line})$$

$$\cos \theta = \frac{|\mathbf{u} \cdot \mathbf{v}|}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{8 - 1}{\sqrt{17} \sqrt{9}} = \frac{7}{3\sqrt{17}} = \frac{7\sqrt{17}}{51}$$

$$\theta \approx 55.5^\circ$$

34. By equating like variables, you have

(i)  $-3t + 1 = 3s + 1$ , (ii)  $4t + 1 = 2s + 4$ , and

(iii)  $2t + 4 = -s + 1$ .

From (i) you have  $s = -t$ , and consequently from (ii),  $t = \frac{1}{2}$  and from (iii),  $t = -3$ . The lines do not intersect.

35. Writing the equations of the lines in parametric form you have

$$x = 3t \qquad y = 2 - t \qquad z = -1 + t$$

$$x = 1 + 4s \qquad y = -2 + s \qquad z = -3 - 3s.$$

For the coordinates to be equal,  $3t = 1 + 4s$  and  $2 - t = -2 + s$ . Solving this system yields  $t = \frac{17}{7}$  and  $s = \frac{11}{7}$ . When using these values for  $s$  and  $t$ , the  $z$  coordinates are not equal. The lines do not intersect.

36. Writing the equations of the lines in parametric form you have

$$\begin{array}{lll} x = 2 - 3t & y = 2 + 6t & z = 3 + t \\ x = 3 + 2s & y = -5 + s & z = -2 + 4s \end{array}$$

By equating like variables, you have  $2 - 3t = 3 + 2s$ ,  $2 + 6t = -5 + s$ ,  $3 + t = -2 + 4s$ . So,  $t = -1$ ,  $s = 1$  and the point of intersection is  $(5, -4, 2)$ .

$$\begin{aligned} \mathbf{u} &= \langle -3, 6, 1 \rangle \quad (\text{First line}) \\ \mathbf{v} &= \langle 2, 1, 4 \rangle \quad (\text{Second line}) \end{aligned}$$

$$\begin{aligned} \cos \theta &= \frac{|\mathbf{u} \cdot \mathbf{v}|}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{4}{\sqrt{46} \sqrt{21}} = \frac{4}{\sqrt{966}} = \frac{2\sqrt{966}}{483} \\ \theta &\approx 82.6^\circ \end{aligned}$$

37.  $x + 2y - 4z - 1 = 0$

(a)  $(-7, 2, -1)$ :  $(-7) + 2(2) - 4(-1) - 1 = 0$

Point is in plane.

(b)  $(5, 2, 2)$ :  $5 + 2(2) - 4(2) - 1 = 0$

Point is in plane.

(c)  $(-6, 1, -1)$ :  $-6 + 2(1) - 4(-1) - 1 = -1 \neq 0$

Point is not in plane.

38.  $2x + y + 3z - 6 = 0$

(a)  $(3, 6, -2)$ :  $2(3) + 6 + 3(-2) - 6 = 0$

Point is in plane.

(b)  $(-1, 5, -1)$ :  $2(-1) + 5 + 3(-1) - 6 = -6 \neq 0$

Point is not in plane.

(c)  $(2, 1, 0)$ :  $2(2) + 1 + 3(0) - 6 = -1 \neq 0$

Point is not in plane.

39. Point:  $(1, 3, -7)$

Normal vector:  $\mathbf{n} = \mathbf{j} = \langle 0, 1, 0 \rangle$

$$0(x - 1) + 1(y - 3) + 0(z - (-7)) = 0$$

$$y - 3 = 0$$

40. Point:  $(0, -1, 4)$

Normal vector:  $\mathbf{n} = \mathbf{k} = \langle 0, 0, 1 \rangle$

$$0(x - 0) + 0(y + 1) + 1(z - 4) = 0$$

$$z - 4 = 0$$

41. Point:  $(3, 2, 2)$

Normal vector:  $\mathbf{n} = 2\mathbf{i} + 3\mathbf{j} - \mathbf{k}$

$$2(x - 3) + 3(y - 2) - 1(z - 2) = 0$$

$$2x + 3y - z - 10 = 0$$

42. Point:  $(0, 0, 0)$

Normal vector:  $\mathbf{n} = -3\mathbf{i} + 2\mathbf{k}$

$$-3(x - 0) + 0(y - 0) + 2(z - 0) = 0$$

$$-3x + 2z = 0$$

43. Point:  $(-1, 4, 0)$

Normal vector:  $\mathbf{v} = \langle 2, -1, -2 \rangle$

$$2(x + 1) - 1(y - 4) - 2(z - 0) = 0$$

$$2x - y - 2z + 6 = 0$$

44. Point:  $(3, 2, 2)$

Normal vector:  $\mathbf{v} = 4\mathbf{i} + \mathbf{j} - 3\mathbf{k}$

$$4(x - 3) + (y - 2) - 3(z - 2) = 0$$

$$4x + y - 3z - 8 = 0$$

45. Let  $\mathbf{u}$  be the vector from  $(0, 0, 0)$  to

$$(2, 0, 3): \mathbf{u} = \langle 2, 0, 3 \rangle$$

Let  $\mathbf{u}$  be the vector from  $(0, 0, 0)$  to

$$(-3, -1, 5): \mathbf{v} = \langle -3, -1, 5 \rangle$$

$$\text{Normal vectors: } \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 0 & 3 \\ -3 & -1 & 5 \end{vmatrix} = \langle 3, -19, -2 \rangle$$

$$3(x - 0) - 19(y - 0) - 2(z - 0) = 0$$

$$3x - 19y - 2z = 0$$

46. Let  $\mathbf{u}$  be the vector from  $(3, -1, 2)$  to  $(2, 1, 5)$ :

$$\mathbf{u} = \langle -1, 2, 3 \rangle$$

Let  $\mathbf{u}$  be the vector from  $(3, -1, 2)$  to  $(1, -2, -2)$ :

$$\mathbf{v} = \langle -2, -1, -4 \rangle$$

Normal vector:

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 2 & 3 \\ -2 & -1 & -4 \end{vmatrix} = \langle -5, -10, 5 \rangle = -5\langle 1, 2, -1 \rangle$$

$$1(x - 3) + 2(y + 1) - (z - 2) = 0$$

$$x + 2y - z + 1 = 0$$

47. Let  $\mathbf{u}$  be the vector from  $(1, 2, 3)$  to

$$(3, 2, 1): \mathbf{u} = 2\mathbf{i} - 2\mathbf{k}$$

Let  $\mathbf{v}$  be the vector from  $(1, 2, 3)$  to

$$(-1, -2, 2): \mathbf{v} = -2\mathbf{i} - 4\mathbf{j} - \mathbf{k}$$

Normal vector:

$$\left(\frac{1}{2}\mathbf{u}\right) \times (-\mathbf{v}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & -1 \\ 2 & 4 & 1 \end{vmatrix} = 4\mathbf{i} - 3\mathbf{j} + 4\mathbf{k}$$

$$4(x - 1) - 3(y - 2) + 4(z - 3) = 0$$

$$4x - 3y + 4z - 10 = 0$$

48.  $(1, 2, 3)$ , Normal vector:

$$\mathbf{v} = \mathbf{i}, l(x - 1) = 0, x - 1 = 0$$

49.  $(1, 2, 3)$ , Normal vector:

$$\mathbf{v} = \mathbf{k}, l(z - 3) = 0, z - 3 = 0$$

50. The plane passes through the three points

$$(0, 0, 0), (0, 1, 0), (\sqrt{3}, 0, 1).$$

The vector from  $(0, 0, 0)$  to  $(0, 1, 0)$ :  $\mathbf{u} = \mathbf{j}$

The vector from  $(0, 0, 0)$  to  $(\sqrt{3}, 0, 1)$ :  $\mathbf{v} = \sqrt{3}\mathbf{i} + \mathbf{k}$

$$\text{Normal vector: } \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 0 \\ \sqrt{3} & 0 & 1 \end{vmatrix} = \mathbf{i} - \sqrt{3}\mathbf{k}$$

$$x - \sqrt{3}z = 0$$

51. The direction vectors for the lines are  $\mathbf{u} = -2\mathbf{i} + \mathbf{j} + \mathbf{k}$ ,

$$\mathbf{v} = -3\mathbf{i} + 4\mathbf{j} - \mathbf{k}.$$

$$\text{Normal vector: } \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 1 & 1 \\ -3 & 4 & -1 \end{vmatrix} = -5(\mathbf{i} + \mathbf{j} + \mathbf{k})$$

Point of intersection of the lines:  $(-1, 5, 1)$

$$(x + 1) + (y - 5) + (z - 1) = 0$$

$$x + y + z - 5 = 0$$

52. The direction of the line is  $\mathbf{u} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$ . Choose any point on the line,  $[(0, 4, 0)$ , for example], and let  $\mathbf{v}$  be the vector from  $(0, 4, 0)$  to the given point  $(2, 2, 1)$ :

$$\mathbf{v} = 2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$$

$$\text{Normal vector: } \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 1 \\ 2 & -2 & 1 \end{vmatrix} = \mathbf{i} - 2\mathbf{k}$$

$$(x - 2) - 2(z - 1) = 0$$

$$x - 2z = 0$$

53. Let  $\mathbf{v}$  be the vector from  $(-1, 1, -1)$  to  $(2, 2, 1)$ :

$$\mathbf{v} = 3\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

Let  $\mathbf{n}$  be a vector normal to the plane  
 $2x - 3y + z = 3$ :  $\mathbf{n} = 2\mathbf{i} - 3\mathbf{j} + \mathbf{k}$

Because  $\mathbf{v}$  and  $\mathbf{n}$  both lie in the plane  $P$ , the normal vector to  $P$  is

$$\mathbf{v} \times \mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 1 & 2 \\ 2 & -3 & 1 \end{vmatrix} = 7\mathbf{i} - \mathbf{j} - 11\mathbf{k}$$

$$7(x - 2) + 1(y - 2) - 11(z - 1) = 0$$

$$7x + y - 11z - 5 = 0$$

54. Let  $\mathbf{v}$  be the vector from  $(3, 2, 1)$  to  $(3, 1, -5)$ :

$$\mathbf{v} = -\mathbf{j} - 6\mathbf{k}$$

Let  $\mathbf{n}$  be the normal to the given plane:  
 $\mathbf{n} = 6\mathbf{i} + 7\mathbf{j} + 2\mathbf{k}$

Because  $\mathbf{v}$  and  $\mathbf{n}$  both lie in the plane  $P$ , the normal vector to  $P$  is:

$$\mathbf{v} \times \mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & -1 & -6 \\ 6 & 7 & 2 \end{vmatrix} = 40\mathbf{i} - 36\mathbf{j} + 6\mathbf{k}$$

$$= 2(20\mathbf{i} - 18\mathbf{j} + 3\mathbf{k})$$

$$20(x - 3) - 18(y - 2) + 3(z - 1) = 0$$

$$20x - 18y + 3z - 27 = 0$$

55. Let  $\mathbf{u} = \mathbf{i}$  and let  $\mathbf{v}$  be the vector from  $(1, -2, -1)$  to  $(2, 5, 6)$ :  $\mathbf{v} = \mathbf{i} + 7\mathbf{j} + 7\mathbf{k}$

Because  $\mathbf{u}$  and  $\mathbf{v}$  both lie in the plane  $P$ , the normal vector to  $P$  is:

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 0 \\ 1 & 7 & 7 \end{vmatrix} = -7\mathbf{j} + 7\mathbf{k} = -7(\mathbf{j} - \mathbf{k})$$

$$[y - (-2)] - [z - (-1)] = 0$$

$$y - z + 1 = 0$$

56. Let  $\mathbf{u} = \mathbf{k}$  and let  $\mathbf{v}$  be the vector from  $(4, 2, 1)$  to  $(-3, 5, 7)$ :  $\mathbf{v} = -7\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}$

Because  $\mathbf{u}$  and  $\mathbf{v}$  both lie in the plane  $P$ , the normal vector to  $P$  is:

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 1 \\ -7 & 3 & 6 \end{vmatrix} = -3\mathbf{i} - 7\mathbf{j} = -(3\mathbf{i} + 7\mathbf{j})$$

$$3(x - 4) + 7(y - 2) = 0$$

$$3x + 7y - 26 = 0$$

57. Let  $(x, y, z)$  be equidistant from  $(2, 2, 0)$  and  $(0, 2, 2)$ .

$$\sqrt{(x - 2)^2 + (y - 2)^2 + (z - 0)^2} = \sqrt{(x - 0)^2 + (y - 2)^2 + (z - 2)^2}$$

$$x^2 - 4x + 4 + y^2 - 4y + 4 + z^2 = x^2 + y^2 - 4y + 4 + z^2 - 4z + 4$$

$$-4x + 8 = -4z + 8$$

$$x - z = 0 \text{ Plane}$$

58. Let  $(x, y, z)$  be equidistant from  $(1, 0, 2)$  and  $(2, 0, 1)$ .

$$\sqrt{(x - 1)^2 + (y - 0)^2 + (z - 2)^2} = \sqrt{(x - 2)^2 + (y - 0)^2 + (z - 1)^2}$$

$$x^2 - 2x + 1 + y^2 + z^2 - 4z + 4 = x^2 - 4x + 4 + y^2 + z^2 - 2z + 1$$

$$-2x - 4z + 5 = -4x - 2z + 5$$

$$2x - 2z = 0$$

$$x - z = 0 \text{ Plane}$$

59. Let  $(x, y, z)$  be equidistant from  $(-3, 1, 2)$  and  $(6, -2, 4)$ .

$$\sqrt{(x + 3)^2 + (y - 1)^2 + (z - 2)^2} = \sqrt{(x - 6)^2 + (y + 2)^2 + (z - 4)^2}$$

$$x^2 + 6x + 9 + y^2 - 2y + 1 + z^2 - 4z + 4 = x^2 - 12x + 36 + y^2 + 4y + 4 + z^2 - 8z + 16$$

$$6x - 2y - 4z + 14 = -12x + 4y - 8z + 56$$

$$18x - 6y + 4z - 42 = 0$$

$$9x - 3y + 2z - 21 = 0 \text{ Plane}$$

60. Let  $(x, y, z)$  be equidistant from  $(-5, 1, -3)$  and  $(2, -1, 6)$ .

$$\sqrt{(x + 5)^2 + (y - 1)^2 + (z + 3)^2} = \sqrt{(x - 2)^2 + (y + 1)^2 + (z - 6)^2}$$

$$x^2 + 10x + 25 + y^2 - 2y + 1 + z^2 + 6z + 9 = x^2 - 4x + 4 + y^2 + 2y + 1 + z^2 - 12z + 36$$

$$10x - 2y + 6z + 35 = -4x + 2y - 12z + 41$$

$$14x - 4y + 18z - 6 = 0$$

$$7x - 2y + 9z - 3 = 0 \text{ Plane}$$

61. First plane:  $\mathbf{n}_1 = \langle -5, 2, -8 \rangle$  and  $P = (0, 3, 0)$  on plane

Second plane:  $\mathbf{n}_2 = \langle 15, -6, 24 \rangle = -3\mathbf{n}_1$  and  $P$  not on plane

Parallel planes

(Note: The equations are not equivalent.)

62. First plane:  $\mathbf{n}_1 = \langle 2, -1, 3 \rangle$  and  $P = (4, 0, 0)$  on plane

Second plane:  $\mathbf{n}_2 = \langle 8, -4, 12 \rangle = 4\mathbf{n}_1$  and  $P$  not on plane.

Parallel planes

(Note: The equations are not equivalent.)

63. First plane:  $\mathbf{n}_1 = \langle 3, -2, 5 \rangle$  and  $P = (0, 0, 2)$  on plane  
 Second plane:  $\mathbf{n}_2 = \langle 75, -50, 125 \rangle = 25\mathbf{n}_1$  and  $P$  on plane

Planes are identical.

(Note: The equations are equivalent.)

64. First plane:  $\mathbf{n}_1 = \langle -1, 4, -1 \rangle$  and  $P = (-6, 0, 0)$  on plane

Second plane:  $\mathbf{n}_2 = \left\langle -\frac{5}{2}, 10, -\frac{5}{2} \right\rangle = \frac{5}{2}\mathbf{n}_1$  and  $P$  on plane

Planes are identical.

(Note: The equations are equivalent.)

65. (a)  $\mathbf{n}_1 = 3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$  and  $\mathbf{n}_2 = \mathbf{i} - 4\mathbf{j} + 2\mathbf{k}$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{|-7|}{\sqrt{14} \sqrt{21}} = \frac{\sqrt{6}}{6}$$

$$\Rightarrow \theta \approx 65.91^\circ$$

(b) The direction vector for the line is

$$\mathbf{n}_2 \times \mathbf{n}_1 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -4 & 2 \\ 3 & 2 & -1 \end{vmatrix} = 7(\mathbf{j} + 2\mathbf{k}).$$

Find a point of intersection of the planes.

$$\begin{aligned} 6x + 4y - 2z &= 14 \\ x - 4y + 2z &= 0 \\ \hline 7x &= 14 \\ x &= 2 \end{aligned}$$

Substituting 2 for  $x$  in the second equation, you have  $-4y + 2z = -2$  or  $z = 2y - 1$ .

Letting  $y = 1$ , a point of intersection is  $(2, 1, 1)$ .

$$x = 2, y = 1 + t, z = 1 + 2t$$

66. (a)  $\mathbf{n}_1 = \langle -2, 1, 1 \rangle$  and  $\mathbf{n}_2 = \langle 6, -3, 2 \rangle$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{|-13|}{\sqrt{6} \sqrt{49}} = \frac{13\sqrt{6}}{42}$$

$$\Rightarrow \theta \approx 40.70^\circ$$

(b) The direction vector for the line is

$$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 1 & 1 \\ 6 & -3 & 2 \end{vmatrix} = 5(\mathbf{i} + 2\mathbf{j}).$$

Find a point of intersection of the planes.

$$\begin{aligned} -6x + 3y + 3z &= 6 \\ 6x - 3y + 2z &= 4 \\ \hline 5z &= 10 \\ z &= 2 \end{aligned}$$

Substituting 2 for  $z$  in the first equation, you have  $-2x + y = 0$  or  $y = 2x$ . Letting  $x = 0$ , a point of intersection is  $(0, 0, 2)$ .

$$x = 5t, y = 10t, z = 2 \text{ or } x = t, y = 2t, z = 2$$

67. (a)  $\mathbf{n}_1 = \langle 3, -1, 1 \rangle$  and  $\mathbf{n}_2 = \langle 4, 6, 3 \rangle$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{|9|}{\sqrt{11} \sqrt{61}} = \frac{9\sqrt{671}}{671}$$

$$\Rightarrow \theta \approx 69.67^\circ$$

(b) The direction vector for the line is

$$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -1 & 1 \\ 4 & 6 & 3 \end{vmatrix} = -9\mathbf{i} - 5\mathbf{j} + 22\mathbf{k}.$$

Find a point of intersection of the planes.

$$\begin{aligned} 18x - 6y + 6z &= 42 \\ 4x + 6y + 3z &= 2 \\ \hline 22x + 9z &= 44 \end{aligned}$$

Let  $z = 0, 22x = 44 \Rightarrow x = 2$  and

$$3(2) - y + 0 = 7 \Rightarrow y = -1.$$

A point of intersection is  $(2, -1, 0)$ .

$$x = 2 - 9t, y = -1 - 5t, z = 22t$$

68. (a)  $\mathbf{n}_1 = 6\mathbf{i} - 3\mathbf{j} + \mathbf{k}$ ,  $\mathbf{n}_2 = -\mathbf{i} + \mathbf{j} + 5\mathbf{k}$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{|-4|}{\sqrt{46} \sqrt{27}} = \frac{2\sqrt{138}}{207}$$

$$\theta \approx 1.6845 \approx 96.52^\circ$$

(b) The direction vector for the line is

$$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 6 & -3 & 1 \\ -1 & 1 & 5 \end{vmatrix} = \langle -16, -31, 3 \rangle.$$

Find a point of intersection of the planes.

$$\begin{aligned} 6x - 3y + z &= 5 \Rightarrow 6x - 3y + z = 5 \\ -x + y + 5z &= 5 \Rightarrow -6x + 6y + 30z = 30 \\ \hline 3y + 31z &= 35 \end{aligned}$$

Let  $y = -9, z = 2 \Rightarrow x = -4 \Rightarrow (-4, -9, 2)$ .

$$x = -4 - 16t, y = -9 - 31t, z = 2 + 3t$$

69. The normal vectors to the planes are

$$\mathbf{n}_1 = \langle 5, -3, 1 \rangle, \mathbf{n}_2 = \langle 1, 4, 7 \rangle, \cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = 0.$$

So,  $\theta = \pi/2$  and the planes are orthogonal.

70. The normal vectors to the planes are

$$\mathbf{n}_1 = \langle 3, 1, -4 \rangle, \mathbf{n}_2 = \langle -9, -3, 12 \rangle.$$

Because  $\mathbf{n}_2 = -3\mathbf{n}_1$ , the planes are parallel, but not equal.

71. The normal vectors to the planes are

$$\mathbf{n}_1 = \mathbf{i} - 3\mathbf{j} + 6\mathbf{k}, \mathbf{n}_2 = 5\mathbf{i} + \mathbf{j} - \mathbf{k},$$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{|5 - 3 - 6|}{\sqrt{46}\sqrt{27}} = \frac{4\sqrt{138}}{414} = \frac{2\sqrt{138}}{207}.$$

$$\text{So, } \theta = \arccos\left(\frac{2\sqrt{138}}{207}\right) \approx 83.5^\circ.$$

72. The normal vectors to the planes are

$$\mathbf{n}_1 = 3\mathbf{i} + 2\mathbf{j} - \mathbf{k}, \mathbf{n}_2 = \mathbf{i} - 4\mathbf{j} + 2\mathbf{k},$$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{|3 - 8 - 2|}{\sqrt{14}\sqrt{21}} = \frac{7\sqrt{6}}{42} = \frac{\sqrt{6}}{6}.$$

$$\text{So, } \theta = \arccos\left(\frac{\sqrt{6}}{6}\right) \approx 65.9^\circ.$$

73. The normal vectors to the planes are  $\mathbf{n}_1 = \langle 1, -5, -1 \rangle$  and  $\mathbf{n}_2 = \langle 5, -25, -5 \rangle$ . Because  $\mathbf{n}_2 = 5\mathbf{n}_1$ , the planes are parallel, but not equal.


74. The normal vectors to the planes are

$$\mathbf{n}_1 = \langle 2, 0, -1 \rangle, \mathbf{n}_2 = \langle 4, 1, 8 \rangle,$$


$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = 0$$

So,  $\theta = \frac{\pi}{2}$  and the planes are orthogonal.


75.  $y \leq -2$




76.  $z = 1$




77.  $x + z = 6$




78.  $2x + y = 8$




79.  $4x + 2y + 6z = 12$




80.  $3x + 6y + 2z = 6$



81.  $2x - y + 3z = 4$



82.  $2x - y + z = 4$



83. Writing the equation of the line in parametric form and substituting into the equation of the plane you have:

$$x = -7 + 2t, y = 4 + t, z = -1 + 5t$$

$$(-7 + 2t) + 3(4 + t) - (-1 + 5t) = 6$$

$$6 = 6$$

The equation is valid for all  $t$ .

The line lies in the plane.

84. Writing the equation of the line in parametric form and substituting into the equation of the plane you have:

$$x = 1 + 4t, y = 2t, z = 3 + 6t$$

$$2(1 + 4t) + 3(2t) = -5, t = \frac{-1}{2}$$

Substituting  $t = -\frac{1}{2}$  into the parametric equations for the line you have the point of intersection  $(-1, -1, 0)$ .

The line does not lie in the plane.

85. Writing the equation of the line in parametric form and substituting into the equation of the plane you have:

$$x = 1 + 3t, y = -1 - 2t, z = 3 + t$$

$$2(1 + 3t) + 3(-1 - 2t) = 10, -1 = 10, \text{ contradiction}$$

So, the line does not intersect the plane.

86. Writing the equation of the line in parametric form and substituting into the equation of the plane you have:

$$x = 4 + 2t, y = -1 - 3t, z = -2 + 5t$$

$$5(4 + 2t) + 3(-1 - 3t) = 17, t = 0$$

Substituting  $t = 0$  into the parametric equations for the line you have the point of intersection  $(4, -1, -2)$ .

The line does not lie in the plane.

87. Point:  $Q(0, 0, 0)$

Plane:  $2x + 3y + z - 12 = 0$

Normal to plane:  $\mathbf{n} = \langle 2, 3, 1 \rangle$

Point in plane:  $P(6, 0, 0)$

Vector  $\overrightarrow{PQ} = \langle -6, 0, 0 \rangle$

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|-12|}{\sqrt{14}} = \frac{6\sqrt{14}}{7}$$

88. Point:  $Q(0, 0, 0)$

Plane:  $5x + y - z - 9 = 0$

Normal to plane:  $\mathbf{n} = \langle 5, 1, -1 \rangle$

Point in plane:  $P(0, 9, 0)$

Vector  $\overrightarrow{PQ} = \langle 0, -9, 0 \rangle$

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|-9|}{\sqrt{27}} = \sqrt{3}$$

89. Point:  $Q(2, 8, 4)$

Plane:  $2x + y + z = 5$

Normal to plane:  $\mathbf{n} = \langle 2, 1, 1 \rangle$

Point in plane:  $P(0, 0, 5)$

Vector:  $\overrightarrow{PQ} = \langle 2, 8, -1 \rangle$

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{11}{\sqrt{6}} = \frac{11\sqrt{6}}{6}$$

90. Point:  $Q(1, 3, -1)$

Plane:  $3x - 4y + 5z - 6 = 0$

Normal to plane:  $\mathbf{n} = \langle 3, -4, 5 \rangle$

Point in plane:  $P(2, 0, 0)$

Vector  $\overrightarrow{PQ}: \langle -1, 3, -1 \rangle$

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|-20|}{\sqrt{50}} = 2\sqrt{2}$$

91. The normal vectors to the planes are  $\mathbf{n}_1 = \langle 1, -3, 4 \rangle$  and

$\mathbf{n}_2 = \langle 1, -3, 4 \rangle$ . Because  $\mathbf{n}_1 = \mathbf{n}_2$ , the planes are

parallel. Choose a point in each plane.

$P(10, 0, 0)$  is a point in  $x - 3y + 4z = 10$ .

$Q(6, 0, 0)$  is a point in  $x - 3y + 4z = 6$ .

$$\overrightarrow{PQ} = \langle -4, 0, 0 \rangle, D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}_1|}{\|\mathbf{n}_1\|} = \frac{4}{\sqrt{26}} = \frac{2\sqrt{26}}{13}$$

92. The normal vectors to the planes are  $\mathbf{n}_1 = \langle 2, 7, 1 \rangle$  and

$\mathbf{n}_2 = \langle 2, 7, 1 \rangle$ . Because  $\mathbf{n}_1 = \mathbf{n}_2$ , the planes are parallel.

Choose a point in each plane.

$P(0, 0, 13)$  is a point in  $2x + 7y + z = 13$ .

$Q(0, 0, 9)$  is a point in  $2x + 7y + z = 9$ .

$\overrightarrow{PQ} = \langle 0, 0, 4 \rangle$

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}_1|}{\|\mathbf{n}_1\|} = \frac{4}{\sqrt{54}} = \frac{2\sqrt{6}}{9}$$

93. The normal vectors to the planes are  $\mathbf{n}_1 = \langle -3, 6, 7 \rangle$  and  $\mathbf{n}_2 = \langle 6, -12, -14 \rangle$ . Because  $\mathbf{n}_2 = -2\mathbf{n}_1$ , the planes are parallel. Choose a point in each plane.

$P(0, -1, 1)$  is a point in  $-3x + 6y + 7z = 1$ .

$$Q\left(\frac{25}{6}, 0, 0\right) \text{ is a point in } 6x - 12y - 14z = 25.$$

$$\overrightarrow{PQ} = \left\langle \frac{25}{6}, 1, -1 \right\rangle$$

$$D = \frac{\|\overrightarrow{PQ} \cdot \mathbf{n}_1\|}{\|\mathbf{n}_1\|} = \frac{|-27/2|}{\sqrt{94}} = \frac{27}{2\sqrt{94}} = \frac{27\sqrt{94}}{188}$$

94. The normal vectors to the planes are  $\mathbf{n}_1 = \langle -1, 6, 2 \rangle$  and  $\mathbf{n}_2 = \left\langle -\frac{1}{2}, 3, 1 \right\rangle$ . Because  $\mathbf{n}_1 = 2\mathbf{n}_2$ , the planes are parallel. Choose a point in each plane.

$P(-3, 0, 0)$  is a point in  $-x + 6y + 2z = 3$ .

$$Q(0, 0, 4) \text{ is a point in } -\frac{1}{2}x + 3y + z = 4.$$

$$\overrightarrow{PQ} = \langle 3, 0, 4 \rangle$$

$$D = \frac{\|\overrightarrow{PQ} \cdot \mathbf{n}_1\|}{\|\mathbf{n}_1\|} = \frac{5}{\sqrt{41}} = \frac{5\sqrt{41}}{41}$$

95.  $\mathbf{u} = \langle 4, 0, -1 \rangle$  is the direction vector for the line.

$Q(1, 5, -2)$  is the given point, and  $P(-2, 3, 1)$  is on the line.

$$\overrightarrow{PQ} = \langle 3, 2, -3 \rangle$$

$$\overrightarrow{PQ} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 2 & -3 \\ 4 & 0 & -1 \end{vmatrix} = \langle -2, -9, -8 \rangle$$

$$D = \frac{\|\overrightarrow{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|} = \frac{\sqrt{149}}{\sqrt{17}} = \frac{\sqrt{2533}}{17}$$

96.  $\mathbf{u} = \langle 2, 1, 2 \rangle$  is the direction vector for the line.

$Q(1, -2, 4)$  is the given point, and  $P(0, -3, 2)$  is a point on the line (let  $t = 0$ ).

$$\overrightarrow{PQ} = \langle 1, 1, 2 \rangle$$

$$\overrightarrow{PQ} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 2 \\ 2 & 1 & 2 \end{vmatrix} = \langle 0, 2, -1 \rangle$$

$$D = \frac{\|\overrightarrow{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|} = \frac{\sqrt{5}}{\sqrt{9}} = \frac{\sqrt{5}}{3}$$

97.  $\mathbf{u} = \langle -1, 1, -2 \rangle$  is the direction vector for the line.

$Q(-2, 1, 3)$  is the given point, and  $P(1, 2, 0)$  is on the line (let  $t = 0$  in the parametric equations for the line).

$$\overrightarrow{PQ} = \langle -3, -1, 3 \rangle$$

$$\overrightarrow{PQ} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -3 & -1 & 3 \\ -1 & 1 & -2 \end{vmatrix} = \langle -1, -9, -4 \rangle$$

$$D = \frac{\|\overrightarrow{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|} = \frac{\sqrt{1+81+16}}{\sqrt{1+1+4}} = \frac{\sqrt{98}}{\sqrt{6}} = \frac{7}{\sqrt{3}} = \frac{7\sqrt{3}}{3}$$

98.  $\mathbf{u} = \langle 0, 3, 1 \rangle$  is the direction vector for the line.

$Q(4, -1, 5)$  is the given point, and  $P(3, 1, 1)$  is on the line.

$$\overrightarrow{PQ} = \langle 1, -2, 4 \rangle$$

$$\overrightarrow{PQ} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -2 & 4 \\ 0 & 3 & 1 \end{vmatrix} = \langle -14, -1, 3 \rangle$$

$$D = \frac{\|\overrightarrow{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|} = \frac{\sqrt{14^2 + 1 + 9}}{\sqrt{9+1}} = \sqrt{\frac{206}{10}} = \sqrt{\frac{103}{5}} = \frac{\sqrt{515}}{5}$$

99. The direction vector for  $L_1$  is  $\mathbf{v}_1 = \langle -1, 2, 1 \rangle$ .

The direction vector for  $L_2$  is  $\mathbf{v}_2 = \langle 3, -6, -3 \rangle$ .

Because  $\mathbf{v}_2 = -3\mathbf{v}_1$ , the lines are parallel.

Let  $Q(2, 3, 4)$  to be a point on  $L_1$  and  $P(0, 1, 4)$  a point on  $L_2$ .  $\overrightarrow{PQ} = \langle 2, 2, 0 \rangle$ .

$\mathbf{u} = \mathbf{v}_2$  is the direction vector for  $L_2$ .

$$\overrightarrow{PQ} \times \mathbf{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 2 & 0 \\ 3 & -6 & -3 \end{vmatrix} = \langle -6, 6, -18 \rangle$$

$$D = \frac{\|\overrightarrow{PQ} \times \mathbf{v}_2\|}{\|\mathbf{v}_2\|} = \frac{\sqrt{36 + 36 + 324}}{\sqrt{9+36+9}} = \sqrt{\frac{396}{54}} = \sqrt{\frac{22}{3}} = \frac{\sqrt{66}}{3}$$

100. The direction vector for  $L_1$  is  $\mathbf{v}_1 = \langle 6, 9, -12 \rangle$ .

The direction vector for  $L_2$  is  $\mathbf{v}_2 = \langle 4, 6, -8 \rangle$ .

Because  $\mathbf{v}_1 = \frac{3}{2}\mathbf{v}_2$ , the lines are parallel.

Let  $Q(3, -2, 1)$  to be a point on  $L_1$  and  $P(-1, 3, 0)$  a point on  $L_2$ .  $\overline{PQ} = \langle 4, -5, 1 \rangle$ .

$\mathbf{u} = \mathbf{v}_2$  is the direction vector for  $L_2$ .

$$\overline{PQ} \times \mathbf{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & -5 & 1 \\ 4 & 6 & -8 \end{vmatrix} = \langle 34, 36, 44 \rangle$$

$$\begin{aligned} D &= \frac{\|\overline{PQ} \times \mathbf{v}_2\|}{\|\mathbf{v}_2\|} \\ &= \frac{\sqrt{34^2 + 36^2 + 44^2}}{\sqrt{16 + 36 + 64}} \\ &= \frac{\sqrt{4388}}{\sqrt{116}} = \sqrt{\frac{1097}{29}} = \frac{\sqrt{31813}}{29} \end{aligned}$$

105.  $z = 0.23x + 0.14y + 6.85$

Year	2009	2010	2011	2012	2013	2014
$z(\text{Approx})$	18.93	19.46	20.31	21.10	21.58	22.62

The approximations are close to the actual values.

- (b) If  $x$  and  $y$  both increase, then so does  $z$ .

106. On one side you have the points  $(0, 0, 0)$ ,  $(18, 0, 0)$ , and  $(-3, -3, 24)$ .


$$\mathbf{n}_1 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 18 & 0 & 0 \\ -3 & -3 & 24 \end{vmatrix} = -432\mathbf{j} - 54\mathbf{k}$$

On the adjacent side you have the points  $(0, 0, 0)$ ,  $(0, 18, 0)$ , and  $(-3, -3, 24)$ .

$$\mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 18 & 0 \\ -3 & -3 & 24 \end{vmatrix} = 432\mathbf{i} + 54\mathbf{k}$$

$$\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{2916}{189,540} = \frac{1}{65}$$

$$\theta = \arccos \frac{1}{65} \approx 89.1^\circ$$



101. Exactly one plane contains the point and line. Select two points on the line and observe that three noncolinear points determine a unique plane.

102. There are an infinite number of planes orthogonal to a given plane in space.

103. Yes, Consider two points on one line, and a third distinct point on another line. Three distinct points determine a unique plane.

104. (a)  $ax + by + d = 0$  matches (iv). The plane is parallel to the  $z$ -axis.  
 (b)  $ax + d = 0$  matches (i). The plane is parallel to the  $yz$ -plane.  
 (c)  $cz + d = 0$  matches (ii). The plane is parallel to the  $xy$ -plane.  
 (d)  $ax + cz + d = 0$  matches (iii). The plane is parallel to the  $y$ -axis.

107.  $L_1: x_1 = 6 + t, y_1 = 8 - t, z_1 = 3 + t$

$L_2: x_2 = 1 + t, y_2 = 2 + t, z_2 = 2t$


(a) At  $t = 0$ , the first insect is at  $P_1(6, 8, 3)$  and the second insect is at  $P_2(1, 2, 0)$ .

$$\text{Distance} = \sqrt{(6-1)^2 + (8-2)^2 + (3-0)^2} = \sqrt{70} \approx 8.37 \text{ centimeters.}$$

$$(b) \text{Distance} = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2} = \sqrt{5^2 + (6-2t)^2 + (3-t)^2} = \sqrt{5t^2 - 30t + 70}, 0 \leq t \leq 10$$

(c) The distance is never zero.

(d) Using a graphing utility, the minimum distance is 5 centimeters when  $t = 3$  minutes.



108. First find the distance D from the point  $Q(-3, 2, 4)$  to the plane. Let  $P(4, 0, 0)$  be on the plane.

$\mathbf{n} = \langle 2, 4, -3 \rangle$  is the normal to the plane.

$$D = \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|\langle -7, 2, 4 \rangle \cdot \langle 2, 4, -3 \rangle|}{\sqrt{4+16+9}} = \frac{|-14+8-12|}{\sqrt{29}} = \frac{18}{\sqrt{29}} = \frac{18\sqrt{29}}{29}$$

The equation of the sphere with center  $(-3, 2, 4)$  and radius  $18\sqrt{29}/29$  is  $(x+3)^2 + (y-2)^2 + (z-4)^2 = \frac{324}{29}$ .

109. The direction vector  $\mathbf{v}$  of the line is the normal to the plane,  $\mathbf{v} = \langle 3, -1, 4 \rangle$ .

The parametric equations of the line are  $x = 5 + 3t$ ,  $y = 4 - t$ ,  $z = -3 + 4t$ .

To find the point of intersection, solve for  $t$  in the following equation:

$$\begin{aligned} 3(5 + 3t) - (4 - t) + 4(-3 + 4t) &= 7 \\ 26t &= 8 \\ t &= \frac{4}{13} \end{aligned}$$

Point of intersection:

$$\left(5 + 3\left(\frac{4}{13}\right), 4 - \frac{4}{13}, -3 + 4\left(\frac{4}{13}\right)\right) = \left(\frac{77}{13}, \frac{48}{13}, -\frac{23}{13}\right)$$

110. The normal to the plane,  $\mathbf{n} = \langle 2, -1, -3 \rangle$  is perpendicular to the direction vector  $\mathbf{v} = \langle 2, 4, 0 \rangle$  of the line because  $\langle 2, -1, -3 \rangle \cdot \langle 2, 4, 0 \rangle = 0$ .

So, the plane is parallel to the line. To find the distance between them, let  $Q(-2, -1, 4)$  be on the line and

$P(2, 0, 0)$  on the plane.  $\overline{PQ} = \langle -4, -1, 4 \rangle$ .

$$\begin{aligned} D &= \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} \\ &= \frac{|\langle -4, -1, 4 \rangle \cdot \langle 2, -1, -3 \rangle|}{\sqrt{4+1+9}} = \frac{19}{\sqrt{14}} = \frac{19\sqrt{14}}{14} \end{aligned}$$

$$111. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -5 & 1 \\ -3 & 1 & 4 \end{vmatrix} = -21\mathbf{i} - 11\mathbf{j} - 13\mathbf{k}$$

Direction numbers: 21, 11, 13

$$x = 21t, y = 1 + 11t, z = 4 + 13t$$

112. The unknown line  $L$  is perpendicular to the normal vector  $\mathbf{n} = \langle 1, 1, 1 \rangle$  of the plane, and perpendicular to the direction vector  $\mathbf{u} = \langle 1, 1, -1 \rangle$ . So, the direction vector of  $L$  is

$$\mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = \langle -2, 2, 0 \rangle.$$

The parametric equations for  $L$  are  $x = 1 - 2t$ ,  $y = 2t$ ,  $z = 2$ .

113. True

114. False. They may be skew lines. (See Section Project.)

115. True

116. False. For example, the lines  $x = t, y = 0, z = 1$  and  $x = 0, y = t, z = 1$  are both parallel to the plane  $z = 0$ , but the lines are not parallel.

117. False. For example, planes  $7x + y - 11z = 5$  and  $5x + 2y - 4z = 1$  are both perpendicular to plane  $2x - 3y + z = 3$ , but are not parallel.

118. True

## Section 11.6 Surfaces in Space

1. Quadric surfaces are the three-dimensional analogs of conic sections.

2. In the  $xz$ -plane,  $z = x^2$  is a parabola.

In three-space,  $z = x^2$  is a cylinder.

3. The trace of a surface is the intersection of the surface with a plane. You find a trace by setting one variable equal to a constant, such as  $x = 0$  or  $z = 2$ .

4. No. For example,  $x^2 + y^2 + z^2 = 0$  is a single point and  $x^2 + y^2 = 1$  is a right circular cylinder.

5. Ellipsoid

Matches graph (c)

6. Hyperboloid of two sheets

Matches graph (e)

7. Hyperboloid of one sheet

Matches graph (f)

8. Elliptic cone

Matches graph (b)

9. Elliptic paraboloid


Matches graph (d)

10. Hyperbolic paraboloid

Matches graph (a)


11.  $y^2 + z^2 = 9$

The  $x$ -coordinate is missing so you have a right circular cylinder with rulings parallel to the  $x$ -axis. The generating curve is a circle.



12.  $y^2 + z = 6$


The  $x$ -coordinate is missing so you have a parabolic cylinder with the rulings parallel to the  $x$ -axis. The generating curve is a parabola.



13.  $4x^2 + y^2 = 4$

$$\frac{x^2}{1} + \frac{y^2}{4} = 1$$


The  $z$ -coordinate is missing so you have an elliptic cylinder with rulings parallel to the  $z$ -axis. The generating curve is an ellipse.



14.  $y^2 - z^2 = 25$

$$\frac{y^2}{25} - \frac{z^2}{25} = 1$$

The  $x$ -coordinate is missing so you have a hyperbolic cylinder with rulings parallel to the  $x$ -axis. The generating curve is a hyperbola.




15.  $4x^2 - y^2 - z^2 = 1$

Hyperboloid of two sheets


xy-trace:  $4x^2 - y^2 = 1$  hyperbola

yz-trace: none

xz-trace:  $4x^2 - z^2 = 1$  hyperbola

16.  $\frac{x^2}{16} + \frac{y^2}{25} + \frac{z^2}{25} = 1$


Ellipsoid

xy-trace:  $\frac{x^2}{16} + \frac{y^2}{25} = 1$  ellipsexz-trace:  $\frac{x^2}{16} + \frac{z^2}{25} = 1$  ellipseyz-trace:  $y^2 + z^2 = 25$  circle

17.  $16x^2 - y^2 + 16z^2 = 4$


$4x^2 - \frac{y^2}{4} + 4z^2 = 1$

Hyperboloid of one sheet

xy-trace:  $4x^2 - \frac{y^2}{4} = 1$  hyperbolaxz-trace:  $4(x^2 + z^2) = 1$  circleyz-trace:  $\frac{-y^2}{4} + 4z^2 = 1$  hyperbola


18.  $z = x^2 + 4y^2$

Elliptic paraboloid

xy-trace: point  $(0, 0, 0)$ xz-trace:  $z = x^2$  parabolayz-trace:  $z = 4y^2$  parabola

19.  $\frac{x^2}{1} + \frac{y^2}{4} + \frac{z^2}{1} = 1$


Ellipsoid

xy-trace:  $\frac{x^2}{1} + \frac{y^2}{4} = 1$  ellipsexz-trace:  $x^2 + z^2 = 1$  circleyz-trace:  $\frac{y^2}{4} + \frac{z^2}{1} = 1$  ellipse

20.  $z^2 - x^2 - \frac{y^2}{4} = 1$


Hyperboloid of two sheets

xy-trace: none

xz-trace:  $z^2 - x^2 = 1$  hyperbolayz-trace:  $z^2 - \frac{y^2}{4} = 1$  hyperbola $z = \pm\sqrt{10}$ :  $\frac{x^2}{9} + \frac{y^2}{36} = 1$  ellipse


21.  $z^2 = x^2 + \frac{y^2}{9}$

Elliptic cone

xy-trace: point  $(0, 0, 0)$ xz-trace:  $z = \pm x$ yz-trace:  $z = \pm \frac{y}{3}$ When  $z = \pm 1$ ,  $x^2 + \frac{y^2}{9} = 1$  ellipse

22.  $3z = -y^2 + x^2$

Hyperbolic paraboloid

xy-trace:  $y = \pm x$ xz-trace:  $z = \frac{1}{3}x^2$ yz-trace:  $z = -\frac{1}{3}y^2$ 


23.  $x^2 - y^2 + z = 0$

Hyperbolic paraboloid  
 $xy$ -trace:  $y = \pm x$

$xz$ -trace:  $z = -x^2$

$yz$ -trace:  $z = y^2$

$$y = \pm 1: z = 1 - x^2$$




24.  $x^2 = 2y^2 + 2z^2$

Elliptic Cone

$xy$ -trace:  $x = \pm\sqrt{2}y$

$xz$ -trace:  $x = \pm\sqrt{2}z$

$yz$ -trace: point:  $(0, 0, 0)$



25.  $x^2 - y + z^2 = 0$

Elliptic paraboloid


$xy$ -trace:  $y = x^2$

$xz$ -trace:  $x^2 + z^2 = 0$ ,

point  $(0, 0, 0)$

$yz$ -trace:  $y = z^2$

$$y = 1: x^2 + z^2 = 1$$



26.  $-8x^2 + 18y^2 + 18z^2 = 2$


$$9y^2 + 9z^2 - 4x^2 = 1$$

Hyperboloid of one sheet

$xy$ -trace:  $9y^2 - 4x^2 = 1$  hyperbola

$yz$ -trace:  $9y^2 + 9z^2 = 1$  circle

$xz$ -trace:  $9z^2 - 4x^2 = 1$  hyperbola



27. These have to be two minus signs in order to have a hyperboloid of two sheets. The number of sheets is the same as the number of minus signs.

28. Yes. Every trace is an ellipse (or circle or point).

29. No. See the table on pages 800 and 801.

30.  $z = x^2 + y^2$

(a) You are viewing the paraboloid from the  $x$ -axis:  
 $(20, 0, 0)$

(b) You are viewing the paraboloid from above, but not on the  $z$ -axis:  $(10, 10, 20)$

(c) You are viewing the paraboloid from the  $z$ -axis:  
 $(0, 0, 20)$

(d) You are viewing the paraboloid from the  $y$ -axis:  
 $(0, 20, 0)$

31.  $x^2 + z^2 = [r(y)]^2$  and  $z = r(y) = 5y$ , so

$$x^2 + z^2 = 25y^2.$$

32.  $x^2 + z^2 = [r(y)]^2$  and  $z = r(y) \pm 3\sqrt{y}$ , so

$$x^2 + z^2 = 9y.$$

33.  $x^2 + y^2 = [r(z)]^2$  and  $y = r(z) = 2z^{1/3}$ , so

$$x^2 + y^2 = 4z^{2/3}.$$

34.  $x^2 + y^2 = [r(z)]^2$  and  $x = r(z) = e^z$ , so

$$x^2 + y^2 = e^{2z}.$$

35.  $y^2 + z^2 = [r(x)]^2$  and  $y = r(x) = \frac{2}{x}$ , so

$$y^2 + z^2 = \left(\frac{2}{x}\right)^2 \Rightarrow y^2 + z^2 = \frac{4}{x^2}.$$

36.  $y^2 + z^2 = [r(x)]^2$  and  $z = r(x) = \frac{1}{2}\sqrt{4 - x^2}$ , so

$$y^2 + z^2 = \frac{1}{4}(4 - x^2) \Rightarrow x^2 + 4y^2 + 4z^2 = 4.$$

37.  $x^2 + y^2 - 2z = 0$

$$x^2 + y^2 = (\sqrt{2z})^2$$

Equation of generating curve:  $y = \sqrt{2z}$  or  $x = \sqrt{2z}$

38.  $x^2 + z^2 = \cos^2 y$

Equation of generating curve:  $x = \cos y$  or  $z = \cos y$

39.  $y^2 + z^2 = 5 - 8x^2 = (\sqrt{5 - 8x^2})^2$

Equation of generating curve:  $y = \sqrt{5 - 8x^2}$  or

$$z = \sqrt{5 - 8x^2}$$


40.  $6x^2 + 2y^2 + 2z^2 = 1$

$$y^2 + z^2 = \frac{1}{2} - 3x^2 = \left(\sqrt{\frac{1}{2} - 3x^2}\right)^2$$

Equation of generating curve:  $y = \sqrt{\frac{1}{2} - 3x^2}$  or


$$z = \sqrt{\frac{1}{2} - 3x^2}$$

41.  $V = 2\pi \int_0^4 x(4x - x^2) dx = 2\pi \left[ \frac{4x^3}{3} - \frac{x^4}{4} \right]_0^4 = \frac{218\pi}{3}$



42.  $V = 2\pi \int_0^\pi y \sin y dy$

$$= 2\pi [\sin y - y \cos y]_0^\pi = 2\pi^2$$



43.  $z = \frac{x^2}{2} + \frac{y^2}{4}$

(a) When  $z = 2$  we have  $2 = \frac{x^2}{2} + \frac{y^2}{4}$ , or

$$1 = \frac{x^2}{4} + \frac{y^2}{8}$$

Major axis:  $2\sqrt{8} = 4\sqrt{2}$

Minor axis:  $2\sqrt{4} = 4$

$$c^2 = a^2 - b^2, c^2 = 4, c = 2$$

Foci:  $(0, \pm 2, 2)$

(b) When  $z = 8$  we have  $8 = \frac{x^2}{2} + \frac{y^2}{4}$ , or

$$1 = \frac{x^2}{16} + \frac{y^2}{32}.$$

Major axis:  $2\sqrt{32} = 8\sqrt{2}$

Minor axis:  $2\sqrt{16} = 8$

$$c^2 = 32 - 16 = 16, c = 4$$

Foci:  $(0, \pm 4, 8)$

44.  $z = \frac{x^2}{2} + \frac{y^2}{4}$

(a) When  $y = 4$  you have  $z = \frac{x^2}{2} + 4$ ,

$$4\left(\frac{1}{2}\right)(z - 4) = x^2.$$

Focus:  $\left(0, 4, \frac{9}{2}\right)$

(b) When  $x = 2$  you have

$$z = 2 + \frac{y^2}{4}, 4(z - 2) = y^2.$$

Focus:  $(2, 0, 3)$

45. If  $(x, y, z)$  is on the surface, then

$$(y + 2)^2 = x^2 + (y - 2)^2 + z^2$$

$$y^2 + 4y + 4 = x^2 + y^2 - 4y + 4 + z^2$$

$$x^2 + z^2 = 8y$$

Elliptic paraboloid

Traces parallel to  $xz$ -plane are circles.

46. If  $(x, y, z)$  is on the surface, then

$$z^2 = x^2 + y^2 + (z - 4)^2$$


$$z^2 = x^2 + y^2 + z^2 - 8z + 16$$

$$8z = x^2 + y^2 + 16 \Rightarrow z = \frac{x^2}{8} + \frac{y^2}{8} + 2$$


Elliptic paraboloid shifted up 2 units. Traces parallel to  $xy$ -plane are circles.

47.  $\frac{x^2}{6378^2} + \frac{y^2}{6378^2} + \frac{z^2}{6357^2} = 1$


48. (a)  $x^2 + y^2 = [r(z)]^2$   
 $= [\sqrt{2(z-1)}]^2$   
 $x^2 + y^2 - 2z + 2 = 0$



(b)  $V = 2\pi \int_0^2 x \left[ 3 - \left( \frac{1}{2}x^2 + 1 \right) \right] dx$   
 $= 2\pi \int_0^2 \left( 2x - \frac{1}{2}x^3 \right) dx$   
 $= 2\pi \left[ x^2 - \frac{x^4}{8} \right]_0^2 = 4\pi \approx 12.6 \text{ cm}^3$



(c)  $V = 2\pi \int_{1/2}^2 x \left[ 3 - \left( \frac{1}{2}x^2 + 1 \right) \right] dx$   
 $= 2\pi \int_{1/2}^2 \left( 2x - \frac{1}{2}x^3 \right) dx$   
 $= 2\pi \left[ x^2 - \frac{x^4}{8} \right]_{1/2}^2$   
 $= 4 - \frac{31\pi}{64} = \frac{225\pi}{64} \approx 11.04 \text{ cm}^3$



49.  $z = \frac{y^2}{b^2} - \frac{x^2}{a^2}, z = bx + ay$   
 $bx + ay = \frac{y^2}{b^2} - \frac{x^2}{a^2}$   
 $\frac{1}{a^2} \left( x^2 + a^2bx + \frac{a^4b^2}{4} \right) = \frac{1}{b^2} \left( y^2 - ab^2y + \frac{a^2b^4}{4} \right)$   
 $\frac{\left( x + \frac{a^2b}{2} \right)^2}{a^2} = \frac{\left( y - \frac{ab^2}{2} \right)^2}{b^2}$   
 $y = \pm \frac{b}{a} \left( x + \frac{a^2b}{2} \right) + \frac{ab^2}{2}$

Letting  $x = at$ , you obtain the two intersecting lines  
 $x = at, y = -bt, z = 0$  and  $x = at,$   
 $y = bt + ab^2, z = 2abt + a^2b^2$ .

50. Equating twice the first equation with the second equation:  
 $2x^2 + 6y^2 - 4z^2 + 4y - 8 = 2x^2 + 6y^2 - 4z^2 - 3x - 2$   
 $4y - 8 = -3x - 2$   
 $3x + 4y = 6$ , a plane

51. The Klein bottle does not have both an “inside” and an “outside.” It is formed by inserting the small open end through the side of the bottle and making it contiguous with the top of the bottle.

## Section 11.7 Cylindrical and Spherical Coordinates

- The cylindrical coordinate system is an extension of the polar coordinate system. In this system, a point P in space is represented by an ordered triple  $(r, \theta, z)$ .  $(r, \theta)$  is a polar representation of the projection of P in the  $xy$ -plane, and  $z$  is the directed distance from  $(r, \theta)$  to P.
- The point is 2 units from the origin, in the  $xz$ -plane, and makes an angle of  $30^\circ$  with the  $z$ -axis.

3.  $(-7, 0, 5)$ , cylindrical

$$x = r \cos \theta = -7 \cos 0 = -7$$

$$y = r \sin \theta = -7 \sin 0 = 0$$

$$z = 5$$

- $(-7, 0, 5)$ , rectangular

4.  $(2, -\pi, -4)$ , cylindrical

$$x = r \cos \theta = 2 \cos(-\pi) = -2$$

$$y = r \sin \theta = 2 \sin(-\pi) = 0$$

$$z = -4$$

 $(-2, 0, -4)$ , rectangular5.  $\left(3, \frac{\pi}{4}, 1\right)$ , cylindrical

$$x = r \cos \theta = 3 \cos \frac{\pi}{4} = \frac{3\sqrt{2}}{2}$$

$$y = r \sin \theta = 3 \sin \frac{\pi}{4} = \frac{3\sqrt{2}}{2}$$

$$z = 1$$

 $\left(\frac{3\sqrt{2}}{2}, \frac{3\sqrt{2}}{2}, 1\right)$ , rectangular6.  $\left(6, -\frac{3\pi}{2}, 2\right)$ , cylindrical

$$x = r \cos \theta = 6 \cos\left(-\frac{3\pi}{2}\right) = 0$$

$$y = r \sin \theta = 6 \sin\left(-\frac{3\pi}{2}\right) = 6$$

$$z = 2$$

 $(0, 6, 2)$ , rectangular7.  $\left(4, \frac{7\pi}{6}, -3\right)$ , cylindrical

$$x = r \cos \theta = 4 \cos \frac{7\pi}{6} = 4\left(-\frac{\sqrt{3}}{2}\right) = -2\sqrt{3}$$

$$y = r \sin \theta = 4 \sin \frac{7\pi}{6} = 4\left(-\frac{1}{2}\right) = -2$$

$$z = -3$$

 $(-2\sqrt{3}, -2, -3)$ , rectangular8.  $\left(-\frac{2}{3}, \frac{4\pi}{3}, 8\right)$ , cylindrical

$$x = r \cos \theta = -\frac{2}{3} \cos \frac{4\pi}{3} = \left(-\frac{2}{3}\right)\left(-\frac{1}{2}\right) = \frac{1}{3}$$

$$y = r \sin \theta = -\frac{2}{3} \sin \frac{4\pi}{3} = \left(-\frac{2}{3}\right)\left(-\frac{\sqrt{3}}{2}\right) = \frac{\sqrt{3}}{3}$$

$$z = 8$$

 $\left(\frac{1}{3}, \frac{\sqrt{3}}{3}, 8\right)$ , rectangular9.  $(0, 5, 1)$ , rectangular

$$r = \sqrt{(0)^2 + (5)^2} = 5$$

$$\tan \theta = \frac{5}{2} \Rightarrow \theta = \arctan \frac{5}{0} = \frac{\pi}{2}$$

$$z = 1$$

 $\left(5, \frac{\pi}{2}, 1\right)$ , cylindrical10.  $(6, 2\sqrt{3}, -1)$ , rectangular

$$r = \sqrt{6^2 + (2\sqrt{3})^2} = \sqrt{36 + 12} = \sqrt{48} = 4\sqrt{3}$$

$$\tan \theta = \frac{2\sqrt{3}}{6} = \frac{\sqrt{3}}{3} \Rightarrow \theta = \arctan \frac{\sqrt{3}}{3} = \frac{\pi}{6}$$

$$z = -1$$

 $\left(4\sqrt{3}, \frac{\pi}{6}, -1\right)$ , cylindrical11.  $(2, -2, -4)$ , rectangular

$$r = \sqrt{2^2 + (-2)^2} = 2\sqrt{2}$$

$$\tan \theta = \frac{-2}{2} \Rightarrow \theta = \arctan(-1) = -\frac{\pi}{4}$$

$$z = -4$$

 $\left(2\sqrt{2}, -\frac{\pi}{4}, -4\right)$ , cylindrical12.  $(3, -3, 7)$ , rectangular

$$r = \sqrt{3^2 + (-3)^2} = \sqrt{18} = 3\sqrt{2}$$

$$\tan \theta = \frac{-3}{3} \Rightarrow \theta = \arctan(-1) = -\frac{\pi}{4}$$

$$z = 7$$

 $\left(3\sqrt{2}, -\frac{\pi}{4}, 7\right)$ , cylindrical13.  $(1, \sqrt{3}, 4)$ , rectangular

$$r = \sqrt{1^2 + (\sqrt{3})^2} = 2$$

$$\tan \theta = \frac{\sqrt{3}}{1} \Rightarrow \theta = \arctan \sqrt{3} = \frac{\pi}{3}$$

$$z = 4$$

 $\left(2, \frac{\pi}{3}, 4\right)$ , cylindrical

14.  $(2\sqrt{3}, -2, 6)$ , rectangular

$$r = \sqrt{12 + 4} = 4$$

$$\theta = \arctan\left(-\frac{1}{\sqrt{3}}\right) = \frac{5\pi}{6}$$

$$z = 6$$

$$\left(4, -\frac{\pi}{6}, 6\right), \text{cylindrical}$$

15.  $z = 4$  is the equation in cylindrical coordinates.  
(plane)

16.  $x = 9$ , rectangular equation

$$r \cos \theta = 9$$

$$r = 9 \sec \theta, \text{cylindrical equation}$$

17.  $x^2 + y^2 - 2z^2 = 5$ , rectangular equation

$$r^2 - 2z^2 = 5, \text{cylindrical equation}$$

18.  $z = x^2 + y^2 - 11$ , rectangular equation

$$z = r^2 - 11, \text{cylindrical equation}$$

19.  $y = x^2$ , rectangular equation

$$r \sin \theta = (r \cos \theta)^2$$

$$\sin \theta = r \cos^2 \theta$$

$$r = \sec \theta \cdot \tan \theta, \text{cylindrical equation}$$

20.  $x^2 + y^2 = 8x$ , rectangular equation

$$r^2 = 8r \cos \theta$$

$$r = 8 \cos \theta, \text{cylindrical equation}$$

21.  $y^2 = 10 - z^2$ , rectangular equation

$$(r \sin \theta)^2 = 10 - z^2$$

$$r^2 \sin^2 \theta + z^2 = 10, \text{cylindrical equation}$$


22.  $x^2 + y^2 + z^2 - 3z = 0$ , rectangular equation

$$r^2 + z^2 - 3z = 0, \text{cylindrical equation}$$

23.  $r = 3$ , cylindrical equation


$$\sqrt{x^2 + y^2} = 3$$

$$x^2 + y^2 = 9, \text{rectangular equation}$$



24.  $z = -2$ , cylindrical equation

$$z = -2, \text{rectangular equation}$$




25.  $\theta = \frac{\pi}{6}$ , cylindrical equation

$$\tan \frac{\pi}{6} = \frac{y}{x}$$

$$\frac{1}{\sqrt{3}} = \frac{y}{x}$$

$$x = \sqrt{3}y$$


$$x - \sqrt{3}y = 0, \text{rectangular equation}$$



26.  $r = \frac{z}{2}$ , cylindrical equation


$$\sqrt{x^2 + y^2} = \frac{z}{2}$$

$$x^2 + y^2 - \frac{z^2}{4} = 0, \text{rectangular equation}$$




27.  $r^2 + z^2 = 5$ , cylindrical equation

$$x^2 + y^2 + z^2 = 5, \text{rectangular equation}$$



28.  $z = r^2 \cos^2 \theta$ , cylindrical equation

$$z = x^2, \text{ rectangular equation}$$



29.


$r = 4 \sin \theta, \text{ cylindrical equation}$

$$r^2 = 4r \sin \theta$$

$$x^2 + y^2 = 4y$$

$$x^2 + y^2 - 4y + 4 = 4$$

$$x^2 + (y - 2)^2 = 4, \text{ rectangular equation}$$



30.


$r = 2 \cos \theta, \text{ cylindrical equation}$

$$r^2 = 2r \cos \theta$$

$$x^2 + y^2 = 2x$$

$$x^2 + y^2 - 2x = 0$$

$$(x - 1)^2 + y^2 = 1, \text{ rectangular equation}$$



31.  $(4, 0, 0)$ , rectangular

$$\rho = \sqrt{4^2 + 0^2 + 0^2} = 4$$

$$\tan \theta = \frac{y}{x} = 0 \Rightarrow \theta = 0$$

$$\phi = \arccos 0 = \frac{\pi}{2}$$

$$\left(4, 0, \frac{\pi}{2}\right), \text{spherical}$$

32.  $(-4, 0, 0)$ , rectangular

$$\rho = \sqrt{(-4)^2 + 0^2 + 0^2} = 4$$

$$\tan \theta = \frac{y}{x} = 0 \Rightarrow \theta = 0$$

$$\phi = \arccos \frac{z}{\rho} = \arccos 0 = \frac{\pi}{2}$$

$$\left(4, 0, \frac{\pi}{2}\right), \text{spherical}$$

33.  $(-2, 2\sqrt{3}, 4)$ , rectangular

$$\rho = \sqrt{(-2)^2 + (2\sqrt{3})^2 + 4^2} = 4\sqrt{2}$$

$$\tan \theta = \frac{y}{x} = \frac{2\sqrt{3}}{-2} = -\sqrt{3}$$

$$\theta = \frac{2\pi}{3}$$

$$\phi = \arccos \frac{z}{\rho} = \arccos \frac{1}{\sqrt{2}} = \frac{\pi}{4}$$

$$\left(4\sqrt{2}, \frac{2\pi}{3}, \frac{\pi}{4}\right), \text{spherical}$$

34.  $(-5, -5, \sqrt{2})$ , rectangular

$$\rho = \sqrt{(-5)^2 + (-5)^2 + (\sqrt{2})^2} = \sqrt{52} = 2\sqrt{13}$$

$$\tan \theta = \frac{y}{x} = \frac{-5}{-5} = 1 \Rightarrow \theta = \frac{\pi}{4}$$

$$\phi = \arccos \frac{z}{\rho} = \arccos \frac{\sqrt{2}}{2\sqrt{13}} = \arccos \frac{\sqrt{26}}{26}$$

$$\left(2\sqrt{13}, \frac{\pi}{4}, \arccos \frac{\sqrt{26}}{26}\right), \text{spherical}$$

35.  $(\sqrt{3}, 1, 2\sqrt{3})$ , rectangular

$$\rho = \sqrt{3 + 1 + 12} = 4$$

$$\tan \theta = \frac{y}{x} = \frac{1}{\sqrt{3}}$$

$$\theta = \frac{\pi}{6}$$

$$\phi = \arccos \frac{z}{\rho} = \arccos \frac{\sqrt{3}}{2} = \frac{\pi}{6}$$

$$\left(4, \frac{\pi}{6}, \frac{\pi}{6}\right), \text{spherical}$$

36.  $(-1, 2, 1)$ , rectangular

$$\rho = \sqrt{(-1)^2 + 2^2 + 1^2} = \sqrt{6}$$

$$\tan \theta = \frac{y}{x} = -2 \Rightarrow \theta = \arctan(-2) + \pi$$

$$\phi = \arccos \frac{z}{\rho} = \arccos \frac{1}{\sqrt{6}}$$

$$\left( \sqrt{6}, \arctan(-2) + \pi, \arccos \frac{1}{\sqrt{6}} \right), \text{spherical}$$

37.  $\left( 4, \frac{\pi}{6}, \frac{\pi}{4} \right)$ , spherical

$$x = \rho \sin \phi \cos \theta = 4 \sin \frac{\pi}{6} \cos \frac{\pi}{4} = \sqrt{6}$$

$$y = \rho \sin \phi \sin \theta = 4 \sin \frac{\pi}{6} \sin \frac{\pi}{4} = \sqrt{2}$$

$$z = \rho \cos \phi = 4 \cos \frac{\pi}{6} = 2\sqrt{2}$$

$$(\sqrt{6}, \sqrt{2}, 2\sqrt{2}), \text{rectangular}$$

38.  $\left( 6, \pi, \frac{\pi}{2} \right)$ , spherical

$$x = \rho \sin \phi \cos \theta = 6 \sin \frac{\pi}{2} \cos \pi = -6$$

$$y = \rho \sin \phi \sin \theta = 6 \sin \frac{\pi}{2} \sin \pi = 0$$

$$z = \rho \cos \phi = 6 \cos \frac{\pi}{2} = 0$$

$$(-6, 0, 0), \text{rectangular}$$

39.  $\left( 12, -\frac{\pi}{4}, 0 \right)$ , spherical

$$x = \rho \sin \phi \cos \theta = 12 \sin 0 \cos \left( -\frac{\pi}{4} \right) = 0$$

$$y = \rho \sin \phi \sin \theta = 12 \sin 0 \sin \left( -\frac{\pi}{4} \right) = 0$$

$$z = \rho \cos \phi = 12 \cos 0 = 12$$

$$(0, 0, 12), \text{rectangular}$$

40.  $\left( 9, \frac{\pi}{4}, \pi \right)$ , spherical

$$x = \rho \sin \phi \cos \theta = 9 \sin \pi \cos \frac{\pi}{4} = 0$$

$$y = \rho \sin \phi \sin \theta = 9 \sin \pi \sin \frac{\pi}{4} = 0$$

$$z = \rho \cos \phi = 9 \cos \pi = -9$$

$$(0, 0, -9), \text{rectangular}$$

41.  $\left( 5, \frac{\pi}{4}, \frac{\pi}{12} \right)$ , spherical

$$x = \rho \sin \phi \cos \theta = 5 \sin \frac{\pi}{12} \cos \frac{\pi}{4} \approx 0.915$$

$$y = \rho \sin \phi \sin \theta = 5 \sin \frac{\pi}{12} \sin \frac{\pi}{4} \approx 0.915$$

$$z = \rho \cos \theta = 5 \cos \frac{\pi}{12} \approx 4.830$$

$$(0.915, 0.915, 4.830), \text{rectangular}$$

42.  $\left( 7, \frac{3\pi}{4}, \frac{\pi}{9} \right)$ , spherical

$$x = \rho \sin \phi \cos \theta = 7 \sin \frac{\pi}{9} \cos \frac{3\pi}{4} \approx -1.693$$

$$y = \rho \sin \phi \sin \theta = 7 \sin \frac{\pi}{9} \sin \frac{3\pi}{4} \approx 1.693$$

$$z = \rho \cos \phi = 7 \cos \frac{\pi}{9} \approx 6.578$$

$$(-1.693, 1.693, 6.578), \text{rectangular}$$

43.  $y = 2$ , rectangular equation

$$\rho \sin \phi \sin \theta = 2$$

$$\rho = 2 \csc \phi \csc \theta, \text{spherical equation}$$

44.  $z = 6$ , rectangular equation

$$\rho \cos \phi = 6$$

$$\rho = 6 \sec \phi, \text{spherical equation}$$

45.  $x^2 + y^2 + z^2 = 49$ , rectangular equation

$$\rho^2 = 49$$

$$\rho = 7, \text{spherical equation}$$

46.  $x^2 + y^2 - 3z^2 = 0$ , rectangular equation

$$x^2 + y^2 + z^2 = 4z^2$$

$$\rho^2 = 4\rho^2 \cos^2 \phi$$

$$1 = 4 \cos^2 \phi$$

$$\cos \phi = \frac{1}{2}$$

$$\phi = \frac{\pi}{3}, (\text{cone}) \text{spherical equation}$$

47.  $x^2 + y^2 = 16$ , rectangular equation

$$\rho^2 \sin^2 \phi \sin^2 \theta + \rho^2 \sin^2 \phi \cos^2 \theta = 16$$

$$\rho^2 \sin^2 \phi (\sin^2 \theta + \cos^2 \theta) = 16$$

$$\rho^2 \sin^2 \phi = 16$$

$$\rho \sin \phi = 4$$

$$\rho = 4 \csc \phi, \text{spherical equation}$$

48.  $x = 13$ , rectangular equation

$$\rho \sin \phi \cos \theta = 13$$

$$\rho = 13 \csc \phi \sec \theta, \text{spherical equation}$$

49.  $x^2 + y^2 = 2z^2$ , rectangular equation

$$\rho^2 \sin^2 \phi \cos^2 \theta + \rho^2 \sin^2 \phi \sin^2 \theta = 2\rho^2 \cos^2 \phi$$

$$\rho^2 \sin^2 \phi [\cos^2 \theta + \sin^2 \theta] = 2\rho^2 \cos^2 \phi$$

$$\rho^2 \sin^2 \phi = 2\rho^2 \cos^2 \theta$$

$$\frac{\sin^2 \phi}{\cos^2 \phi} = 2$$

$$\tan^2 \phi = 2$$

$$\tan \phi = \pm \sqrt{2}, \text{spherical equation}$$

50.  $x^2 + y^2 + z^2 - 9z = 0$ , rectangular equation

$$\rho^2 - 9\rho \cos \phi = 0$$

$$\rho = 9 \cos \phi, \text{spherical equation}$$


53.  $\phi = \frac{\pi}{6}$ , spherical equation

$$\cos \phi = \frac{z}{\sqrt{x^2 + y^2 + z^2}}$$

$$\frac{\sqrt{3}}{2} = \frac{z}{\sqrt{x^2 + y^2 + z^2}}$$

$$\frac{3}{4} = \frac{z^2}{x^2 + y^2 + z^2}$$

$3x^2 + 3y^2 - z^2 = 0, z \geq 0$ , rectangular equation





52.  $\theta = \frac{3\pi}{4}$ , spherical equation

$$\tan \theta = \frac{y}{x}$$

$$-1 = \frac{y}{x}$$

$x + y = 0$ , rectangular equation




54.  $\phi = \frac{\pi}{2}$ , spherical equation

$$\cos \phi = \frac{z}{\sqrt{x^2 + y^2 + z^2}}$$

$$0 = \frac{z}{\sqrt{x^2 + y^2 + z^2}}$$

$z = 0$ , rectangular equation

$xy$ -plane




55.  $\rho = 4 \cos \phi$ , spherical equation

$$\sqrt{x^2 + y^2 + z^2} = \frac{4z}{\sqrt{x^2 + y^2 + z^2}}$$

$$x^2 + y^2 + z^2 - 4z = 0$$


$$x^2 + y^2 + (z - 2)^2 = 4, z \geq 0, \text{ rectangular equation}$$



56.  $\rho = 2 \sec \phi$ , spherical equation

$$\rho \cos \phi = 2$$

$$z = 2, \text{ rectangular equation}$$




57.  $\rho = \csc \phi$ , spherical equation

$$\rho \sin \phi = 1$$

$$\sqrt{x^2 + y^2} = 1$$

$$x^2 + y^2 = 1, \text{ rectangular equation}$$




58.  $\rho = 4 \csc \phi \sec \phi$ , spherical equation

$$= \frac{4}{\sin \phi \cos \theta}$$

$$\rho \sin \phi \cos \theta = 4$$

$$x = 4, \text{ rectangular equation}$$



59.  $\left(4, \frac{\pi}{4}, 0\right)$ , cylindrical

$$\rho = \sqrt{4^2 + 0^2} = 4$$

$$\theta = \frac{\pi}{4}$$

$$\phi = \arccos \frac{z}{\sqrt{r^2 + z^2}} = \arccos 0 = \frac{\pi}{2}$$

$\left(4, \frac{\pi}{4}, \frac{\pi}{2}\right)$ , spherical

60.  $\left(3, -\frac{\pi}{4}, 0\right)$ , cylindrical

$$\rho = \sqrt{3^2 + 0^2} = 3$$

$$\theta = -\frac{\pi}{4}$$

$$\phi = \arccos \frac{z}{\sqrt{r^2 + z^2}} = \arccos \frac{0}{9} = \frac{\pi}{2}$$

$\left(3, -\frac{\pi}{4}, \frac{\pi}{2}\right)$ , spherical

61.  $\left(6, \frac{\pi}{2}, -6\right)$ , cylindrical

$$\rho = \sqrt{6^2 + (-6)^2} = \sqrt{72} = 6\sqrt{2}$$

$$\theta = \frac{\pi}{2}$$

$$\phi = \arccos \frac{z}{\sqrt{r^2 + z^2}} = \arccos \left( \frac{-6}{6\sqrt{2}} \right) = \arccos \left( \frac{-1}{\sqrt{2}} \right) = \frac{3\pi}{4}$$

$$\left(6\sqrt{2}, \frac{\pi}{2}, \frac{3\pi}{4}\right)$$
, spherical

62.  $\left(-4, \frac{\pi}{3}, 4\right)$ , cylindrical

$$\rho = \sqrt{(-4)^2 + 4^2} = 4\sqrt{2}$$

$$\theta = \frac{\pi}{3}$$

$$\phi = \arccos \frac{z}{\sqrt{r^2 + z^2}} = \arccos \frac{1}{\sqrt{2}} = \frac{\pi}{4}$$

$$\left(4\sqrt{2}, \frac{\pi}{3}, \frac{\pi}{4}\right)$$
, spherical

63.  $(12, \pi, 5)$ , cylindrical

$$\rho = \sqrt{12^2 + 5^2} = 13$$

$$\theta = \pi$$

$$\phi = \arccos \frac{z}{\sqrt{r^2 + z^2}} = \arccos \frac{5}{13}$$

$$\left(13, \pi, \arccos \frac{5}{13}\right)$$
, spherical

64.  $\left(4, \frac{\pi}{2}, 3\right)$ , cylindrical

$$\rho = \sqrt{4^2 + 3^2} = 5$$

$$\theta = \frac{\pi}{2}$$

$$\phi = \arccos \frac{z}{\sqrt{r^2 + z^2}} = \arccos \frac{3}{5}$$

$$\left(5, \frac{\pi}{2}, \arccos \frac{3}{5}\right)$$
, spherical

65.  $\left(10, \frac{\pi}{6}, \frac{\pi}{2}\right)$ , spherical

$$r = 10 \sin \frac{\pi}{2} = 10$$

$$\theta = \frac{\pi}{6}$$

$$z = 10 \cos \frac{\pi}{2} = 0$$

$$\left(10, \frac{\pi}{6}, 0\right)$$
, cylindrical

66.  $\left(4, \frac{\pi}{18}, \frac{\pi}{2}\right)$ , spherical

$$r = 4 \sin \frac{\pi}{2} = 4$$

$$\theta = \frac{\pi}{18}$$

$$z = 4 \cos \frac{\pi}{2} = 0$$

$$\left(4, \frac{\pi}{18}, 0\right)$$
, cylindrical

67.  $\left(6, -\frac{\pi}{6}, \frac{\pi}{3}\right)$ , spherical

$$r = 6 \sin \frac{\pi}{3} = 3\sqrt{3}$$

$$\theta = -\frac{\pi}{6}$$

$$z = 6 \cos \frac{\pi}{3} = 3$$

$$\left(3\sqrt{3}, -\frac{\pi}{6}, 3\right)$$
, cylindrical

68.  $\left(5, -\frac{5\pi}{6}, \pi\right)$ , spherical

$$r = 5 \sin \pi = 0$$

$$\theta = -\frac{5\pi}{6}$$

$$z = 5 \cos \pi = -5$$

$$\left(0, -\frac{5\pi}{6}, -5\right)$$
, cylindrical

69.  $\left(8, \frac{7\pi}{6}, \frac{\pi}{6}\right)$ , spherical

$$r = 8 \sin \frac{\pi}{6} = 4$$

$$\theta = \frac{7\pi}{6}$$

$$z = 8 \cos \frac{\pi}{6} = \frac{8\sqrt{3}}{2}$$

$$\left(4, \frac{7\pi}{6}, 4\sqrt{3}\right)$$
, cylindrical

70.  $\left(7, \frac{\pi}{4}, \frac{3\pi}{4}\right)$ , spherical

$$r = 7 \sin \frac{3\pi}{4} = \frac{7\sqrt{2}}{2}$$

$$\theta = \frac{\pi}{4}$$

$$z = 7 \cos \frac{3\pi}{4} = -\frac{7\sqrt{2}}{2}$$

$$\left(\frac{7\sqrt{2}}{2}, \frac{\pi}{4}, -\frac{7\sqrt{2}}{2}\right)$$
, cylindrical

71.  $r = 5$

Cylinder

Matches graph (d)

72.  $\theta = \frac{\pi}{4}$

Plane

Matches graph (e)

73.  $\rho = 5$

Sphere

Matches graph (c)

74.  $\phi = \frac{\pi}{4}$

Cone

Matches graph (a)

75.  $r^2 = z, x^2 + y^2 = z$

Paraboloid

Matches graph (f)

76.  $\rho = 4 \sec \phi, z = \rho \cos \phi = 4$

Plane

Matches graph (b)

77.  $\theta = c$  is a half-plane because of the restriction  $r \geq 0$ .

78. (a) The surface is a cone. The equation is (i)

$$x^2 + y^2 = \frac{4}{9}z^2.$$

In cylindrical coordinates, the equation is

$$x^2 + y^2 = \frac{4}{9}z^2$$

$$r^2 = \frac{4}{9}z^2$$

$$r = \frac{2}{3}z.$$

(b) The surface is a hyperboloid of one sheet. The equation is (ii)  $x^2 + y^2 - z^2 = 2$ .

In cylindrical coordinates, the equation is

$$x^2 + y^2 - z^2 = 2$$

$$r^2 - z^2 = 2$$

$$r^2 = z^2 + 2.$$

79.  $x^2 + y^2 + z^2 = 27$

(a)  $r^2 + z^2 = 27$

(b)  $\rho^2 = 27 \Rightarrow \rho = 3\sqrt{3}$

80.  $4(x^2 + y^2) = z^2$

(a)  $4r^2 = z^2 \Rightarrow 2r = z$

(b)  $4(\rho^2 \sin^2 \phi \cos^2 \theta + \rho^2 \sin^2 \phi \sin^2 \theta) = \rho^2 \cos^2 \phi$

$$4 \sin^2 \phi = \cos^2 \phi,$$

$$\tan^2 \phi = \frac{1}{4},$$

$$\tan \phi = \frac{1}{2} \Rightarrow \phi = \arctan \frac{1}{2}$$

81.  $x^2 + y^2 + z^2 - 2z = 0$

(a)  $r^2 + z^2 - 2z = 0 \Rightarrow r^2 + (z - 1)^2 = 1$

(b)  $\rho^2 - 2\rho \cos \phi = 0$

$$\rho(\rho - 2 \cos \phi) = 0$$

$$\rho = 2 \cos \phi$$

82.  $x^2 + y^2 = z$

(a)  $r^2 = z$

(b)  $\rho^2 \sin^2 \phi = \rho \cos \phi$

$$\rho \sin^2 \phi = \cos \phi$$

$$\rho = \frac{\cos \phi}{\sin^2 \phi}$$

$$\rho = \csc \phi \cot \phi$$

83.  $x^2 + y^2 = 4y$

(a)  $r^2 = 4r \sin \theta, r = 4 \sin \theta$

(b)  $\rho^2 \sin^2 \phi = 4\rho \sin \phi \sin \theta$

$$\rho \sin \phi (\rho \sin \phi - 4 \sin \theta) = 0$$

$$\rho = \frac{4 \sin \theta}{\sin \phi}$$

$$\rho = 4 \sin \theta \csc \phi$$

84.  $x^2 + y^2 = 45$

(a)  $r^2 = 45$  or  $r = 3\sqrt{5}$

(b)  $\rho^2 \sin^2 \phi \cos^2 \theta + \rho^2 \sin^2 \phi \sin^2 \theta = 45$

$$\rho^2 \sin^2 \phi = 45$$

$$\rho = 3\sqrt{5} \csc \phi$$

85.  $x^2 - y^2 = 9$

(a)  $r^2 \cos^2 \theta - r^2 \sin^2 \theta = 9$

$$r^2 = \frac{9}{\cos^2 \theta - \sin^2 \theta}$$

(b)  $\rho^2 \sin^2 \phi \cos^2 \theta - \rho^2 \sin^2 \phi \sin^2 \theta = 9$

$$\rho^2 \sin^2 \phi = \frac{9}{\cos^2 \theta - \sin^2 \theta}$$


$$\rho^2 = \frac{9 \csc^2 \phi}{\cos^2 \theta - \sin^2 \theta}$$

86.  $y = 4$


(a)  $r \sin \theta = 4 \Rightarrow r = 4 \csc \theta$

(b)  $\rho \sin \phi \sin \theta = 4$ ,  
 $\rho = 4 \csc \phi \csc \theta$

87.  $0 \leq \theta \leq \frac{\pi}{2}$   
 $0 \leq r \leq 2$   
 $0 \leq z \leq 4$




88.  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$   
 $0 \leq r \leq 3$   
 $0 \leq z \leq r \cos \theta$



89.  $0 \leq \theta \leq 2\pi$

$$0 \leq r \leq a$$


$$r \leq z \leq a$$



90.  $0 \leq \theta \leq 2\pi$

$$2 \leq r \leq 4$$


$$z^2 \leq -r^2 + 6r - 8$$



91.  $0 \leq \theta \leq 2\pi$

$$0 \leq \phi \leq \frac{\pi}{6}$$


$$0 \leq \rho \leq a \sec \phi$$



92.  $0 \leq \theta \leq 2\pi$

$$\frac{\pi}{4} \leq \phi \leq \frac{\pi}{2}$$


$$0 \leq \rho \leq 1$$



93.  $0 \leq \theta \leq \frac{\pi}{2}$

$0 \leq \phi \leq \frac{\pi}{2}$


$0 \leq \rho \leq 2$



94.  $0 \leq \theta \leq \pi$

$0 \leq \phi \leq \frac{\pi}{2}$

$1 \leq \rho \leq 3$




95. Rectangular

$0 \leq x \leq 10$

$0 \leq y \leq 10$

$0 \leq z \leq 10$



96. Cylindrical:


$0.75 \leq r \leq 1.25$

$0 \leq z \leq 8$



97. Spherical

$4 \leq \rho \leq 6$




98. Cylindrical

$\frac{1}{2} \leq r \leq 3$

$0 \leq \theta \leq 2\pi$


$-\sqrt{9 - r^2} \leq z \leq \sqrt{9 - r^2}$



99. Cylindrical coordinates:

$r^2 + z^2 \leq 9,$

$r \leq 3 \cos \theta, 0 \leq \theta \leq \pi$




100. Spherical coordinates:

$\rho \geq 2$

$\rho \leq 3$

$0 \leq \phi \leq \frac{\pi}{4}$



- 101.** False.  $(r, \theta, z) = (0, 0, 1)$  and  $(r, \theta, z) = (0, \pi, 1)$

represent the same point  $(x, y, z) = (0, 0, 1)$ .

- 102.** True (except for the origin).

- 103.**  $z = \sin \theta, r = 1$

$$z = \sin \theta = \frac{y}{r} = \frac{y}{1} = y$$

The curve of intersection is the ellipse formed by the intersection of the plane  $z = y$  and the cylinder  $r = 1$ .

- 104.**  $\rho = 2 \sec \phi \Rightarrow \rho \cos \phi = 2 \Rightarrow z = 2$  plane

$\rho = 4$  sphere

The intersection of the plane and the sphere is a circle.

## Review Exercises for Chapter 11

**1.**  $P = (1, 2), Q = (4, 1), R = (5, 4)$

(a)  $\mathbf{u} = \overrightarrow{PQ} = \langle 4 - 1, 1 - 2 \rangle = \langle 3, -1 \rangle$

$\mathbf{v} = \overrightarrow{PR} = \langle 5 - 1, 4 - 2 \rangle = \langle 4, 2 \rangle$

(b)  $\mathbf{u} = 3\mathbf{i} - \mathbf{j}, \mathbf{v} = 4\mathbf{i} + 2\mathbf{j}$

(c)  $\|\mathbf{u}\| = \sqrt{3^2 + (-1)^2} = \sqrt{10} \quad \|\mathbf{v}\| = \sqrt{4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$

(d)  $-3\mathbf{u} + \mathbf{v} = -3\langle 3, -1 \rangle + \langle 4, 2 \rangle = \langle -5, 5 \rangle$

**2.**  $P = (-2, -1), Q = (5, -1), R = (2, 4)$

(a)  $\mathbf{u} = \overrightarrow{PQ} = \langle 5 - (-2), -1 - (-1) \rangle = \langle 7, 0 \rangle$

$\mathbf{v} = \overrightarrow{PR} = \langle 2 - (-2), 4 - (-1) \rangle = \langle 4, 5 \rangle$

(b)  $\mathbf{u} = 7\mathbf{i}, \mathbf{v} = 4\mathbf{i} + 5\mathbf{j}$

(c)  $\|\mathbf{u}\| = \sqrt{7^2 + 0^2} = \sqrt{49} = 7 \quad \|\mathbf{v}\| = \sqrt{4^2 + 5^2} = \sqrt{41}$

(d)  $-3\mathbf{u} + \mathbf{v} = -3\langle 7, 0 \rangle + \langle 4, 5 \rangle = \langle -17, 5 \rangle$

**3.**  $\mathbf{v} = \|\mathbf{v}\|(\cos \theta \mathbf{i} + \sin \theta \mathbf{j})$

$= 8(\cos 60^\circ \mathbf{i} + \sin 60^\circ \mathbf{j})$

$= 8\left(\frac{1}{2}\mathbf{i} + \frac{\sqrt{3}}{2}\mathbf{j}\right) = 4\mathbf{i} + 4\sqrt{3}\mathbf{j} = \langle 4, 4\sqrt{3} \rangle$

**4.**  $\mathbf{v} = \|\mathbf{v}\| \cos \theta \mathbf{i} + \|\mathbf{v}\| \sin \theta \mathbf{j}$

$= \frac{1}{2} \cos 225^\circ \mathbf{i} + \frac{1}{2} \sin 225^\circ \mathbf{j}$

$= -\frac{\sqrt{2}}{4}\mathbf{i} - \frac{\sqrt{2}}{4}\mathbf{j} = \left\langle -\frac{\sqrt{2}}{4}, -\frac{\sqrt{2}}{4} \right\rangle$

**5.**  $z = 0, y = 4, x = -5: (-5, 4, 0)$

- 6.**  $y = 3$  describes a plane parallel to the  $xz$ -plane and passing through  $(0, 3, 0)$ .

**7.**  $d = \sqrt{(-2 - 1)^2 + (3 - 6)^2 + (5 - 3)^2}$   
 $= \sqrt{9 + 9 + 4} = \sqrt{22}$

**8.**  $d = \sqrt{(4 - (-2))^2 + (-1 - 1)^2 + (-1 - (-5))^2}$   
 $= \sqrt{36 + 4 + 16} = \sqrt{56} = 2\sqrt{14}$

**9.**  $(x - 3)^2 + (y + 2)^2 + (z - 6)^2 = 4^2$   
 $(x - 3)^2 + (y + 2)^2 + (z - 6)^2 = 16$

**10.** Center:  $\left(\frac{0+4}{2}, \frac{0+6}{2}, \frac{4+0}{2}\right) = (2, 3, 2)$

Radius:


$$\sqrt{(2 - 0)^2 + (3 - 0)^2 + (2 - 4)^2} = \sqrt{4 + 9 + 4} = \sqrt{17}$$

$$(x - 2)^2 + (y - 3)^2 + (z - 2)^2 = 17$$

11.  $(x^2 - 4x + 4) + (y^2 - 6y + 9) + z^2 = -4 + 4 + 9$   
 $(x - 2)^2 + (y - 3)^2 + z^2 = 9$

Center:  $(2, 3, 0)$ 


Radius: 3




12.  $(x^2 - 10x + 25) + (y^2 + 6y + 9) + (z^2 - 4z + 4) = -34 + 25 + 9 + 4$   
 $(x - 5)^2 + (y + 3)^2 + (z - 2)^2 = 4$

Center:  $(5, -3, 2)$ 

Radius: 2




13. (a), (d)



(b)  $\mathbf{v} = \langle 4 - 2, 4 - (-1), -7 - 3 \rangle = \langle 2, 5, -10 \rangle$

(c)  $\mathbf{v} = 2\mathbf{i} + 5\mathbf{j} - 10\mathbf{k}$

14. (a), (d)



(b)  $\mathbf{v} = \langle 3 - 6, -3 - 2, 8 - 0 \rangle = \langle -3, -5, 8 \rangle$

(c)  $\mathbf{v} = -3\mathbf{i} - 5\mathbf{j} + 8\mathbf{k}$

15.  $z = -\mathbf{u} + 3\mathbf{v} + \frac{1}{2}\mathbf{w}$   
 $= -\langle 5, -2, 3 \rangle + 3\langle 0, 2, 1 \rangle + \frac{1}{2}\langle -6, -6, 2 \rangle$   
 $= \langle -5, 2, -3 \rangle + \langle 0, 6, 3 \rangle + \langle -3, -3, 1 \rangle$   
 $= \langle -8, 5, 1 \rangle$

16.  $\mathbf{u} - \mathbf{v} + \mathbf{w} - 2\mathbf{z} = 0$

$$\begin{aligned} z &= \frac{1}{2}(\mathbf{u} - \mathbf{v} + \mathbf{w}) \\ &= \frac{1}{2}(\langle 5, -2, 3 \rangle - \langle 0, 2, 1 \rangle + \langle -6, -6, 2 \rangle) \\ &= \frac{1}{2}\langle -1, -10, 4 \rangle \\ &= \left\langle -\frac{1}{2}, -5, 2 \right\rangle \end{aligned}$$

17.  $\mathbf{v} = \langle -1 - 3, 6 - 4, 9 + 1 \rangle = \langle -4, 2, 10 \rangle$

$\mathbf{w} = \langle 5 - 3, 3 - 4, -6 + 1 \rangle = \langle 2, -1, -5 \rangle$

Because  $-2\mathbf{w} = \mathbf{v}$ , the points lie in a straight line.

18.  $\mathbf{v} = \langle 8 - 5, -5 + 4, 5 - 7 \rangle = \langle 3, -1, -2 \rangle$

$\mathbf{w} = \langle 11 - 5, 6 + 4, 3 - 7 \rangle = \langle 6, 10, -4 \rangle$

Because  $\mathbf{v}$  and  $\mathbf{w}$  are not parallel, the points do not lie in a straight line.

19. Unit vector:  $\frac{\mathbf{u}}{\|\mathbf{u}\|} = \left\langle \frac{2}{\sqrt{38}}, \frac{3}{\sqrt{38}}, \frac{5}{\sqrt{38}} \right\rangle$

20.  $8 \frac{\langle 6, -3, 2 \rangle}{\sqrt{49}} = \frac{8}{7} \langle 6, -3, 2 \rangle = \left\langle \frac{48}{7}, -\frac{24}{7}, \frac{16}{7} \right\rangle$

21.  $P = \langle 5, 0, 0 \rangle, Q = \langle 4, 4, 0 \rangle, R = \langle 2, 0, 6 \rangle$

(a)  $\mathbf{u} = \overrightarrow{PQ} = \langle -1, 4, 0 \rangle$

$\mathbf{v} = \overrightarrow{PR} = \langle -3, 0, 6 \rangle$

(b)  $\mathbf{u} \cdot \mathbf{v} = (-1)(-3) + 4(0) + 0(6) = 3$

(c)  $\mathbf{v} \cdot \mathbf{v} = 9 + 36 = 45$

22.  $P = \langle 2, -1, 3 \rangle, Q = \langle 0, 5, 1 \rangle, R = \langle 5, 5, 0 \rangle$

(a)  $\mathbf{u} = \overrightarrow{PQ} = \langle -2, 6, -2 \rangle$

$\mathbf{v} = \overrightarrow{PR} = \langle 3, 6, -3 \rangle$

(b)  $\mathbf{u} \cdot \mathbf{v} = (-2)(3) + (6)(6) + (-2)(-3) = 36$

(c)  $\mathbf{v} \cdot \mathbf{v} = 9 + 36 + 9 = 54$

23.  $\mathbf{u} = 5\left(\cos \frac{3\pi}{4}\mathbf{i} + \sin \frac{3\pi}{4}\mathbf{j}\right) = \frac{5\sqrt{2}}{2}[-\mathbf{i} + \mathbf{j}]$

$$\mathbf{v} = 2\left(\cos \frac{2\pi}{3}\mathbf{i} + \sin \frac{2\pi}{3}\mathbf{j}\right) = -\mathbf{i} + \sqrt{3}\mathbf{j}$$

$$\mathbf{u} \cdot \mathbf{v} = \frac{5\sqrt{2}}{2}(1 + \sqrt{3})$$

$$\|\mathbf{u}\| = \sqrt{\frac{25}{2} + \frac{25}{2}} = 5 \quad \|\mathbf{v}\| = \sqrt{1 + 3} = 2$$

$$\cos \theta = \frac{|\mathbf{u} \cdot \mathbf{v}|}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{(5\sqrt{2}/2)(1 + \sqrt{3})}{5(2)} = \frac{\sqrt{2} + \sqrt{6}}{4}$$

$$(a) \theta = \arccos \frac{\sqrt{2} + \sqrt{6}}{4} = \frac{\pi}{12} \approx 0.262$$

$$(b) \theta \approx 15^\circ$$

24.  $\mathbf{u} = \langle 1, 0, -3 \rangle$

$$\mathbf{v} = (2, -2, 1)$$

$$\mathbf{u} \cdot \mathbf{v} = -1$$

$$\|\mathbf{u}\| = \sqrt{1 + 9} = \sqrt{10}$$

$$\|\mathbf{v}\| = \sqrt{4 + 4 + 1} = 3$$

$$\cos \theta = \frac{|\mathbf{u} \cdot \mathbf{v}|}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{1}{3\sqrt{10}}$$

$$(a) \theta = \arccos \left( \frac{1}{3\sqrt{10}} \right) \approx 1.465$$

$$(b) \theta = 83.9^\circ$$

25.  $\mathbf{u} = \langle 7, -2, 3 \rangle, \mathbf{v} = \langle -1, 4, 5 \rangle$

Because  $\mathbf{u} \cdot \mathbf{v} = 0$ , the vectors are orthogonal.

26.  $\mathbf{u} = \langle -3, 0, 9 \rangle = -3\langle 1, 0, -3 \rangle = -3\mathbf{v}$

The vectors are parallel.

27.  $\mathbf{u} = \langle 4, 2 \rangle, \mathbf{v} = \langle 3, 4 \rangle$

$$(a) \mathbf{w}_1 = \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v}$$

$$= \left( \frac{20}{25} \right) \langle 3, 4 \rangle$$

$$= \frac{4}{5} \langle 3, 4 \rangle = \left\langle \frac{12}{5}, \frac{16}{5} \right\rangle$$

$$(b) \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 4, 2 \rangle - \left\langle \frac{12}{5}, \frac{16}{5} \right\rangle = \left\langle \frac{8}{5}, -\frac{6}{5} \right\rangle$$

28.  $\mathbf{u} = \langle 1, -1, 1 \rangle, \mathbf{v} = \langle 2, 0, 2 \rangle$

$$(a) \mathbf{w}_1 = \text{proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2} \right) \mathbf{v}$$

$$= \frac{4}{8} \langle 2, 0, 2 \rangle = \langle 1, 0, 1 \rangle$$

$$(b) \mathbf{w}_2 = \mathbf{u} - \mathbf{w}_1 = \langle 1, -1, 1 \rangle - \langle 1, 0, 1 \rangle = \langle 0, -1, 0 \rangle$$

29. There are many correct answers.

For example:  $\mathbf{v} = \pm \langle 6, -5, 0 \rangle$ .

30.  $W = \mathbf{F} \cdot \overline{PQ} = \|\mathbf{F}\| \|\overline{PQ}\| \cos \theta = (75)(8) \cos 30^\circ$

$$= 300\sqrt{3} \text{ J}$$

31. (a)  $\mathbf{u} \times \mathbf{v} = \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & 3 & 6 \\ 5 & 2 & 1 \end{bmatrix} = -9\mathbf{i} + 26\mathbf{j} - 7\mathbf{k}$

$$(b) \mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = 9\mathbf{i} - 26\mathbf{j} + 7\mathbf{k}$$

$$(c) \mathbf{v} \times \mathbf{v} = \mathbf{0}$$

32. (a)  $\mathbf{u} \times \mathbf{v} = \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 2 & 1 \\ 1 & -3 & 4 \end{bmatrix} = 11\mathbf{i} + \mathbf{j} - 2\mathbf{k}$

$$(b) \mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = -11\mathbf{i} - \mathbf{j} + 2\mathbf{k}$$

$$(c) \mathbf{v} \times \mathbf{v} = \mathbf{0}$$

33.  $\mathbf{u} \times \mathbf{v} = \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -10 & 8 \\ 4 & 6 & -8 \end{bmatrix} = 32\mathbf{i} + 48\mathbf{j} + 52\mathbf{k}$

$$\|\mathbf{u} \times \mathbf{v}\| = \sqrt{6032} = 4\sqrt{377}$$

$$\text{Unit vector: } \frac{1}{\sqrt{377}} \langle 8, 12, 13 \rangle$$

34.  $\mathbf{u} = \langle 3, -1, 5 \rangle, \mathbf{v} = \langle 2, -4, 1 \rangle$

$$\mathbf{u} \times \mathbf{v} = \begin{bmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -1 & 5 \\ 2 & -4 & 1 \end{bmatrix} = 19\mathbf{i} + 7\mathbf{j} - 10\mathbf{k}$$

$$A = \|\mathbf{u} \times \mathbf{v}\| = \sqrt{19^2 + 7^2 + (-10)^2} = \sqrt{510}$$


35.  $\mathbf{F} = -40\mathbf{k}$  ( $25 \text{ cm} = \frac{1}{4} \text{ m}$ )

$$\overrightarrow{PQ} = \frac{1}{4}(\cos 60^\circ \mathbf{j} + \sin 60^\circ \mathbf{k}) = \frac{1}{8}\mathbf{j} + \frac{\sqrt{3}}{8}\mathbf{k}$$

The moment of  $\mathbf{F}$  about  $P$  is

$$M = \overrightarrow{PQ} \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & \frac{1}{8} & \frac{\sqrt{3}}{8} \\ 0 & 0 & -40 \end{vmatrix} = -5\mathbf{i}$$

Torque = 5 N-m



36.  $V = |\mathbf{u} \cdot (\mathbf{v} \cdot \mathbf{w})| = \begin{vmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & -1 & 2 \end{vmatrix} = 2(5) = 10$

37.  $\mathbf{v} = \langle 9 - 3, 11 - 0, 6 - 2 \rangle = \langle 6, 11, 4 \rangle$

(a) Parametric equations:

$$x = 3 + 6t, y = 11t, z = 2 + 4t$$

(b) Symmetric equations:  $\frac{x - 3}{6} = \frac{y}{11} = \frac{z - 2}{4}$

38.  $\mathbf{v} = \langle 8 + 1, 10 - 4, 5 - 3 \rangle = \langle 9, 6, 2 \rangle$

(a) Parametric equations:

$$x = -1 + 9t, y = 4 + 6t, z = 3 + 2t$$

(b) Symmetric equations:  $\frac{x + 1}{9} = \frac{y - 4}{6} = \frac{z - 3}{2}$

39.  $P = (-6, -8, 2)$

$$\mathbf{v} = \mathbf{j} = \langle 0, 1, 0 \rangle$$

$$x = -6, y = -8 + t, z = 2$$

40. Direction numbers: 1, 1, 1,  $\mathbf{v} = \langle 1, 1, 1 \rangle$

$$P(1, 2, 3)$$

$$x = 1 + t, y = 2 + t, z = 3 + t$$

41.  $P = (-3, -4, 2), Q = (-3, 4, 1), R = (1, 1, -2)$

$$\overrightarrow{PQ} = \langle 0, 8, -1 \rangle, \overrightarrow{PR} = [4, 5, -4]$$

$$\mathbf{n} = \overrightarrow{PQ} \times \overrightarrow{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 8 & -1 \\ 4 & 5 & -4 \end{vmatrix} = -27\mathbf{i} - 4\mathbf{j} - 32\mathbf{k}$$

$$-27(x + 3) - 4(y + 4) - 32(z - 2) = 0 \\ 27x + 4y + 32z = -33$$

42.  $\mathbf{n} = 3\mathbf{i} - \mathbf{j} + \mathbf{k}$

$$3(x + 2) - 1(y - 3) + 1(z - 1) = 0$$

$$3x - y + z + 8 = 0$$

43. The two lines are parallel as they have the same direction numbers,  $-2, 1, 1$ . Therefore, a vector parallel to the plane is  $\mathbf{v} = -2\mathbf{i} + \mathbf{j} + \mathbf{k}$ . A point on the first line is  $(1, 0, -1)$  and a point on the second line is  $(-1, 1, 2)$ .

The vector  $\mathbf{u} = 2\mathbf{i} - \mathbf{j} - 3\mathbf{k}$  connecting these two points is also parallel to the plane. Therefore, a normal to the plane is

$$\mathbf{v} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -2 & 1 & 1 \\ 2 & -1 & -3 \end{vmatrix} = -2\mathbf{i} - 4\mathbf{j} = -2(\mathbf{i} + 2\mathbf{j}).$$

Equation of the plane:  $(x - 1) + 2y = 0$

$$x + 2y = 1$$

44. Let  $\mathbf{v} = \langle 5 - 2, 1 + 2, 3 - 1 \rangle = \langle 3, 3, 2 \rangle$  be the direction vector for the line through the two points. Let  $\mathbf{n} = \langle 2, 1, -1 \rangle$  be the normal vector to the plane. Then

$$\mathbf{v} \times \mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & 3 & 2 \\ 2 & 1 & -1 \end{vmatrix} = \langle -5, 7, -3 \rangle$$

is the normal to the unknown plane.

$$-5(x - 5) + 7(y - 1) - 3(z - 3) = 0$$

$$-5x + 7y - 3z + 27 = 0$$

45.  $Q(1, 0, 2)$  point

$$2x - 3y + 6z = 6$$

A point  $P$  on the plane is  $(3, 0, 0)$ .

$$\overrightarrow{PQ} = \langle -2, 0, 2 \rangle$$

$\mathbf{n} = \langle 2, -3, 6 \rangle$  normal to plane

$$D = \frac{|\overrightarrow{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{8}{7}$$

46.  $Q(3, -2, 4)$  point

$$2x - 5y + z = 10$$

A point  $P$  on the plane is  $(5, 0, 0)$ .

$$\overline{PQ} = \langle -2, -2, 4 \rangle$$

 $\mathbf{n} = \langle 2, -5, 1 \rangle$  normal to plane

$$D = \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{10}{\sqrt{30}} = \frac{\sqrt{30}}{3}$$

47. The normal vectors to the planes are the same,

$$\mathbf{n} = \langle 5, -3, 1 \rangle.$$

Choose a point in the first plane  $P(0, 0, 2)$ . Choose a point in the second plane,  $Q(0, 0, -3)$ .

$$\overline{PQ} = \langle 0, 0, -5 \rangle$$

$$D = \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} = \frac{|-5|}{\sqrt{35}} = \frac{5}{\sqrt{35}} = \frac{\sqrt{35}}{7}$$

48.  $Q(-5, 1, 3)$  point

$$\mathbf{u} = \langle 1, -2, -1 \rangle$$
 direction vector

 $P(1, 3, 5)$  point on line





$$\overline{PQ} = \langle -6, -2, -2 \rangle$$

$$\overline{PQ} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -6 & -2 & -2 \\ 1 & -2 & -1 \end{vmatrix} = \langle -2, -8, 14 \rangle$$

$$D = \frac{\|\overline{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|} = \frac{\sqrt{264}}{\sqrt{6}} = 2\sqrt{11}$$

49.  $x + 2y + 3z = 6$ 

Plane


Intercepts:  $(6, 0, 0), (0, 3, 0), (0, 0, 2)$ ,50.  $y = z^2$ Because the  $x$ -coordinate is missing, you have a cylindrical surface with rulings parallel to the  $x$ -axis. The generating curve is a parabola in the  $yz$ -coordinate plane.51.  $y = \frac{1}{2}z$ Plane with rulings parallel to the  $x$ -axis.52.  $y = \cos z$ Because the  $x$ -coordinate is missing, you have a cylindrical surface with rulings parallel to the  $x$ -axis. The generating curve is  $y = \cos z$ .53.  $\frac{x^2}{16} + \frac{y^2}{9} + z^2 = 1$ 

Ellipsoid

$$xy\text{-trace: } \frac{x^2}{16} + \frac{y^2}{9} = 1$$

$$xz\text{-trace: } \frac{x^2}{16} + z^2 = 1$$

$$yz\text{-trace: } \frac{y^2}{9} + z^2 = 1$$



54.  $16x^2 + 16y^2 - 9z^2 = 0$


Cone

$xy$ -trace: point  $(0, 0, 0)$

$xz$ -trace:  $z = \pm \frac{4x}{3}$

$yz$ -trace:  $z = \pm \frac{4y}{3}$

$z = 4, x^2 + y^2 = 9$



55.  $\frac{x^2}{16} - \frac{y^2}{9} + z^2 = -1$


$\frac{y^2}{9} - \frac{x^2}{16} - z^2 = 1$

Hyperboloid of two sheets

$xy$ -trace:  $\frac{y^2}{9} - \frac{x^2}{16} = 1$

$xz$ -trace: None

$yz$ -trace:  $\frac{y^2}{9} - z^2 = 1$




56.  $\frac{x^2}{25} + \frac{y^2}{4} - \frac{z^2}{100} = 1$

Hyperboloid of one sheet

$xy$ -trace:  $\frac{x^2}{25} + \frac{y^2}{4} = 1$


$xz$ -trace:  $\frac{x^2}{25} - \frac{z^2}{100} = 1$

$yz$ -trace:  $\frac{y^2}{4} - \frac{z^2}{100} = 1$




57.  $x^2 + z^2 = 4$ .

Cylinder of radius 2 about  $y$ -axis



58.  $y^2 + z^2 = 16$ .

Cylinder of radius 4 about  $x$ -axis



59.  $z^2 = 2y$  revolved about  $y$ -axis

$z = \pm \sqrt{2y}$

$x^2 + z^2 = [r(y)]^2 = 2y$

$x^2 + z^2 = 2y$

60.  $2x + 3z = 1$  revolved about the  $x$ -axis

$z = \frac{1 - 2x}{3}$

$y^2 + z^2 = [r(x)]^2 = \left(\frac{1 - 2x}{3}\right)^2$ , Cone

61.  $(-\sqrt{3}, 3, -5)$ , rectangular

(a)  $r = \sqrt{(-\sqrt{3})^2 + 3^2} = \sqrt{12} = 2\sqrt{3}$

$\tan \theta = \frac{-3}{\sqrt{3}} \Rightarrow \theta = -\frac{\pi}{3}$

$z = -5$

$\left(2\sqrt{3}, -\frac{\pi}{3}, -5\right)$ , cylindrical

(b)  $\rho = \sqrt{(-\sqrt{3})^2 + 3^2 + (-5)^2} = \sqrt{37}$

$\tan \theta = -\frac{3}{\sqrt{3}} \Rightarrow \theta = -\frac{\pi}{3}$

$\phi = \arccos \frac{z}{\rho} = \arccos \left( \frac{-5}{\sqrt{37}} \right)$

$\left(\sqrt{37}, -\frac{\pi}{3}, \arccos \left( -\frac{5\sqrt{37}}{37} \right) \right)$ , spherical

62.  $(8, 8, 1)$ , rectangular

(a)  $r = \sqrt{8^2 + 8^2} = 8\sqrt{2}$

$\tan \theta = \frac{8}{8} = 1 \Rightarrow \theta = \frac{\pi}{4}$

$z = 1$

$\left(8\sqrt{2}, \frac{\pi}{4}, 1\right)$ , cylindrical

(b)  $\rho = \sqrt{8^2 + 8^2 + 1^2} = \sqrt{129}$

$\tan \theta = \frac{8}{8} = 1 \Rightarrow \theta = \frac{\pi}{4}$

$\phi = \arccos \frac{z}{\rho} = \arccos \frac{1}{\sqrt{129}}$

$\left(\sqrt{129}, \frac{\pi}{4}, \arccos \frac{1}{\sqrt{129}}\right)$ , spherical

63.  $(5, \pi, 1)$ , cylindrical

$x = r \cos \theta = 5 \cos \pi = -5$

$y = r \sin \theta = 5 \sin \pi = 0$

$z = 1$

$(-5, 0, 1)$ , rectangular

66.  $\left(8, -\frac{\pi}{6}, \frac{\pi}{3}\right)$ , spherical

$x = \rho \sin \phi \cos \theta = 8 \sin \frac{\pi}{3} \cos \left(-\frac{\pi}{6}\right) = 8 \left(\frac{\sqrt{3}}{2}\right) \left(\frac{\sqrt{3}}{2}\right) = 6$

$y = \rho \sin \phi \sin \theta = 8 \sin \frac{\pi}{3} \sin \left(-\frac{\pi}{6}\right) = 8 \left(\frac{\sqrt{3}}{2}\right) \left(-\frac{1}{2}\right) = -2\sqrt{3}$

$z = \rho \cos \phi = 8 \cos \frac{\pi}{3} = 4$

$(6, -2\sqrt{3}, 4)$ , rectangular

67.  $x^2 - y^2 = 2z$ 

(a) Cylindrical:

$r^2 \cos^2 \theta - r^2 \sin^2 \theta = 2z \Rightarrow r^2 \cos 2\theta = 2z$

(b) Spherical:

$\rho^2 \sin^2 \phi \cos^2 \theta - \rho^2 \sin^2 \phi \sin^2 \theta = 2\rho \cos \phi$

$\rho \sin^2 \phi \cos 2\theta - 2 \cos \phi = 0$

$\rho = 2 \sec 2\theta \cos \phi \csc^2 \phi$

68.  $x^2 + y^2 + z^2 = 16$ (a) Cylindrical:  $r^2 + z^2 = 16$ (b) Spherical:  $\rho = 4$ 64.  $\left(-2, \frac{\pi}{3}, 3\right)$ , cylindrical

$x = r \cos \theta = -2 \cos \frac{\pi}{3} = -1$

$y = r \sin \theta = -2 \sin \frac{\pi}{3} = -\sqrt{3}$

$z = 3$

$(-1, -\sqrt{3}, 3)$ , rectangular


65.  $\left(4, \pi, \frac{\pi}{4}\right)$ , spherical

$x = \rho \sin \phi \cos \theta = 4 \sin \frac{\pi}{4} \cos \pi = -2\sqrt{2}$

$y = \rho \sin \phi \sin \theta = 4 \sin \frac{\pi}{4} \sin \pi = 0$

$z = \rho \cos \phi = 4 \cos \frac{\pi}{4} = 2\sqrt{2}$


$(-2\sqrt{2}, 0, 2\sqrt{2})$ , rectangular

69.  $z = r^2 \sin^2 \theta + 3r \cos \theta$ , cylindrical equation $z = y^2 + 3x$ , rectangular equation

70.  $r = -5z$ , cylindrical equation

$$\sqrt{x^2 + y^2} = -5z$$


$x^2 + y^2 - 25z^2 = 0$ , rectangular equation



72.  $\rho = 9 \sec \theta$ , spherical equation

$$\rho \cos \theta = 9$$

$z = 9$ , rectangular equation



71.  $\phi = \frac{\pi}{4}$ , spherical equation


$$\phi = \arccos \frac{z}{\sqrt{x^2 + y^2 + z^2}} = \frac{\pi}{4}$$

$$\frac{z}{\sqrt{x^2 + y^2 + z^2}} = \cos \frac{\pi}{4} = \frac{\sqrt{2}}{2}$$

$$z^2 = \frac{1}{2}(x^2 + y^2 + z^2)$$

$$2z^2 = x^2 + y^2 + z^2$$

$x^2 + y^2 - z^2 = 0$ , rectangular equation



## Problem Solving for Chapter 11

1.  $\mathbf{a} + \mathbf{b} + \mathbf{c} = \mathbf{0}$

$$\mathbf{b} \times (\mathbf{a} + \mathbf{b} + \mathbf{c}) = \mathbf{0}$$

$$(\mathbf{b} \times \mathbf{a}) + (\mathbf{b} \times \mathbf{c}) = \mathbf{0}$$


$$\|\mathbf{a} \times \mathbf{b}\| = \|\mathbf{b} \times \mathbf{c}\|$$

$$\|\mathbf{b} \times \mathbf{c}\| = \|\mathbf{b}\| \|\mathbf{c}\| \sin A$$

$$\|\mathbf{a} \times \mathbf{b}\| = \|\mathbf{a}\| \|\mathbf{b}\| \sin C$$

Then,


$$\begin{aligned} \frac{\sin A}{\|\mathbf{a}\|} &= \frac{\|\mathbf{b} \times \mathbf{c}\|}{\|\mathbf{a}\| \|\mathbf{b}\| \|\mathbf{c}\|} \\ &= \frac{\|\mathbf{a} \times \mathbf{b}\|}{\|\mathbf{a}\| \|\mathbf{b}\| \|\mathbf{c}\|} \\ &= \frac{\sin C}{\|\mathbf{c}\|}. \end{aligned}$$



The other case,  $\frac{\sin A}{\|\mathbf{a}\|} = \frac{\sin B}{\|\mathbf{b}\|}$  is similar.

2.  $f(x) = \int_0^x \sqrt{t^4 + 1} dt$

(a)



(b)  $f'(x) = \sqrt{x^4 + 1}$

$$f'(0) = 1 = \tan \theta$$

$$\theta = \frac{\pi}{4}$$

$$\mathbf{u} = \frac{1}{\sqrt{2}}(\mathbf{i} + \mathbf{j}) = \left\langle \frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right\rangle$$

(c)  $\pm \left\langle \frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2} \right\rangle$

(d) The line is  $y = x$ :  $x = t$ ,  $y = t$ .


3. Label the figure as indicated.

From the figure, you see that

$$\overline{SP} = \frac{1}{2}\mathbf{a} - \frac{1}{2}\mathbf{b} = \overline{RQ} \text{ and } \overline{SR} = \frac{1}{2}\mathbf{a} + \frac{1}{2}\mathbf{b} = \overline{PQ}.$$

Because  $\overline{SP} = \overline{RQ}$  and  $\overline{SR} = \overline{PQ}$ ,

$PSRQ$  is a parallelogram.




4. Label the figure as indicated.

$$\overline{PR} = \mathbf{a} + \mathbf{b}$$

$$\overline{SQ} = \mathbf{b} - \mathbf{a}$$


$$(\mathbf{a} + \mathbf{b}) \cdot (\mathbf{b} - \mathbf{a}) = \|\mathbf{b}\|^2 - \|\mathbf{a}\|^2 = 0, \text{ because}$$

$\|\mathbf{a}\| = \|\mathbf{b}\|$  in a rhombus.



5. (a)  $\mathbf{u} = \langle 0, 1, 1 \rangle$  is the direction vector of the line determined by  $P_1$  and  $P_2$ .

$$\begin{aligned} D &= \frac{\|\overline{P_1Q} \times \mathbf{u}\|}{\|\mathbf{u}\|} \\ &= \frac{\|\langle 2, 0, -1 \rangle \times \langle 0, 1, 1 \rangle\|}{\sqrt{2}} \\ &= \frac{\|\langle 1, -2, 2 \rangle\|}{\sqrt{2}} = \frac{3}{\sqrt{2}} = \frac{3\sqrt{2}}{2} \end{aligned}$$




- (b) The shortest distance to the line segment is  $\|P_1Q\| = \|\langle 2, 0, -1 \rangle\| = \sqrt{5}$ .

$$6. (\mathbf{n} + \overrightarrow{PP_0}) \perp (\mathbf{n} - \overrightarrow{PP_0})$$

Figure is a square.

So,  $\|\overrightarrow{PP_0}\| = \|\mathbf{n}\|$  and the points  $P$  form a circle of radius  $\|\mathbf{n}\|$  in the plane with center at  $P_0$ .



$$7. (a) V = \pi \int_0^1 (\sqrt{z})^2 dz = \left[ \pi \frac{z^2}{2} \right]_0^1 = \frac{1}{2}\pi$$

$$\text{Note: } \frac{1}{2}(\text{base})(\text{altitude}) = \frac{1}{2}\pi(1) = \frac{1}{2}\pi$$

$$(b) \frac{x^2}{a^2} + \frac{y^2}{b^2} = z: \text{(slice at } z = c)$$

$$\frac{x^2}{(\sqrt{ca})^2} + \frac{y^2}{(\sqrt{cb})^2} = 1$$

At  $z = c$ , figure is ellipse of area

$$\pi(\sqrt{ca})(\sqrt{cb}) = \pi abc.$$

$$V = \int_0^k \pi abc \cdot dc = \left[ \frac{\pi abc^2}{2} \right]_0^k = \frac{\pi abk^2}{2}$$

$$(c) V = \frac{1}{2}(\pi abk)k = \frac{1}{2}(\text{area of base})(\text{height})$$

$$8. (a) V = 2 \int_0^r \pi(r^2 - x^2) dx = 2\pi \left[ r^2x - \frac{x^3}{3} \right]_0^r = \frac{4}{3}\pi r^3$$

- (b) At height  $z = d > 0$ ,

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{d^2}{c^2} = 1$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 - \frac{d^2}{c^2} = \frac{c^2 - d^2}{c^2}$$

$$\frac{x^2}{a^2(c^2 - d^2)} + \frac{y^2}{b^2(c^2 - d^2)} = 1.$$

$$\text{Area} = \pi \sqrt{\left( \frac{a^2(c^2 - d^2)}{c^2} \right) \left( \frac{b^2(c^2 - d^2)}{c^2} \right)}$$

$$= \frac{\pi ab}{c^2}(c^2 - d^2)$$

$$V = 2 \int_0^c \frac{\pi ab}{c^2}(c^2 - d^2) dd$$

$$= \frac{2\pi ab}{c^2} \left[ c^2 d - \frac{d^3}{3} \right]_0^c = \frac{4}{3}\pi abc$$

9. From Exercise 54, Section 11.4,

$$(\mathbf{u} \times \mathbf{v}) \times (\mathbf{w} \times \mathbf{z}) = [(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{z}] \mathbf{w} - [(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}] \mathbf{z}.$$

10.  $x = -t + 3, y = \frac{1}{2}t + 1, z = 2t - 1; Q = (4, 3, s)$


(a)  $\mathbf{u} = \langle -2, 1, 4 \rangle$  direction vector for line

$P = (3, 1, -1)$  point on line

$$\overline{PQ} = \langle 1, 2, s + 1 \rangle$$

$$\begin{aligned}\overline{PQ} \times \mathbf{u} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & s+1 \\ -2 & 1 & 4 \end{vmatrix} \\ &= (7-s)\mathbf{i} + (-6-2s)\mathbf{j} + 5\mathbf{k} \\ D &= \frac{\|\overline{PQ} \times \mathbf{u}\|}{\|\mathbf{u}\|} = \frac{\sqrt{(7-s)^2 + (-6-2s)^2 + 25}}{\sqrt{21}}\end{aligned}$$

(b)



The minimum is  $D \approx 2.2361$  at  $s = -1$ .


(c) Yes, there are slant asymptotes. Using  $s = x$ , you have

$$D(s) = \frac{1}{\sqrt{21}} \sqrt{5x^2 + 10x + 110} = \frac{\sqrt{5}}{\sqrt{21}} \sqrt{x^2 + 2x + 22} = \frac{\sqrt{5}}{\sqrt{21}} \sqrt{(x+1)^2 + 21} \rightarrow \pm \sqrt{\frac{5}{21}}(x+1)$$

$$y = \pm \frac{\sqrt{105}}{21}(s+1) \text{ slant asymptotes.}$$


11. (a)  $\rho = 2 \sin \phi$

Torus




(b)  $\rho = 2 \cos \phi$

Sphere



12. (a)  $r = 2 \cos \theta$


Cylinder



(b)  $z = r^2 \cos 2\theta$

$$\begin{aligned}
 &= r^2(\cos^2 \theta - \sin^2 \theta) \\
 &= (r \cos \theta)^2 - (r \sin \theta)^2 \\
 &= x^2 - y^2
 \end{aligned}$$

Hyperbolic paraboloid



13. (a)  $\mathbf{u} = \|\mathbf{u}\|(\cos 0 \mathbf{i} + \sin 0 \mathbf{j}) = \|\mathbf{u}\|\mathbf{i}$

Downward force  $\mathbf{w} = -3\mathbf{j}$ 

$\mathbf{T} = \|\mathbf{T}\|(\cos(90^\circ + \theta)\mathbf{i} + \sin(90^\circ + \theta)\mathbf{j})$

$= \|\mathbf{T}\|(-\sin \theta \mathbf{i} + \cos \theta \mathbf{j})$

$\mathbf{0} = \mathbf{u} + \mathbf{w} + \mathbf{T} = \|\mathbf{u}\|\mathbf{i} - 3\mathbf{j} + \|\mathbf{T}\|(-\sin \theta \mathbf{i} + \cos \theta \mathbf{j})$

$\|\mathbf{u}\| = \sin \theta \|\mathbf{T}\|$


$3 = \cos \theta \|\mathbf{T}\|$

If  $\theta = 30^\circ$ ,  $\|\mathbf{u}\| = (1/2)\|\mathbf{T}\|$  and  $3 = (\sqrt{3}/2)\|\mathbf{T}\| \Rightarrow \|\mathbf{T}\| = 2\sqrt{3} \approx 3.4641$  N and  $\|\mathbf{u}\| = \frac{1}{2}(2\sqrt{3}) \approx 1.7321$  N

(b) From part (a),  $\|\mathbf{u}\| = 3 \tan \theta$  and  $\|\mathbf{T}\| = 3 \sec \theta$ .Domain:  $0^\circ \leq \theta < 90^\circ$ 

$\theta$	$0^\circ$	$10^\circ$	$20^\circ$	$30^\circ$	$40^\circ$	$50^\circ$	$60^\circ$
$\mathbf{T}$	3	3.0463	3.1925	3.4641	3.9162	4.6672	6
$\ \mathbf{u}\ $	0	0.5290	1.0919	1.7321	2.5173	3.5753	5.1962

(d)



(e) Both are increasing functions.

(f)  $\lim_{\theta \rightarrow \pi/2^-} T = \infty$  and  $\lim_{\theta \rightarrow \pi/2^-} \|\mathbf{u}\| = \infty$ .

Yes. As  $\theta$  increases, both  $T$  and  $\|\mathbf{u}\|$  increase.

14. (a) The tension  $T$  is the same in each tow line.

$$\begin{aligned} 6000\mathbf{i} &= T(\cos 20^\circ + \cos(-20^\circ))\mathbf{i} + T(\sin 20^\circ + \sin(-20^\circ))\mathbf{j} \\ &= 2T\cos 20^\circ \mathbf{i} \\ \Rightarrow T &= \frac{6000}{2\cos 20^\circ} \approx 3192.5 \text{ N} \end{aligned}$$

- (b) As in part (a),  $6000\mathbf{i} = 2T\cos \theta$


$$\Rightarrow T = \frac{3000}{\cos \theta}$$

Domain:  $0^\circ < \theta < 90^\circ$

(c)

$\theta$	$10^\circ$	$20^\circ$	$30^\circ$	$40^\circ$	$50^\circ$	$60^\circ$
$T$	3046.3	3192.5	3464.1	3916.2	4667.2	6000.0

(d)



- (e) As  $\theta$  increases, there is less force applied in the direction of motion.

15. Let  $\theta = \alpha - \beta$ , the angle between  $\mathbf{u}$  and  $\mathbf{v}$ . Then

$$\sin(\alpha - \beta) = \frac{\|\mathbf{u} \times \mathbf{v}\|}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{\|\mathbf{v} \times \mathbf{u}\|}{\|\mathbf{u}\| \|\mathbf{v}\|}.$$

For  $\mathbf{u} = \langle \cos \alpha, \sin \alpha, 0 \rangle$  and  $\mathbf{v} = \langle \cos \beta, \sin \beta, 0 \rangle$ ,  $\|\mathbf{u}\| = \|\mathbf{v}\| = 1$  and

$$\mathbf{v} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \cos \beta & \sin \beta & 0 \\ \cos \alpha & \sin \alpha & 0 \end{vmatrix} = (\sin \alpha \cos \beta - \cos \alpha \sin \beta)\mathbf{k}.$$

So,  $\sin(\alpha - \beta) = \|\mathbf{v} \times \mathbf{u}\| = \sin \alpha \cos \beta - \cos \alpha \sin \beta$ .

16. (a) Los Angeles:  $(6371, -118.24^\circ, 55.95^\circ)$

Rio de Janeiro:  $(6371, -43.23^\circ, 112.90^\circ)$

- (b) Los Angeles:  $x = 6371 \sin(55.95^\circ) \cos(-118.24^\circ)$

Rio de Janeiro:  $x = 6371 \sin(112.90^\circ) \cos(-43.23^\circ)$

$$y = 6371 \sin(55.95^\circ) \sin(-118.24^\circ)$$

$$y = 6371 \sin(112.90^\circ) \sin(-43.23^\circ)$$

$$z = 6371 \cos(55.95^\circ)$$

$$z = 6371 \cos(112.90^\circ)$$

$$(x, y, z) \approx (-2497.7, -4650.4, 3567.2)$$

$$(x, y, z) \approx (4276.1, -4019.8, -2479.1)$$

$$(c) \cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{(-2497.7)(4276.1) + (-4650.4)(-4019.8) + (3567.2)(-2479.1)}{(6371)(6371)} \approx -0.02045$$

$$\theta \approx 91.17^\circ \text{ or } 1.59 \text{ radians}$$

$$(d) s = r\theta = 6371(1.59) \approx 10,130 \text{ km}$$

(e) For Boston and Honolulu:

a. Boston:  $(6371, -71.06^\circ, 47.64^\circ)$

Honolulu:  $(6371, -157.86^\circ, 68.69^\circ)$

b. Boston:  $x = 6371 \sin 47.64^\circ \cos(-71.06^\circ)$

Honolulu:  $x = 6371 \sin 68.69^\circ \cos(-157.86^\circ)$

$$y = 6371 \sin 47.64^\circ \sin(-71.06^\circ)$$

$$y = 6371 \sin 68.69^\circ \sin(-157.86^\circ)$$

$$z = 6371 \cos 47.64^\circ$$

$$z = 6371 \cos 68.69^\circ$$

$$(1528.0, -4452.8, 4292.7)$$

$$(-5497.8, -2236.9, 2315.3)$$

c.  $\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} = \frac{(1528.0)(-5497.8) + (-4452.8)(-2236.9) + (4292.7)(2315.3)}{(6371)(6371)} \approx 0.28329$

$$\theta \approx 73.54^\circ \text{ or } 1.28 \text{ radians}$$

d.  $s = r\theta = 6371(1.28) \approx 8155 \text{ km}$

17. From Theorem 11.13 and Theorem 11.7 (6) you have

$$\begin{aligned} D &= \frac{|\overline{PQ} \cdot \mathbf{n}|}{\|\mathbf{n}\|} \\ &= \frac{|\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})|}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{|(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}|}{\|\mathbf{u} \times \mathbf{v}\|} = \frac{|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|}{\|\mathbf{u} \times \mathbf{v}\|}. \end{aligned}$$

18. Assume one of  $a, b, c$ , is not zero, say  $a$ . Choose a point in the first plane such as  $(-d_1/a, 0, 0)$ . The distance between this point and the second plane is

$$\begin{aligned} D &= \frac{|a(-d_1/a) + b(0) + c(0) + d_2|}{\sqrt{a^2 + b^2 + c^2}} \\ &= \frac{|-d_1 + d_2|}{\sqrt{a^2 + b^2 + c^2}} = \frac{|d_1 - d_2|}{\sqrt{a^2 + b^2 + c^2}}. \end{aligned}$$

19.  $x^2 + y^2 = 1$  cylinder

$z = 2y$  plane

Introduce a coordinate system in the plane  $z = 2y$ .

The new  $u$ -axis is the original  $x$ -axis.

The new  $v$ -axis is the line  $z = 2y, x = 0$ .

Then the intersection of the cylinder and plane satisfies the equation of an ellipse:

$$x^2 + y^2 = 1$$

$$x^2 + \left(\frac{z}{2}\right)^2 = 1$$

$$x^2 + \frac{z^2}{4} = 1 \quad \text{ellipse}$$

