

Exam

Name _____

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the average rate of change of the function over the given interval.

1) $y = x^2 + 4x$, $[4, 7]$

A) 15

B) $\frac{45}{7}$

C) 11

D) $\frac{77}{3}$

Answer: A

2) $y = 5x^3 - 3x^2 - 8$, $[-8, -5]$

A) 684

B) - 236

C) $\frac{708}{5}$

D) $-\frac{2052}{5}$

Answer: A

3) $y = \sqrt{2x}$, $[2, 8]$

A) 7

B) $\frac{1}{3}$

C) $-\frac{3}{10}$

D) 2

Answer: B

4) $y = \frac{3}{x-2}$, $[4, 7]$

A) $\frac{1}{3}$

B) $-\frac{3}{10}$

C) 7

D) 2

Answer: B

5) $y = 4x^2$, $\left[0, \frac{7}{4}\right]$

A) 2

B) $-\frac{3}{10}$

C) 7

D) $\frac{1}{3}$

Answer: C

6) $y = -3x^2 - x$, $[5, 6]$

A) $\frac{1}{2}$

B) -2

C) -34

D) $-\frac{1}{6}$

Answer: C

7) $h(t) = \sin(4t)$, $\left[0, \frac{\pi}{8}\right]$

A) $\frac{4}{\pi}$

B) $-\frac{8}{\pi}$

C) $\frac{\pi}{8}$

D) $\frac{8}{\pi}$

Answer: D

8) $g(t) = 4 + \tan t, \left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$

A) $-\frac{4}{\pi}$

B) $\frac{4}{\pi}$

C) $-\frac{3}{2}$

D) 0

Answer: B

Find the slope of the curve at the given point P and an equation of the tangent line at P.

9) $y = x^2 + 5x, P(4, 36)$

A) slope is $\frac{1}{20}; y = \frac{x}{20} + \frac{1}{5}$

B) slope is $-\frac{4}{25}; y = -\frac{4x}{25} + \frac{8}{5}$

C) slope is -39; $y = -39x - 80$

D) slope is 13; $y = 13x - 16$

Answer: D

10) $y = x^2 + 11x - 15, P(1, -3)$

A) slope is $-\frac{4}{25}; y = -\frac{4x}{25} + \frac{8}{5}$

B) slope is -39; $y = -39x - 80$

C) slope is $\frac{1}{20}; y = \frac{x}{20} + \frac{1}{5}$

D) slope is 13; $y = 13x - 16$

Answer: D

11) $y = x^3 - 9x, P(1, -8)$

A) slope is -6; $y = -6x$

B) slope is 3; $y = 3x - 11$

C) slope is 3; $y = 3x - 7$

D) slope is -6; $y = -6x - 2$

Answer: D

12) $y = x^3 - 2x^2 + 4, P(3, 13)$

A) slope is 1; $y = x - 32$

B) slope is 15; $y = 15x + 13$

C) slope is 15; $y = 15x - 32$

D) slope is 0; $y = -32$

Answer: C

13) $y = 4 - x^3, (-1, 5)$

A) slope is -1; $y = -x + 2$

B) slope is 0; $y = 2$


C) slope is -3; $y = -3x + 2$

D) slope is 3; $y = 3x + 2$

Answer: C

Use the slopes of UQ, UR, US, and UT to estimate the rate of change of y at the specified value of x .

14) $x = 5$



A) 0


B) 1

C) 5

D) 2

Answer: D

15) $x = 3$



A) 2


B) 0

C) 6

D) 4

Answer: D

16) $x = 5$



A) $\frac{25}{4}$


B) 0

C) $\frac{5}{2}$

D) $\frac{5}{4}$

Answer: C

17) $x = 2$



A) 0


B) 6

C) 4

D) 3

Answer: B

18) $x = 2.5$



A) 1.25

B) 3.75

C) 7.5

D) 0

Answer: B

Use the table to estimate the rate of change of y at the specified value of x .

19) $x = 1$.

x	y
0	0
0.2	0.02
0.4	0.08
0.6	0.18
0.8	0.32
1.0	0.5
1.2	0.72
1.4	0.98

A) 0.5

B) 1

C) 2

D) 1.5

Answer: B

20) $x = 1$.

x	y
0	0
0.2	0.01
0.4	0.04
0.6	0.09
0.8	0.16
1.0	0.25
1.2	0.36
1.4	0.49

A) 2

B) 1

C) 0.5

D) 1.5

Answer: C

21) $x = 1$.

x	y
0	0
0.2	0.12
0.4	0.48
0.6	1.08
0.8	1.92
1.0	3
1.2	4.32
1.4	5.88

A) 8

B) 6

C) 2

D) 4

Answer: B

22) $x = 2$.

x	y
0	10
0.5	38
1.0	58
1.5	70
2.0	74
2.5	70
3.0	58
3.5	38
4.0	10

A) 0

B) 8

C) -8

D) 4

Answer: A

23) $x = 1$.

x	y
0.900	-0.05263
0.990	-0.00503
0.999	-0.0005
1.000	0.0000
1.001	0.0005
1.010	0.00498
1.100	0.04762

A) 0

B) 1

C) 0.5

D) -0.5


Answer: C

Solve the problem.


24) When exposed to ethylene gas, green bananas will ripen at an accelerated rate. The number of days for ripening becomes shorter for longer exposure times. Assume that the table below gives average ripening times of bananas for several different ethylene exposure times:

Exposure time (minutes)	Ripening Time (days)
10	4.2
15	3.5
20	2.6
25	2.1
30	1.1

Plot the data and then find a line approximating the data. With the aid of this line, find the limit of the average ripening time as the exposure time to ethylene approaches 0. Round your answer to the nearest tenth.




A)





5.8 days

B)



2.6 days



0.1 day


37.5 minutes

Answer: A


- 25) When exposed to ethylene gas, green bananas will ripen at an accelerated rate. The number of days for ripening becomes shorter for longer exposure times. Assume that the table below gives average ripening times of bananas for several different ethylene exposure times.

Exposure time (minutes)	Ripening Time (days)
10	4.3
15	3.2
20	2.7
25	2.1
30	1.3


Plot the data and then find a line approximating the data. With the aid of this line, determine the rate of change of ripening time with respect to exposure time. Round your answer to two significant digits.



A)




B)




-6.7 days per minute

C)



5.6 days

D)




38 minutes

-0.14 day per minute

Answer: D

26) The graph below shows the number of tuberculosis deaths in the United States from 1989 to 1998.




Estimate the average rate of change in tuberculosis deaths from 1993 to 1995.

- A) About -150 deaths per year
- B) About -300 deaths per year
- C) About -1 deaths per year
- D) About -80 deaths per year

Answer: A

Use the graph to evaluate the limit.


27) $\lim_{x \rightarrow 1} f(x)$



- A) $\frac{3}{4}$
- B) ∞
- C) -1
- D) $-\frac{3}{4}$

Answer: A

28) $\lim_{x \rightarrow 0} f(x)$



A) 2


B) -2

C) does not exist

D) 0

Answer: C

29) $\lim_{x \rightarrow 0} f(x)$



A) -3


B) 0

C) does not exist

D) 3

Answer: B

30) $\lim_{x \rightarrow 0} f(x)$



A) 6


B) -1

C) does not exist

D) 0

Answer: D

31) $\lim_{x \rightarrow 0} f(x)$



A) does not exist


B) ∞

C) 1

D) -1

Answer: A

32) $\lim_{x \rightarrow 0} f(x)$



A) 1


B) does not exist

C) ∞

D) -1

Answer: B

33) $\lim_{x \rightarrow 0} f(x)$



A) -2


B) 0

C) 2

D) does not exist

Answer: A

34) $\lim_{x \rightarrow 0} f(x)$



A) 1


B) -2

C) 0

D) does not exist

Answer: B

35) $\lim_{x \rightarrow 0} f(x)$



A) 2


B) -1

C) does not exist

D) -2

Answer: D

36) $\lim_{x \rightarrow 0} f(x)$



A) 0

B) -1

C) does not exist

D) 1

Answer: D

Find the limit.

37) $\lim_{x \rightarrow 2} (8x + 3)$

A) 19

B) -13

C) 11

D) 3

Answer: A

38) $\lim_{x \rightarrow 2} (x^2 + 8x - 2)$

A) does not exist

B) 0

C) 18

D) -18

Answer: C

39) $\lim_{x \rightarrow 0} (x^2 - 5)$

A) does not exist

B) 0

C) -5

D) 5

Answer: C

40) $\lim_{x \rightarrow 0} (\sqrt{x} - 2)$

A) -2

B) does not exist

C) 0

D) 2

Answer: A

41) $\lim_{x \rightarrow 2} (x^3 + 5x^2 - 7x + 1)$

A) 0

B) does not exist

C) 29

D) 15

Answer: D

42) $\lim_{x \rightarrow 2} (3x^5 - 3x^4 - 4x^3 + x^2 + 5)$

A) -23

B) 25

C) 121

D) 89

Answer: B

$$43) \lim_{x \rightarrow 5} \sqrt{x^2 + 10x + 25}$$

A) ± 10

B) 10

C) 100

D) does not exist

Answer: B

$$44) \lim_{x \rightarrow 1} \frac{x}{3x + 2}$$

A) does not exist

B) $-\frac{1}{5}$

C) 0

D) 1

Answer: D

Find the limit if it exists.

$$45) \lim_{x \rightarrow 1} \sqrt{10}$$

A) $\sqrt{11}$

B) 11

C) $\sqrt{10}$

D) 10

Answer: C

$$46) \lim_{x \rightarrow 3} (4x - 4)$$

A) 8

B) -8

C) 16

D) -16

Answer: D

$$47) \lim_{x \rightarrow 9} (22 - 6x)$$

A) -32

B) -76

C) 32

D) 76

Answer: D

$$48) \lim_{x \rightarrow 8} (5x^2 - 3x - 6)$$

A) 302

B) 290

C) 350

D) 338

Answer: B

$$49) \lim_{x \rightarrow 1} 9x(x + 4)(x - 4)$$

A) -225

B) -135

C) -81

D) 135

Answer: D

$$50) \lim_{x \rightarrow \frac{1}{6}} 6x \left(x - \frac{2}{5} \right)$$

A) $\frac{17}{30}$

B) $-\frac{7}{30}$

C) $-\frac{7}{5}$

D) $-\frac{7}{180}$

Answer: B

51) $\lim_{x \rightarrow 625} x^{3/4}$

A) 125

B) $\frac{3}{4}$

C) 625

D) $\frac{1875}{4}$

Answer: A

52) $\lim_{x \rightarrow 1} (x + 3)^2(x - 2)^3$

A) -4

B) -16

C) 108

D) 432

Answer: B

53) $\lim_{x \rightarrow 6} \sqrt{4x + 98}$

A) 122

B) -122

C) $-\sqrt{122}$

D) $\sqrt{122}$

Answer: D

54) $\lim_{x \rightarrow 3} (x + 3128)^{3/5}$

A) 125

B) 625

C) -125

D) 25

Answer: A

Find the limit, if it exists.

55) $\lim_{x \rightarrow 14} \frac{1}{x - 14}$

A) 14

B) 28

C) Does not exist

D) 0

Answer: C

56) $\lim_{x \rightarrow 0} \frac{x^3 - 6x + 8}{x - 2}$

A) 4

B) -4

C) Does not exist

D) 0

Answer: B

57) $\lim_{x \rightarrow 1} \frac{2x - 7}{4x + 5}$

A) $-\frac{5}{9}$

B) Does not exist

C) $-\frac{1}{2}$

D) $-\frac{7}{5}$

Answer: A

58) $\lim_{x \rightarrow 1} \frac{3x^2 + 7x - 2}{3x^2 - 4x - 2}$

A) Does not exist

B) $-\frac{8}{3}$

C) $-\frac{7}{4}$

D) 0

Answer: B

67) $\lim_{x \rightarrow 7} \frac{x^2 - 49}{x - 7}$

A) 14

B) 7

C) 1

D) Does not exist

Answer: A

68) $\lim_{x \rightarrow -6} \frac{x^2 + 10x + 24}{x + 6}$

A) -2

B) 120

C) Does not exist

D) 10

Answer: A

69) $\lim_{x \rightarrow 7} \frac{x^2 + 2x - 63}{x - 7}$

A) 0

B) Does not exist

C) 16

D) 2

Answer: C

70) $\lim_{x \rightarrow 8} \frac{x^2 + 2x - 80}{x^2 - 64}$

A) $\frac{9}{8}$

B) $-\frac{1}{8}$

C) 0

D) Does not exist

Answer: A

71) $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x^2 - 3x + 2}$

A) 0

B) 2

C) Does not exist

D) 4

Answer: D

72) $\lim_{x \rightarrow 6} \frac{x^2 + 3x - 18}{x^2 + 2x - 24}$

A) $\frac{3}{10}$

B) Does not exist

C) $-\frac{9}{10}$

D) $\frac{9}{10}$

Answer: D

73) $\lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h}$

A) Does not exist

B) $3x^2 + 3xh + h^2$

C) $3x^2$

D) 0

Answer: C

74) $\lim_{x \rightarrow 7} \frac{|7-x|}{7-x}$

A) 0

B) -1

C) Does not exist

D) 1

Answer: C

81) Let $\lim_{x \rightarrow 9} f(x) = 7$ and $\lim_{x \rightarrow 9} g(x) = -10$. Find $\lim_{x \rightarrow 9} \frac{f(x)}{g(x)}$.

A) $-\frac{10}{7}$

B) 9

C) $-\frac{7}{10}$

D) 17

Answer: C

82) Let $\lim_{x \rightarrow 8} f(x) = 8$. Find $\lim_{x \rightarrow 8} \log_2 f(x)$.

A) 8

B) 3

C) 9

D) $\frac{3}{2}$

Answer: B

83) Let $\lim_{x \rightarrow 9} f(x) = 49$. Find $\lim_{x \rightarrow 9} \sqrt{f(x)}$.

A) 49

B) -9

C) 7

D) 2.6458

Answer: C

84) Let $\lim_{x \rightarrow 7} f(x) = -7$ and $\lim_{x \rightarrow 7} g(x) = -5$. Find $\lim_{x \rightarrow 7} [f(x) + g(x)]^2$.

A) 144

B) -2

C) -12

D) 74

Answer: A

85) Let $\lim_{x \rightarrow 9} f(x) = 4$. Find $\lim_{x \rightarrow 9} (-3)^{f(x)}$.

A) 81

B) -3

C) -19,683

D) 4

Answer: A

86) Let $\lim_{x \rightarrow 8} f(x) = 32$. Find $\lim_{x \rightarrow 8} \sqrt[5]{f(x)}$.

A) 2

B) 32

C) 8

D) 5

Answer: A

87) Let $\lim_{x \rightarrow 4} f(x) = 7$ and $\lim_{x \rightarrow 4} g(x) = -7$. Find $\lim_{x \rightarrow 4} \left[\frac{-4f(x) - 7g(x)}{9 + g(x)} \right]$.

A) $-\frac{77}{2}$

B) $-\frac{91}{9}$

C) -4

D) $\frac{21}{2}$

Answer: D

Evaluate $\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$ for the given x_0 and function f .

88) $f(x) = 3x^2$ for $x_0 = 9$

A) 54

B) 243

C) 27

D) Does not exist

Answer: A

89) $f(x) = 3x^2 + 4$ for $x_0 = 4$

A) 48

B) Does not exist

C) 24

D) 28

Answer: C

90) $f(x) = 4x + 5$ for $x_0 = 2$

A) Does not exist

B) 8

C) 13

D) 4

Answer: D

91) $f(x) = \frac{x}{3} + 4$ for $x_0 = 4$

A) $\frac{16}{3}$

B) $\frac{4}{3}$

C) Does not exist

D) $\frac{1}{3}$

Answer: D

92) $f(x) = \frac{5}{x}$ for $x_0 = 6$

A) $-\frac{5}{36}$

B) Does not exist

C) $\frac{5}{6}$

D) -30

Answer: A

93) $f(x) = 4\sqrt{x}$ for $x_0 = 4$

A) Does not exist

B) 8

C) 4

D) 1

Answer: D

94) $f(x) = \sqrt{x}$ for $x_0 = 13$

A) $\frac{\sqrt{13}}{26}$

B) $\frac{13}{2}$

C) $\frac{\sqrt{13}}{13}$

D) Does not exist

Answer: A

95) $f(x) = 3\sqrt{x} + 8$ for $x_0 = 9$

A) $\frac{27}{2}$

B) $\frac{1}{2}$

C) Does not exist

D) $\frac{9}{2}$

Answer: B

Provide an appropriate response.

96) It can be shown that the inequalities $-x \leq x \cos\left(\frac{1}{x}\right) \leq x$ hold for all values of $x \geq 0$.

Find $\lim_{x \rightarrow 0} x \cos\left(\frac{1}{x}\right)$ if it exists.

A) 1

B) 0

C) 0.0007

D) does not exist

Answer: B

100) Let $f(x) = \frac{x - 4}{\sqrt{x} - 2}$, find $\lim_{x \rightarrow 4} f(x)$.

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)						

A)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	3.97484	3.99750	3.99975	4.00025	4.00250	4.02485

; limit = 4.0

B)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.19245	1.19925	1.19993	1.20007	1.20075	1.20745

; limit = 1.20

C)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	1.19245	1.19925	1.19993	1.20007	1.20075	1.20745

; limit = ∞

D)

x	3.9	3.99	3.999	4.001	4.01	4.1
f(x)	5.07736	5.09775	5.09978	5.10022	5.10225	5.12236

; limit = 5.10

Answer: A

101) Let $f(x) = x^2 - 5$, find $\lim_{x \rightarrow 0} f(x)$.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(x)						

A)

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(x)	-2.9910	-2.9999	-3.0000	-3.0000	-2.9999	-2.9910

; limit = -3.0

B)

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(x)	-1.4970	-1.4999	-1.5000	-1.5000	-1.4999	-1.4970

; limit = -15.0

C)

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(x)	-1.4970	-1.4999	-1.5000	-1.5000	-1.4999	-1.4970

; limit = ∞

D)

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(x)	-4.9900	-4.9999	-5.0000	-5.0000	-4.9999	-4.9900

; limit = -5.0

Answer: D

102) Let $f(x) = \frac{x+5}{x^2+7x+10}$, find $\lim_{x \rightarrow 5} f(x)$.

x	-5.1	-5.01	-5.001	-4.999	-4.99	-4.9
f(x)						

A)

x	-5.1	-5.01	-5.001	-4.999	-4.99	-4.9
f(x)	-0.4226	-0.4322	-0.4332	-0.4334	-0.4344	-0.4448

; limit = -0.4333

B)

x	-5.1	-5.01	-5.001	-4.999	-4.99	-4.9
f(x)	-0.2226	-0.2322	-0.2332	-0.2334	-0.2344	-0.2448

; limit = -0.2333

C)

x	-5.1	-5.01	-5.001	-4.999	-4.99	-4.9
f(x)	0.3226	0.3322	0.3332	0.3334	0.3344	0.3448

; limit = 0.3333

D)

x	-5.1	-5.01	-5.001	-4.999	-4.99	-4.9
f(x)	-0.3226	-0.3322	-0.3332	-0.3334	-0.3344	-0.3448

; limit = -0.3333

Answer: D

103) Let $f(x) = \frac{x^2 - 4x - 5}{x^2 - 7x + 10}$, find $\lim_{x \rightarrow 5} f(x)$.

x	4.9	4.99	4.999	5.001	5.01	5.1
f(x)						

A)

x	4.9	4.99	4.999	5.001	5.01	5.1
f(x)	2.1345	2.1033	2.1003	2.0997	2.0967	2.0677

; limit = 2.1

B)

x	4.9	4.99	4.999	5.001	5.01	5.1
f(x)	1.9345	1.9033	1.9003	1.8997	1.8967	1.8677

; limit = 1.9

C)

x	4.9	4.99	4.999	5.001	5.01	5.1
f(x)	2.0345	2.0033	2.0003	1.9997	1.9967	1.9677

; limit = 2

D)

x	4.9	4.99	4.999	5.001	5.01	5.1
f(x)	0.5775	0.5720	0.5715	0.5714	0.5708	0.5652

; limit = 0.5714

Answer: C

104) Let $f(x) = \frac{\sin(6x)}{x}$, find $\lim_{x \rightarrow 0} f(x)$.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(x)		5.99640065			5.99640065	

A) limit = 6

B) limit = 5.5

C) limit does not exist

D) limit = 0

Answer: A

105) Let $f(\theta) = \frac{\cos(4\theta)}{\theta}$, find $\lim_{\theta \rightarrow 0} f(\theta)$.

x	-0.1	-0.01	-0.001	0.001	0.01	0.1
f(θ)	-9.2106099					9.2106099

- A) limit = 9.2106099 B) limit does not exist C) limit = 4 D) limit = 0

Answer: B

Find the limit.

106) If $\lim_{x \rightarrow 4} \frac{f(x) - 3}{x - 4} = 4$, find $\lim_{x \rightarrow 4} f(x)$.

- A) 3 B) 1 C) -9 D) Does not exist

Answer: C

107) If $\lim_{x \rightarrow 2} \frac{f(x)}{x} = 3$, find $\lim_{x \rightarrow 2} f(x)$.

- A) 2 B) 3 C) 6 D) Does not exist

Answer: C

108) If $\lim_{x \rightarrow 2} \frac{f(x)}{x^2} = 4$, find $\lim_{x \rightarrow 2} \frac{f(x)}{x}$.

- A) 2 B) 8 C) 16 D) 4

Answer: B

109) If $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$, find $\lim_{x \rightarrow 0} f(x)$.

- A) 0 B) 2 C) 1 D) Does not exist

Answer: A

110) If $\lim_{x \rightarrow 0} \frac{f(x)}{x^2} = 1$, find $\lim_{x \rightarrow 0} \frac{f(x)}{x}$.

- A) 1 B) 0 C) 2 D) Does not exist

Answer: B

111) If $\lim_{x \rightarrow 1} \frac{f(x) - 3}{x - 1} = 2$, find $\lim_{x \rightarrow 1} f(x)$.

- A) 1 B) 2 C) 3 D) Does not exist

Answer: C

Use a CAS to plot the function near the point x_0 being approached. From your plot guess the value of the limit.

112) $\lim_{x \rightarrow 36} \frac{\sqrt{x} - 6}{x - 36}$

- A) $\frac{1}{12}$ B) $\frac{1}{6}$ C) 6 D) 0

Answer: A

$$113) \lim_{x \rightarrow 36} \frac{6 - \sqrt{x}}{36 - x}$$

A) 12

B) 0

C) $\frac{1}{12}$

D) 6

Answer: C

$$114) \lim_{x \rightarrow 0} \frac{\sqrt{81+x} - \sqrt{81-x}}{x}$$

A) $\frac{1}{9}$

B) 9

C) $\frac{1}{18}$

D) 0

Answer: A

$$115) \lim_{x \rightarrow 0} \frac{\sqrt{1-x} - 1}{x}$$

A) 2

B) $\frac{1}{2}$

C) 1

D) $-\frac{1}{2}$

Answer: D

$$116) \lim_{x \rightarrow 0} \frac{\sqrt{36+2x} - 6}{x}$$

A) $\frac{1}{6}$

B) 36

C) $\frac{1}{3}$

D) $\frac{1}{12}$

Answer: A

$$117) \lim_{x \rightarrow 0} \frac{\sqrt{6+6x} - \sqrt{6}}{x}$$

A) 0

B) $\frac{\sqrt{6}}{2}$

C) $\sqrt{6}$

D) $\frac{1}{2}$

Answer: B

$$118) \lim_{x \rightarrow 0} \frac{4 - \sqrt{16-x^2}}{x}$$

A) 8

B) $\frac{1}{8}$

C) $\frac{1}{4}$

D) 0

Answer: D

$$119) \lim_{x \rightarrow 3} \frac{x^2 - 9}{\sqrt{x^2 + 7} - 4}$$

A) 3

B) 8

C) $\frac{1}{4}$

D) 4

Answer: B

120) $\lim_{x \rightarrow 1} \frac{x^2 - 1}{\sqrt{x^2 + 3} - 2}$
 A) $\frac{1}{4}$

B) 4

C) 2

D) 1

Answer: B

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Provide an appropriate response.

121) It can be shown that the inequalities $1 - \frac{x^2}{6} < \frac{x \sin(x)}{2 - 2 \cos(x)} < 1$ hold for all values of x close to zero. What, if anything, does this tell you about $\frac{x \sin(x)}{2 - 2 \cos(x)}$? Explain.

Answer: Answers may vary. One possibility: $\lim_{x \rightarrow 0} 1 - \frac{x^2}{6} = \lim_{x \rightarrow 0} 1 = 1$. According to the squeeze theorem, the function $\frac{x \sin(x)}{2 - 2 \cos(x)}$, which is squeezed between $1 - \frac{x^2}{6}$ and 1, must also approach 1 as x approaches 0. Thus, $\lim_{x \rightarrow 0} \frac{x \sin(x)}{2 - 2 \cos(x)} = 1$.

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

122) Write the formal notation for the principle "the limit of a quotient is the quotient of the limits" and include a statement of any restrictions on the principle.

A) $\lim_{x \rightarrow a} \frac{g(x)}{f(x)} = \frac{g(a)}{f(a)}$.

B) If $\lim_{x \rightarrow a} g(x) = M$ and $\lim_{x \rightarrow a} f(x) = L$, then $\lim_{x \rightarrow a} \frac{g(x)}{f(x)} = \frac{\lim_{x \rightarrow a} g(x)}{\lim_{x \rightarrow a} f(x)} = \frac{M}{L}$, provided that $f(a) \neq 0$.

C) $\lim_{x \rightarrow a} \frac{g(x)}{f(x)} = \frac{g(a)}{f(a)}$, provided that $f(a) \neq 0$.

D) If $\lim_{x \rightarrow a} g(x) = M$ and $\lim_{x \rightarrow a} f(x) = L$, then $\lim_{x \rightarrow a} \frac{g(x)}{f(x)} = \frac{\lim_{x \rightarrow a} g(x)}{\lim_{x \rightarrow a} f(x)} = \frac{M}{L}$, provided that $L \neq 0$.

Answer: D

123) What conditions, when present, are sufficient to conclude that a function $f(x)$ has a limit as x approaches some value of a ?

A) The limit of $f(x)$ as $x \rightarrow a$ from the left exists, the limit of $f(x)$ as $x \rightarrow a$ from the right exists, and at least one of these limits is the same as $f(a)$.

B) The limit of $f(x)$ as $x \rightarrow a$ from the left exists, the limit of $f(x)$ as $x \rightarrow a$ from the right exists, and these two limits are the same.

C) Either the limit of $f(x)$ as $x \rightarrow a$ from the left exists or the limit of $f(x)$ as $x \rightarrow a$ from the right exists

D) $f(a)$ exists, the limit of $f(x)$ as $x \rightarrow a$ from the left exists, and the limit of $f(x)$ as $x \rightarrow a$ from the right exists.

Answer: B

124) Provide a short sentence that summarizes the general limit principle given by the formal notation $\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x) = L \pm M$, given that $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow a} g(x) = M$.

- A) The limit of a sum or a difference is the sum or the difference of the functions.
- B) The sum or the difference of two functions is continuous.
- C) The limit of a sum or a difference is the sum or the difference of the limits.
- D) The sum or the difference of two functions is the sum of two limits.

Answer: C

125) The statement "the limit of a constant times a function is the constant times the limit" follows from a combination of two fundamental limit principles. What are they?

- A) The limit of a function is a constant times a limit, and the limit of a constant is the constant.
- B) The limit of a product is the product of the limits, and a constant is continuous.
- C) The limit of a product is the product of the limits, and the limit of a quotient is the quotient of the limits.
- D) The limit of a constant is the constant, and the limit of a product is the product of the limits.

Answer: D

Given the interval (a, b) on the x -axis with the point x_0 inside, find the greatest value for $\delta > 0$ such that for all x , $0 < |x - x_0| < \delta \Rightarrow a < x < b$.

126) $a = 9, b = 19, x_0 = 16$

- A) 1
- B) 3
- C) 4
- D) 7

Answer: B

127) $a = \frac{3}{9}, b = \frac{9}{9}, x_0 = \frac{4}{9}$

- A) $\delta = \frac{5}{9}$
- B) $\delta = \frac{1}{9}$
- C) $\delta = 0$
- D) $\delta = 1$

Answer: B


128) $a = 1.317, b = 2.713, x_0 = 1.976$

- A) $\delta = 1$
- B) $\delta = 0.659$
- C) $\delta = 1.396$
- D) $\delta = 0.737$

Answer: B

Use the graph to find a $\delta > 0$ such that for all x , $0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \epsilon$.

129)



$$y = 2x + 3$$

$$f(x) = 2x + 3$$

$$x_0 = 2$$

$$L = 7$$

$$\epsilon = 0.2$$

1.9 2 2.1

NOT TO SCALE

A) 0.1


B) 5

C) 0.4

D) 0.2

Answer: A

130)



$$y = 4x - 2$$

$$f(x) = 4x - 2$$

$$x_0 = 2$$

$$L = 6$$

$$\epsilon = 0.2$$

2
1.95 2.05

NOT TO SCALE

A) 0.05

B) 0.1


C) 0.5

D) 4

Answer: A

131)

$f(x) = -5x - 3$
 $x_0 = -2$
 $L = 7$
 $\varepsilon = 0.2$



7.2

7

6.8

$$\begin{array}{c} \swarrow -2 \searrow \\ -2.04 \quad -1.96 \end{array}$$

NOT TO SCALE

A) 15

B) 0.4


C) 0.04

D) -0.04

Answer: C

132)

$f(x) = -x + 3$
 $x_0 = -1$
 $L = 4$
 $\varepsilon = 0.2$



4.2

4

3.8

$$-1.2 \quad -1 \quad -0.8$$

NOT TO SCALE

A) 5


B) 0.4

C) 0.2

D) -0.2

Answer: C

133)



$$y = \frac{3}{2}x + 3$$

$$f(x) = \frac{3}{2}x + 3$$

$$x_0 = 1$$

$$L = 4.5$$

$$\varepsilon = 0.2$$

0.9 1 1.1

NOT TO SCALE

A) 0.2

B) 0.1

C) 3.5

D) -0.2

Answer: B


134)

$$f(x) = -\frac{4}{3}x + 1$$

$$x_0 = -1$$

$$L = 2.3$$

$$\varepsilon = 0.2$$



2.5

2.3

2.1

-1.1 -1 -0.8

NOT TO SCALE

A) 0.3


B) 3.3

C) -0.3

D) 0.1

Answer: D

135)



$$y = 2\sqrt{x}$$

$$f(x) = 2\sqrt{x}$$

$$x_0 = 3$$

$$L = 2\sqrt{3}$$

$$\varepsilon = \frac{1}{4}$$

- 3.71
- 3.46
- 3.21

2.5831 3 3.4481

NOT TO SCALE

A) 0.4481


B) 0.46

C) 0.865

D) 0.4169

Answer: D

136)



$$y = \sqrt{x-2}$$

$$f(x) = \sqrt{x-2}$$

$$x_0 = 3$$

$$L = 1$$

$$\varepsilon = \frac{1}{4}$$

- 1.25
- 1
- 0.75

2.5625 3 3.5625

NOT TO SCALE

A) 0.5625


B) 2

C) 1

D) 0.4375

Answer: D

137)



$$y = x^2$$

$$f(x) = x^2$$

$$x_0 = 2$$

$$L = 4$$

$$\varepsilon = 1$$

3

$$\begin{array}{c} / \quad 2 \quad \backslash \\ 1.73 \quad 2.24 \end{array}$$

NOT TO SCALE

A) 0.51


B) 0.27

C) 0.24

D) 2

Answer: C

138)



$$y = x^2 - 3$$

$$f(x) = x^2 - 3$$

$$x_0 = 3$$

$$L = 6$$

$$\varepsilon = 1$$

$$\begin{array}{c} / \quad 3 \quad \backslash \\ 2.83 \quad 3.16 \end{array}$$

NOT TO SCALE

A) 0.17

B) 3

C) 0.33

D) 0.16

Answer: D

A function $f(x)$, a point x_0 , the limit of $f(x)$ as x approaches x_0 , and a positive number ε is given. Find a number $\delta > 0$ such that for all x , $0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon$.

139) $f(x) = 5x + 2$, $L = 17$, $x_0 = 3$, and $\varepsilon = 0.01$

A) 0.003333

B) 0.004

C) 0.002

D) 0.01

Answer: C

140) $f(x) = 3x - 2$, $L = 1$, $x_0 = 1$, and $\varepsilon = 0.01$

A) 0.003333

B) 0.001667

C) 0.006667

D) 0.01

Answer: A

141) $f(x) = -10x + 5$, $L = -15$, $x_0 = 2$, and $\epsilon = 0.01$

A) 0.004

B) 0.001

C) 0.002

D) -0.005

Answer: B

142) $f(x) = -10x - 4$, $L = -24$, $x_0 = 2$, and $\epsilon = 0.01$

A) 0.002

B) 0.0005

C) -0.005

D) 0.001

Answer: D

143) $f(x) = \sqrt{x + 5}$, $L = 3$, $x_0 = 4$, and $\epsilon = 1$

A) 16

B) 5

C) 4

D) 7

Answer: B

144) $f(x) = \sqrt{18 - x}$, $L = 4$, $x_0 = 2$, and $\epsilon = 1$

A) -9

B) 9

C) 7

D) 12

Answer: C

145) $f(x) = 3x^2$, $L = 243$, $x_0 = 9$, and $\epsilon = 0.5$

A) 0.00926

B) 9.00925

C) 0.00925

D) 8.99074

Answer: C

146) $f(x) = 1/x$, $L = 1/7$, $x_0 = 7$, and $\epsilon = 0.4$

A) 0.7368

B) 5.1579

C) -27.2222

D) -10.8889

Answer: B

147) $f(x) = mx$, $m > 0$, $L = 4m$, $x_0 = 4$, and $\epsilon = 0.07$

A) $\delta = 4 - m$

B) $\delta = 0.07$

C) $\delta = \frac{0.07}{m}$

D) $\delta = 4 + \frac{0.07}{m}$

Answer: C

148) $f(x) = mx + b$, $m > 0$, $L = (m/8) + b$, $x_0 = 1/8$, and $\epsilon = c > 0$

A) $\delta = \frac{8}{m}$

B) $\delta = \frac{1}{8} + \frac{c}{m}$

C) $\delta = \frac{c}{m}$

D) $\delta = \frac{c}{8}$

Answer: C

Find the limit L for the given function f , the point x_0 , and the positive number ϵ . Then find a number $\delta > 0$ such that, for all x , $0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \epsilon$.

149) $f(x) = 6x + 3$, $x_0 = -5$, $\epsilon = 0.06$

A) $L = -27$; $\delta = 0.01$

B) $L = -33$; $\delta = 0.01$

C) $L = -27$; $\delta = 0.02$

D) $L = 33$; $\delta = 0.02$

Answer: A

150) $f(x) = \frac{x^2 + 4x - 21}{x + 7}$, $x_0 = -7$, $\epsilon = 0.03$

A) $L = 0$; $\delta = 0.03$

B) $L = -6$; $\delta = 0.04$

C) $L = 4$; $\delta = 0.04$

D) $L = -10$; $\delta = 0.03$

Answer: D

151) $f(x) = \sqrt{24 - 3x}$, $x_0 = -4$, $\varepsilon = 0.3$

A) $L = -5$; $\delta = 0.57$

B) $L = 6$; $\delta = 1.17$

C) $L = 7$; $\delta = 1.17$

D) $L = 6$; $\delta = 1.23$

Answer: B

152) $f(x) = \frac{10}{x}$, $x_0 = 5$, $\varepsilon = 0.1$

A) $L = 2$; $\delta = 0.53$

B) $L = 2$; $\delta = 0.24$

C) $L = 2$; $\delta = 0.26$

D) $L = 2$; $\delta = 2.63$

Answer: C

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Prove the limit statement

153) $\lim_{x \rightarrow 5} (5x - 3) = 22$

Answer:

Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon/5$. Then $0 < |x - 5| < \delta$ implies that

$$\begin{aligned} |(5x - 3) - 22| &= |5x - 25| \\ &= |5(x - 5)| \\ &= 5|x - 5| < 5\delta = \varepsilon \end{aligned}$$

Thus, $0 < |x - 5| < \delta$ implies that $|(5x - 3) - 22| < \varepsilon$

154) $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = 4$

Answer: Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon$. Then $0 < |x - 2| < \delta$ implies that

$$\begin{aligned} \left| \frac{x^2 - 4}{x - 2} - 4 \right| &= \left| \frac{(x - 2)(x + 2)}{x - 2} - 4 \right| \\ &= |(x + 2) - 4| \quad \text{for } x \neq 2 \\ &= |x - 2| < \delta = \varepsilon \end{aligned}$$

Thus, $0 < |x - 2| < \delta$ implies that $\left| \frac{x^2 - 4}{x - 2} - 4 \right| < \varepsilon$

155) $\lim_{x \rightarrow 6} \frac{3x^2 - 14x - 24}{x - 6} = 22$

Answer: Let $\varepsilon > 0$ be given. Choose $\delta = \varepsilon/3$. Then $0 < |x - 6| < \delta$ implies that

$$\begin{aligned} \left| \frac{3x^2 - 14x - 24}{x - 6} - 22 \right| &= \left| \frac{(x - 6)(3x + 4)}{x - 6} - 22 \right| \\ &= |(3x + 4) - 22| \quad \text{for } x \neq 6 \\ &= |3x - 18| \\ &= |3(x - 6)| \\ &= 3|x - 6| < 3\delta = \varepsilon \end{aligned}$$

Thus, $0 < |x - 6| < \delta$ implies that $\left| \frac{3x^2 - 14x - 24}{x - 6} - 22 \right| < \varepsilon$

$$156) \lim_{x \rightarrow 5} \frac{1}{x} = \frac{1}{5}$$

Answer: Let $\varepsilon > 0$ be given. Choose $\delta = \min\{5/2, 25\varepsilon/2\}$. Then $0 < |x - 5| < \delta$ implies that

$$\begin{aligned} \left| \frac{1}{x} - \frac{1}{5} \right| &= \left| \frac{5-x}{5x} \right| \\ &= \frac{1}{|x|} \cdot \frac{1}{5} \cdot |x-5| \\ &< \frac{1}{5/2} \cdot \frac{1}{5} \cdot \frac{25\varepsilon}{2} = \varepsilon \end{aligned}$$

Thus, $0 < |x - 5| < \delta$ implies that $\left| \frac{1}{x} - \frac{1}{5} \right| < \varepsilon$

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Solve the problem.

- 157) You are asked to make some circular cylinders, each with a cross-sectional area of 9 cm^2 . To do this, you need to know how much deviation from the ideal cylinder diameter of $x_0 = 1.93 \text{ cm}$ you can allow and still have the area come within 0.1 cm^2 of the required 9 cm^2 . To find out, let $A = \pi \left(\frac{x}{2} \right)^2$ and look for the interval in which you must hold x to make $|A - 9| < 0.1$. What interval do you find?
- A) $(3.3663, 3.4039)$ B) $(0.5642, 0.5642)$ C) $(2.3803, 2.4069)$ D) $(5.9666, 6.0332)$

Answer: A

- 158) Ohm's Law for electrical circuits is stated $V = RI$, where V is a constant voltage, R is the resistance in ohms and I is the current in amperes. Your firm has been asked to supply the resistors for a circuit in which V will be 12 volts and I is to be 3 ± 0.1 amperes. In what interval does R have to lie for I to be within 0.1 amps of the target value $I_0 = 3$?

A) $\left(\frac{31}{120}, \frac{29}{120} \right)$ B) $\left(\frac{120}{29}, \frac{120}{31} \right)$ C) $\left(\frac{10}{29}, \frac{10}{31} \right)$ D) $\left(\frac{120}{31}, \frac{120}{29} \right)$

Answer: D

- 159) The cross-sectional area of a cylinder is given by $A = \pi D^2/4$, where D is the cylinder diameter. Find the tolerance range of D such that $|A - 10| < 0.01$ as long as $D_{\min} < D < D_{\max}$.

A) $D_{\min} = 3.567, D_{\max} = 3.578$ B) $D_{\min} = 3.567, D_{\max} = 3.570$
 C) $D_{\min} = 3.558, D_{\max} = 3.570$ D) $D_{\min} = 3.558, D_{\max} = 3.578$

Answer: B

- 160) The current in a simple electrical circuit is given by $I = V/R$, where I is the current in amperes, V is the voltage in volts, and R is the resistance in ohms. When $V = 12$ volts, what is a 12Ω resistor's tolerance for the current to be within 1 ± 0.01 amp?

A) 0.1% B) 1% C) 10% D) 0.01%

Answer: B

161) Select the correct statement for the definition of the limit: $\lim_{x \rightarrow x_0} f(x) = L$

means that _____

- A) if given any number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x , $0 < |x - x_0| < \varepsilon$ implies $|f(x) - L| > \delta$.
- B) if given any number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x , $0 < |x - x_0| < \varepsilon$ implies $|f(x) - L| < \delta$.
- C) if given a number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x , $0 < |x - x_0| < \delta$ implies $|f(x) - L| > \varepsilon$.
- D) if given any number $\varepsilon > 0$, there exists a number $\delta > 0$, such that for all x , $0 < |x - x_0| < \delta$ implies $|f(x) - L| < \varepsilon$.

Answer: D

162) Identify the incorrect statements about limits.


- I. The number L is the limit of $f(x)$ as x approaches x_0 if $f(x)$ gets closer to L as x approaches x_0 .
- II. The number L is the limit of $f(x)$ as x approaches x_0 if, for any $\varepsilon > 0$, there corresponds a $\delta > 0$ such that $|f(x) - L| < \varepsilon$ whenever $0 < |x - x_0| < \delta$.
- III. The number L is the limit of $f(x)$ as x approaches x_0 if, given any $\varepsilon > 0$, there exists a value of x for which $|f(x) - L| < \varepsilon$.

- A) I and II
- B) II and III
- C) I and III
- D) I, II, and III

Answer: C

Use the graph to estimate the specified limit.


163) Find $\lim_{x \rightarrow (-1)^-} f(x)$ and $\lim_{x \rightarrow (-1)^+} f(x)$



- A) -7; -5
- B) -5; -2
- C) -2; -7
- D) -7; -2

Answer: C

164) Find $\lim_{x \rightarrow 0} f(x)$




A) does not exist
Answer: A

B) -2

C) 2

D) 0

165) Find $\lim_{x \rightarrow 0} f(x)$




A) -1
Answer: D

B) 1

C) does not exist

D) 0

166) Find $\lim_{x \rightarrow 3^-} f(x)$



A) $-\frac{2\sqrt{3}}{3}$


B) -3

C) $\frac{2}{3}$

D) $\frac{2\sqrt{3}}{3}$

Answer: D

167) Find $\lim_{x \rightarrow (\pi/2)^-} f(x)$ and $\lim_{x \rightarrow (\pi/2)^+} f(x)$



A) 1; 4


B) π ; π

C) 4; 1

D) $\frac{\pi}{2}$; $\frac{\pi}{2}$

Answer: C

168) Find $\lim_{x \rightarrow 0^-} f(x)$ and $\lim_{x \rightarrow 0^+} f(x)$



A) 2; 1


B) -2; -1

C) 2; -1

D) -1; 2

Answer: C

169) Find $\lim_{x \rightarrow 2^-} f(x)$ and $\lim_{x \rightarrow 2^+} f(x)$



A) 5; -3


C) -3; 5

B) 1; 1

D) does not exist; does not exist

Answer: A


170) Find $\lim_{x \rightarrow 0} f(x)$



- A) does not exist B) -1 C) 0 D) 6
 Answer: C


Determine the limit by sketching an appropriate graph.

171) $\lim_{x \rightarrow 3^-} f(x)$, where $f(x) = \begin{cases} -4x + 1 & \text{for } x < 3 \\ 2x + 2 & \text{for } x \geq 3 \end{cases}$




- A) 2 B) 3 C) -11 D) 8
 Answer: C

172) $\lim_{x \rightarrow 7^+} f(x)$, where $f(x) = \begin{cases} -2x - 7 & \text{for } x < 7 \\ 2x - 6 & \text{for } x \geq 7 \end{cases}$



- A) -21 B) 8 C) -6 D) -5
 Answer: B

173) $\lim_{x \rightarrow 4^+} f(x)$, where $f(x) = \begin{cases} x^2 + 3 & \text{for } x \neq 4 \\ 0 & \text{for } x = 4 \end{cases}$



A) 13


B) 19

C) 0

D) 16

Answer: B

174) $\lim_{x \rightarrow 2^-} f(x)$, where $f(x) = \begin{cases} \sqrt{1-x^2} & 0 \leq x < 1 \\ 1 & 1 \leq x < 2 \\ 2 & x = 2 \end{cases}$



A) 2


B) 0

C) Does not exist

D) 1

Answer: D

175) $\lim_{x \rightarrow 7^+} f(x)$, where $f(x) = \begin{cases} x & -7 \leq x < 0, \text{ or } 0 < x \leq 3 \\ 1 & x = 0 \\ 0 & x < -7 \text{ or } x > 3 \end{cases}$



A) -7

B) -0

C) 1

D) Does not exist

Answer: A

Find the limit.

$$176) \lim_{x \rightarrow 0.5^-} \sqrt{\frac{x+7}{x+5}}$$

A) $\sqrt{\frac{13}{9}}$

B) $\sqrt{\frac{15}{11}}$

C) Does not exist

D) $\frac{15}{11}$

Answer: B

$$177) \lim_{x \rightarrow 2^+} \sqrt{\frac{7x^2}{5+x}}$$

A) $\sqrt{\frac{28}{3}}$

B) $\frac{28}{3}$

C) Does not exist

D) $\sqrt{-\frac{14}{3}}$

Answer: A

$$178) \lim_{x \rightarrow 1^+} \left(\frac{x}{x+9} \right) \left(\frac{-3x+8}{x^2+9x} \right)$$

A) Does not exist

B) $\frac{11}{82}$

C) $\frac{11}{64}$

D) $\frac{1}{11}$

Answer: C

$$179) \lim_{h \rightarrow 0^+} \frac{\sqrt{h^2 + 13h + 14} - \sqrt{14}}{h}$$

A) $\frac{13}{\sqrt{28}}$

B) $\frac{13}{2\sqrt{14}}$

C) $\frac{13}{28}$

D) Does not exist

Answer: B

$$180) \lim_{h \rightarrow 0^-} \frac{\sqrt{10} - \sqrt{2h^2 + 7h + 10}}{h}$$

A) $\frac{-7}{2\sqrt{10}}$

B) $\frac{7}{2\sqrt{10}}$

C) $\frac{-7}{\sqrt{20}}$

D) Does not exist

Answer: A

$$181) \lim_{x \rightarrow 2^+} (x+5) \left(\frac{|x+2|}{x+2} \right)$$

A) Does not exist

B) 3

C) -3

D) 7

Answer: B

$$182) \lim_{x \rightarrow 1^-} (x+3) \left(\frac{|x+1|}{x+1} \right)$$

A) 2

B) Does not exist

C) -2

D) 4

Answer: C

$$183) \lim_{x \rightarrow 5^-} \frac{\sqrt{3x}(x-5)}{|x-5|}$$

A) Does not exist

B) 0

C) $-\sqrt{15}$

D) $\sqrt{15}$

Answer: C

$$184) \lim_{x \rightarrow 3^+} \frac{\sqrt{5x}(x-3)}{|x-3|}$$

A) 0

B) $-\sqrt{15}$

C) $\sqrt{15}$

D) Does not exist

Answer: C

Use the graph of the greatest integer function $y = \lfloor x \rfloor$ to find the limit.

$$185) \lim_{x \rightarrow 2^-} \frac{\lfloor x \rfloor}{x}$$

A) 2

B) 0

C) 1

D) -2

Answer: C

$$186) \lim_{x \rightarrow 6^-} (x - \lfloor x \rfloor)$$

A) 12

B) 6

C) 0

D) -12

Answer: C

Find the limit using $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$.

$$187) \lim_{x \rightarrow 0} \frac{\sin 5x}{x}$$

A) does not exist

B) 1

C) $\frac{1}{5}$

D) 5

Answer: D

$$188) \lim_{x \rightarrow 0} \frac{x}{\sin 3x}$$

A) 3

B) 1

C) $\frac{1}{3}$

D) does not exist

Answer: C

$$189) \lim_{x \rightarrow 0} \frac{\tan 4x}{x}$$

A) $\frac{1}{4}$

B) 4

C) 1

D) does not exist

Answer: B

190) $\lim_{x \rightarrow 0} \frac{\sin 5x}{\sin 4x}$

- A) does not exist B) $\frac{4}{5}$ C) $\frac{5}{4}$ D) 0

Answer: C

191) $\lim_{x \rightarrow 0} \frac{\sin 4x}{\sin 5x}$

- A) does not exist B) $\frac{5}{4}$ C) $\frac{4}{5}$ D) 0

Answer: C

192) $\lim_{x \rightarrow 0} \frac{\sin x \cos 4x}{x + x \cos 5x}$

- A) 0 B) $\frac{4}{5}$ C) $\frac{1}{2}$ D) does not exist

Answer: C

193) $\lim_{x \rightarrow 0} 6x^2(\cot 3x)(\csc 2x)$

- A) 1 B) does not exist C) $\frac{1}{3}$ D) $\frac{1}{2}$

Answer: A

194) $\lim_{x \rightarrow 0} \frac{x^2 - 2x + \sin x}{x}$

- A) does not exist B) 0 C) -1 D) 1

Answer: C

195) $\lim_{x \rightarrow 0} \frac{\sin(\sin x)}{\sin x}$

- A) 1 B) -1 C) does not exist D) 0

Answer: A

196) $\lim_{x \rightarrow 0} \frac{\sin 3x \cot 4x}{\cot 5x}$

- A) 0 B) $\frac{15}{4}$ C) does not exist D) $\frac{12}{5}$

Answer: A

Provide an appropriate response.

197) Given $\lim_{x \rightarrow \theta^-} f(x) = L_I$, $\lim_{x \rightarrow \theta^+} f(x) = L_R$, and $L_I \neq L_R$, which of the following statements is true?

- I. $\lim_{x \rightarrow \theta} f(x) = L_I$
- II. $\lim_{x \rightarrow \theta} f(x) = L_R$
- III. $\lim_{x \rightarrow \theta} f(x)$ does not exist.

A) I B) none C) II D) III

Answer: D

198) Given $\lim_{x \rightarrow \theta^-} f(x) = L_I$, $\lim_{x \rightarrow \theta^+} f(x) = L_R$, and $L_I = L_R$, which of the following statements is false?

- I. $\lim_{x \rightarrow \theta} f(x) = L_I$
- II. $\lim_{x \rightarrow \theta} f(x) = L_R$
- III. $\lim_{x \rightarrow \theta} f(x)$ does not exist.

A) II B) none C) III D) I

Answer: C

199) If $\lim_{x \rightarrow \theta} f(x) = L$, which of the following expressions are true?

- I. $\lim_{x \rightarrow \theta^-} f(x)$ does not exist.
- II. $\lim_{x \rightarrow \theta^+} f(x)$ does not exist.
- III. $\lim_{x \rightarrow \theta^-} f(x) = L$
- IV. $\lim_{x \rightarrow \theta^+} f(x) = L$

A) I and II only B) II and III only C) III and IV only D) I and IV only

Answer: C

200) If $\lim_{x \rightarrow \theta^-} f(x) = 1$ and $f(x)$ is an odd function, which of the following statements are true?

- I. $\lim_{x \rightarrow \theta} f(x) = 1$
- II. $\lim_{x \rightarrow \theta^+} f(x) = -1$
- III. $\lim_{x \rightarrow \theta} f(x)$ does not exist.

A) II and III only B) I and III only C) I, II, and III D) I and II only

Answer: A

201) If $\lim_{x \rightarrow 1^-} f(x) = 1$, $\lim_{x \rightarrow 1^+} f(x) = -1$, and $f(x)$ is an even function, which of the following statements are true?

- I. $\lim_{x \rightarrow 1^-} f(x) = -1$
- II. $\lim_{x \rightarrow 1^+} f(x) = -1$
- III. $\lim_{x \rightarrow 1} f(x)$ does not exist.

- A) I and III only B) I and II only C) I, II, and III D) II and III only

Answer: A

202) Given $\epsilon > 0$, find an interval $I = (1, 1 + \delta)$, $\delta > 0$, such that if x lies in I , then $\sqrt{x - 1} < \epsilon$. What limit is being verified and what is its value?

- A) $\lim_{x \rightarrow 1^+} \sqrt{x} = 1$ B) $\lim_{x \rightarrow 0^-} \sqrt{x - 1} = 0$ C) $\lim_{x \rightarrow 1^+} \sqrt{x - 1} = 0$ D) $\lim_{x \rightarrow 1^-} \sqrt{x - 1} = 0$

Answer: C


203) Given $\epsilon > 0$, find an interval $I = (5 - \delta, 5)$, $\delta > 0$, such that if x lies in I , then $\sqrt{5 - x} < \epsilon$. What limit is being verified and what is its value?

- A) $\lim_{x \rightarrow 5^-} \sqrt{x} = 5$ B) $\lim_{x \rightarrow 5^+} \sqrt{5 - x} = 0$ C) $\lim_{x \rightarrow 0^-} \sqrt{5 - x} = 0$ D) $\lim_{x \rightarrow 5^-} \sqrt{5 - x} = 0$

Answer: D

Find all points where the function is discontinuous.


204)



- A) $x = 4$ B) $x = 4, x = 2$ C) None D) $x = 2$

Answer: A


205)



- A) $x = -2, x = 1$ B) $x = 1$ C) None D) $x = -2$

Answer: B

206)



A) $x = 2$


B) $x = -2, x = 0, x = 2$

C) $x = -2, x = 0$

D) $x = 0, x = 2$

Answer: B

207)



A) None


B) $x = -2$

C) $x = 6$

D) $x = -2, x = 6$

Answer: C

208)



A) $x = 1, x = 4, x = 5$


B) None

C) $x = 1, x = 5$

D) $x = 4$

Answer: B

209)



A) $x = 1$


B) $x = 0$

C) None

D) $x = 0, x = 1$

Answer: C

210)



A) $x = 0$


B) $x = 3$

C) $x = 0, x = 3$

D) None

Answer: B

211)



A) $x = 2$


B) None

C) $x = -2, x = 2$

D) $x = -2$

Answer: C

212)



A) $x = -2, x = 0, x = 2$

B) $x = 0$

C) None


D) $x = -2, x = 2$

Answer: B

Answer the question.

213) Does $\lim_{x \rightarrow (-1)^+} f(x)$ exist?

$$f(x) = \begin{cases} -x^2 + 1, & -1 \leq x < 0 \\ 2x, & 0 < x < 1 \\ -4, & x = 1 \\ -2x + 4, & 1 < x < 3 \\ 5, & 3 < x < 5 \end{cases}$$




A) Yes

B) No

Answer: A

214) Does $\lim_{x \rightarrow 1^+} f(x) = f(-1)$?

$$f(x) = \begin{cases} -x^2 + 1, & -1 \leq x < 0 \\ 3x, & 0 < x < 1 \\ -2, & x = 1 \\ -3x + 6, & 1 < x < 3 \\ 1, & 3 < x < 5 \end{cases}$$




A) No
Answer: B

B) Yes

215) Does $\lim_{x \rightarrow 1} f(x)$ exist?

$$f(x) = \begin{cases} -x^2 + 1, & -1 \leq x < 0 \\ 2x, & 0 < x < 1 \\ -2, & x = 1 \\ -2x + 4, & 1 < x < 3 \\ 3, & 3 < x < 5 \end{cases}$$




A) No
Answer: B

B) Yes

216) Is f continuous at $f(1)$?

$$f(x) = \begin{cases} -x^2 + 1, & -1 \leq x < 0 \\ 2x, & 0 < x < 1 \\ -5, & x = 1 \\ -2x + 4, & 1 < x < 3 \\ 1, & 3 < x < 5 \end{cases}$$




A) No

B) Yes

Answer: A

217) Is f continuous at $f(3)$?

$$f(x) = \begin{cases} -x^2 + 1, & -1 \leq x < 0 \\ 4x, & 0 < x < 1 \\ -5, & x = 1 \\ -4x + 8, & 1 < x < 3 \\ 5, & 3 < x < 5 \end{cases}$$




A) Yes

B) No

Answer: B

218) Does $\lim_{x \rightarrow 0} f(x)$ exist?

$$f(x) = \begin{cases} x^3, & -2 < x \leq 0 \\ -3x, & 0 \leq x < 2 \\ 5, & 2 < x \leq 4 \\ 0, & x = 2 \end{cases}$$




A) Yes

B) No

Answer: A

219) Does $\lim_{x \rightarrow 2} f(x) = f(2)$?

$$f(x) = \begin{cases} x^3, & -2 < x \leq 0 \\ -2x, & 0 \leq x < 2 \\ 6, & 2 < x \leq 4 \\ 0, & x = 2 \end{cases}$$




A) No

B) Yes

Answer: A

220) Is f continuous at $x = 0$?




A) Yes

B) No

Answer: A

221) Is f continuous at $x = 4$?




A) Yes

B) No

Answer: A

222) Is f continuous on $(-2, 4]$?



A) Yes

B) No

Answer: B

$$230) y = \frac{\sin(3\theta)}{4\theta}$$

A) continuous everywhere

B) discontinuous only when $\theta = \frac{\pi}{2}$

C) discontinuous only when $\theta = \pi$

D) discontinuous only when $\theta = 0$

Answer: D

$$231) y = \frac{2 \cos \theta}{\theta + 1}$$

A) discontinuous only when $\theta = 1$

B) discontinuous only when $\theta = -1$

C) continuous everywhere

D) discontinuous only when $\theta = \frac{\pi}{2}$

Answer: B

$$232) y = \sqrt{4x + 6}$$

A) continuous on the interval $\left[\frac{3}{2}, \infty\right)$

B) continuous on the interval $\left(-\infty, -\frac{3}{2}\right]$

C) continuous on the interval $\left[-\frac{3}{2}, \infty\right)$

D) continuous on the interval $\left[-\frac{3}{2}, \infty\right)$

Answer: D

$$233) y = \sqrt[4]{10x - 5}$$

A) continuous on the interval $\left[-\frac{1}{2}, \infty\right)$

B) continuous on the interval $\left[\frac{1}{2}, \infty\right)$

C) continuous on the interval $\left[-\infty, \frac{1}{2}\right]$

D) continuous on the interval $\left[\frac{1}{2}, \infty\right)$

Answer: D

$$234) y = \sqrt{x^2 - 2}$$

A) continuous on the interval $[-\sqrt{2}, \sqrt{2}]$

B) continuous everywhere

C) continuous on the intervals $(-\infty, -\sqrt{2}]$ and $[\sqrt{2}, \infty)$

D) continuous on the interval $[\sqrt{2}, \infty)$

Answer: C

Find the limit and determine if the function is continuous at the point being approached.

$$235) \lim_{x \rightarrow 4\pi} \sin(2x - \sin 2x)$$

A) 0; no

B) does not exist; no

C) does not exist; yes

D) 0; yes

Answer: D

$$236) \lim_{x \rightarrow \pi/2} \cos(5x - \cos 5x)$$

A) 0; no

B) does not exist; yes

C) does not exist; no

D) 0; yes

Answer: D

$$237) \lim_{x \rightarrow 2\pi} \sin\left(\frac{-3\pi}{2} \cos(\tan x)\right)$$

A) 1; no

B) does not exist; no

C) does not exist; yes

D) 1; yes

Answer: D

$$238) \lim_{x \rightarrow \pi/2} \cos\left(\frac{3\pi}{2} \cos(\tan x)\right)$$

A) does not exist; yes

B) 1; no

C) does not exist; no

D) 1; yes

Answer: C

$$239) \lim_{x \rightarrow 4} \sec(x \sec^2 x - x \tan^2 x - 1)$$

A) sec 3; no

B) does not exist; no

C) sec 3; yes

D) csc 3; yes

Answer: C

$$240) \lim_{x \rightarrow 1} \sin(x \sin^2 x + x \cos^2 x + 2)$$

A) sin 1; yes

B) does not exist; no

C) sin 3; yes

D) sin 3; no

Answer: C

$$241) \lim_{\theta \rightarrow \pi} \tan\left(\frac{-3\pi}{4} \cos(\sin \theta)\right)$$

A) 1; yes

B) 1; no

C) does not exist; no

D) 0; yes

Answer: A

$$242) \lim_{\theta \rightarrow 2\pi} \tan(\sin(2\pi \cos(\sin \theta)))$$

A) 1; yes

B) 0; no

C) does not exist; no

D) 0; yes

Answer: D

$$243) \lim_{x \rightarrow 1} \cos\left(\frac{2\pi}{3} \ln(e^x)\right)$$

A) $-\frac{1}{2}$; yes

B) 1; yes

C) $-\frac{1}{2}$; no

D) does not exist; no

Answer: A

$$244) \lim_{x \rightarrow 0} \sin^{-1}(e^{x^8})$$

A) does not exist; no

B) $\frac{\pi}{2}$; yes

C) $\frac{\pi}{4}$; yes

D) $\frac{\pi}{4}$; no

Answer: B

252)

$$f(x) = \begin{cases} x^2, & \text{if } x \leq 8 \\ x + k, & \text{if } x > 8 \end{cases}$$

A) $k = 72$

B) $k = -8$

C) $k = 56$

D) Impossible

Answer: C

253)

$$f(x) = \begin{cases} x^2, & \text{if } x \leq 8 \\ kx, & \text{if } x > 8 \end{cases}$$

A) $k = 64$

B) $k = \frac{1}{8}$

C) $k = 8$

D) Impossible

Answer: C

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Provide an appropriate response.

254) Use the Intermediate Value Theorem to prove that $3x^3 - 7x^2 - 9x + 6 = 0$ has a solution between 3 and 4.

Answer: Let $f(x) = 3x^3 - 7x^2 - 9x + 6$ and let $y_0 = 0$. $f(3) = -3$ and $f(4) = 50$. Since f is continuous on $[3, 4]$ and since $y_0 = 0$ is between $f(3)$ and $f(4)$, by the Intermediate Value Theorem, there exists a c in the interval $(3, 4)$ with the property that $f(c) = 0$. Such a c is a solution to the equation $3x^3 - 7x^2 - 9x + 6 = 0$.

255) Use the Intermediate Value Theorem to prove that $10x^4 - 7x^3 - 4x - 10 = 0$ has a solution between -1 and 0.

Answer: Let $f(x) = 10x^4 - 7x^3 - 4x - 10$ and let $y_0 = 0$. $f(-1) = 11$ and $f(0) = -10$. Since f is continuous on $[-1, 0]$ and since $y_0 = 0$ is between $f(-1)$ and $f(0)$, by the Intermediate Value Theorem, there exists a c in the interval $(-1, 0)$ with the property that $f(c) = 0$. Such a c is a solution to the equation $10x^4 - 7x^3 - 4x - 10 = 0$.

256) Use the Intermediate Value Theorem to prove that $x(x - 3)^2 = 3$ has a solution between 2 and 4.

Answer: Let $f(x) = x(x - 3)^2$ and let $y_0 = 3$. $f(2) = 2$ and $f(4) = 4$. Since f is continuous on $[2, 4]$ and since $y_0 = 3$ is between $f(2)$ and $f(4)$, by the Intermediate Value Theorem, there exists a c in the interval $(2, 4)$ with the property that $f(c) = 3$. Such a c is a solution to the equation $x(x - 3)^2 = 3$.

257) Use the Intermediate Value Theorem to prove that $4 \sin x = x$ has a solution between $\frac{\pi}{2}$ and π .

Answer: Let $f(x) = \frac{\sin x}{x}$ and let $y_0 = \frac{1}{4}$. $f\left(\frac{\pi}{2}\right) \approx 0.6366$ and $f(\pi) = 0$. Since f is continuous on $\left[\frac{\pi}{2}, \pi\right]$ and since $y_0 = \frac{1}{4}$ is between $f\left(\frac{\pi}{2}\right)$ and $f(\pi)$, by the Intermediate Value Theorem, there exists a c in the interval $\left(\frac{\pi}{2}, \pi\right)$, with the property that $f(c) = \frac{1}{4}$. Such a c is a solution to the equation $4 \sin x = x$.

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

- 258) Use a calculator to graph the function f to see whether it appears to have a continuous extension to the origin. If it does, use Trace and Zoom to find a good candidate for the extended function's value at $x = 0$. If the function does not appear to have a continuous extension, can it be extended to be continuous at the origin from the right or from the left? If so, what do you think the extended function's value(s) should be?

$$f(x) = \frac{8^x - 1}{x}$$

- A) continuous extension exists at origin; $f(0) = 0$
- B) continuous extension exists from the left; $f(0) \approx 2.0766$
- C) continuous extension exists from the right; $f(0) \approx 2.0766$
- D) continuous extension exists at origin; $f(0) \approx 2.0766$

Answer: D

- 259) Use a calculator to graph the function f to see whether it appears to have a continuous extension to the origin. If it does, use Trace and Zoom to find a good candidate for the extended function's value at $x = 0$. If the function does not appear to have a continuous extension, can it be extended to be continuous at the origin from the right or from the left? If so, what do you think the extended function's value(s) should be?


$$f(x) = \frac{4 \sin x}{|x|}$$

- A) continuous extension exists from the right; $f(0) = 1$
continuous extension exists from the left; $f(0) = -1$
- B) continuous extension exists at origin; $f(0) = 0$
- C) continuous extension exists from the right; $f(0) = 4$
continuous extension exists from the left; $f(0) = -4$
- D) continuous extension exists at origin; $f(0) = 4$

Answer: C


SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

260) A function $y = f(x)$ is continuous on $[-5, 3]$. It is known to be positive at $x = -5$ and negative at $x = 3$. What, if anything, does this indicate about the equation $f(x) = 0$? Illustrate with a sketch.



Answer: The Intermediate Value Theorem implies that there is at least one solution to $f(x) = 0$ on the interval $[-5, 3]$.

Possible graph:



261) Explain why the following five statements ask for the same information.

- Find the roots of $f(x) = 4x^3 - 1x - 5$.
- Find the x -coordinate of the points where the curve $y = 4x^3$ crosses the line $y = 1x + 5$.
- Find all the values of x for which $4x^3 - 1x = 5$.
- Find the x -coordinates of the points where the cubic curve $y = 4x^3 - 1x$ crosses the line $y = 5$.
- Solve the equation $4x^3 - 1x - 5 = 0$.

Answer: The roots of $f(x)$ are the solutions to the equation $f(x) = 0$. Statement (b) is asking for the solution to the equation $4x^3 = 1x + 5$. Statement (d) is asking for the solution to the equation $4x^3 - 1x = 5$. These three equations are equivalent to the equations in statements (c) and (e). As five equations are equivalent, their solutions are the same.

262) If $f(x) = 2x^3 - 5x + 5$, show that there is at least one value of c for which $f(x)$ equals π .

Answer: Notice that $f(0) = 5$ and $f(1) = 2$. As f is continuous on $[0, 1]$, the Intermediate Value Theorem implies that there is a number c such that $f(c) = \pi$.

263) If functions $f(x)$ and $g(x)$ are continuous for $0 \leq x \leq 4$, could $\frac{f(x)}{g(x)}$ possibly be discontinuous at a point of $[0,4]$?

Provide an example.

Answer: Yes, if $f(x) = 1$ and $g(x) = x - 2$, then $h(x) = \frac{1}{x - 2}$ is discontinuous at $x = 2$.

264) Give an example of a function $f(x)$ that is continuous at all values of x except at $x = 7$, where it has a removable discontinuity. Explain how you know that f is discontinuous at $x = 7$ and how you know the discontinuity is removable.

Answer: Let $f(x) = \frac{\sin(x - 7)}{(x - 7)}$ be defined for all $x \neq 7$. The function f is continuous for all $x \neq 7$. The function is not defined at $x = 7$ because division by zero is undefined; hence f is not continuous at $x = 7$. This discontinuity is removable because $\lim_{x \rightarrow 7} \frac{\sin(x - 7)}{x - 7} = 1$. (We can extend the function to $x = 7$ by defining its value to be 1.)


265) Give an example of a function $f(x)$ that is continuous for all values of x except $x = 6$, where it has a nonremovable discontinuity. Explain how you know that f is discontinuous at $x = 6$ and why the discontinuity is nonremovable.

Answer: Let $f(x) = \frac{1}{(x - 6)^2}$, for all $x \neq 6$. The function f is continuous for all $x \neq 6$, and $\lim_{x \rightarrow 6} \frac{1}{(x - 6)^2} = \infty$. As f is unbounded as x approaches 6, f is discontinuous at $x = 6$, and, moreover, this discontinuity is nonremovable.

MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

For the function f whose graph is given, determine the limit.

266) Find $\lim_{x \rightarrow 1^-} f(x)$ and $\lim_{x \rightarrow 1^+} f(x)$.



A) -7; -5


B) -7; -2

C) -2; -7

D) -5; -2

Answer: C


267) Find $\lim_{x \rightarrow 2^-} f(x)$ and $\lim_{x \rightarrow 2^+} f(x)$.



- A) does not exist; does not exist
 - C) -1; 3
- Answer: D

- B) 1; 1
- D) 3; -1

268) Find $\lim_{x \rightarrow 2^-} f(x)$.



- A) 4
- Answer: C


B) -1

C) 1.3

D) 2.3

269) Find $\lim_{x \rightarrow 1^-} f(x)$.

$x \rightarrow 1^-$



A) does not exist

B) -1


C) $\frac{1}{2}$

D) 2

Answer: D

270) Find $\lim_{x \rightarrow 1^+} f(x)$.

$x \rightarrow 1^+$



A) does not exist


B) 3

C) $3\frac{1}{2}$

D) 4

Answer: B

271) Find $\lim_{x \rightarrow 0} f(x)$.




A) does not exist
Answer: A

B) 0

C) -2

D) 2

272) Find $\lim_{x \rightarrow 0} f(x)$.




A) 0
Answer: A

B) does not exist

C) 1

D) -1

273) Find $\lim_{x \rightarrow 0} f(x)$.



A) 0


B) does not exist

C) -5

D) 5

Answer: B

274) Find $\lim_{x \rightarrow 1} f(x)$.



A) -1


B) $\frac{4}{5}$

C) $-\frac{4}{5}$

D) does not exist

Answer: B

275) Find $\lim_{x \rightarrow \infty} f(x)$.



A) ∞

B) does not exist

C) 0

D) -2

Answer: D

Find the limit.

276) $\lim_{x \rightarrow \infty} \frac{1}{x} - 2$

A) 2

B) -1

C) -3

D) -2

Answer: D

277) $\lim_{x \rightarrow \infty} \frac{2}{4 - (7/x^2)}$

A) $-\frac{2}{3}$

B) ∞

C) $\frac{1}{2}$

D) 2

Answer: C

278) $\lim_{x \rightarrow \infty} \frac{-5 + (2/x)}{5 - (1/x^2)}$

A) -1

B) ∞

C) 1

D) ∞

Answer: A

279) $\lim_{x \rightarrow \infty} \frac{x^2 + 8x + 6}{x^3 - 9x^2 + 5}$

A) 0

B) ∞

C) $\frac{6}{5}$

D) 1

Answer: A

280) $\lim_{x \rightarrow \infty} \frac{-9x^2 - 8x + 2}{-3x^2 + 3x + 16}$

A) 1

B) 3

C) $\frac{1}{8}$

D) ∞

Answer: B

$$281) \lim_{x \rightarrow \infty} \frac{6x + 1}{9x - 7}$$

$$A) \frac{2}{3}$$

$$B) \infty$$

$$C) -\frac{1}{7}$$

$$D) 0$$

Answer: A

$$282) \lim_{x \rightarrow \infty} \frac{2x^3 - 4x^2 + 3x}{-x^3 - 2x + 5}$$

$$A) -2$$

$$B) \frac{3}{2}$$

$$C) 2$$

$$D) \infty$$

Answer: A

$$283) \lim_{x \rightarrow \infty} \frac{5x^3 + 3x^2}{x - 6x^2}$$

$$A) \infty$$

$$B) 5$$

$$C) -\infty$$

$$D) -\frac{1}{2}$$

Answer: A

$$284) \lim_{x \rightarrow \infty} \frac{\cos 2x}{x}$$

$$A) 2$$

$$B) 0$$

$$C) 1$$

$$D) \infty$$

Answer: B

Divide numerator and denominator by the highest power of x in the denominator to find the limit.

$$285) \lim_{x \rightarrow \infty} \sqrt{\frac{49x^2}{2 + 4x^2}}$$

$$A) \frac{7}{2}$$

$$B) \frac{49}{4}$$

C) does not exist

$$D) \frac{49}{2}$$

Answer: A

$$286) \lim_{x \rightarrow \infty} \sqrt{\frac{64x^2 + x - 3}{(x - 17)(x + 1)}}$$

$$A) 8$$

$$B) 0$$

$$C) 64$$

$$D) \infty$$

Answer: A

$$287) \lim_{x \rightarrow \infty} \frac{-3\sqrt{x} + x^{-1}}{3x + 3}$$

$$A) -1$$

$$B) \frac{1}{3}$$

$$C) \infty$$

$$D) 0$$

Answer: D

$$288) \lim_{x \rightarrow \infty} \frac{-5x^{-1} + 4x^{-3}}{-4x^{-2} + x^{-5}}$$

- A) ∞ B) $\frac{5}{4}$ C) 0 D) ∞

Answer: A

$$289) \lim_{x \rightarrow \infty} \frac{\sqrt[3]{x} - 5x + 7}{-7x + x^{2/3} + 2}$$

- A) ∞ B) $\frac{5}{7}$ C) 0 D) $\frac{7}{5}$

Answer: B

$$290) \lim_{t \rightarrow \infty} \frac{\sqrt{9t^2 - 27}}{t - 3}$$

- A) 9 B) does not exist C) 3 D) 27

Answer: C

$$291) \lim_{t \rightarrow \infty} \frac{\sqrt{25t^2 - 125}}{t - 5}$$

- A) does not exist B) 125 C) 25 D) 5

Answer: D

$$292) \lim_{x \rightarrow \infty} \frac{7x + 5}{\sqrt{5x^2 + 1}}$$

- A) 0 B) $\frac{7}{5}$ C) ∞ D) $\frac{7}{\sqrt{5}}$

Answer: D

Find the limit.

$$293) \lim_{x \rightarrow 2} \frac{1}{x + 2}$$

- A) 1/2 B) Does not exist C) $-\infty$ D) ∞

Answer: B

$$294) \lim_{x \rightarrow 7^+} \frac{1}{x - 7}$$

- A) -1 B) 0 C) ∞ D) ∞

Answer: D

$$295) \lim_{x \rightarrow 2^+} \frac{1}{x + 2}$$

- A) ∞ B) ∞ C) -1 D) 0

Answer: A

$$296) \lim_{x \rightarrow 10^+} \frac{1}{(x - 10)^2}$$

A) $-\infty$

B) 0

C) -1

D) ∞

Answer: D

$$297) \lim_{x \rightarrow 3^-} \frac{4}{x^2 - 9}$$

A) $-\infty$

B) ∞

C) 0

D) -1

Answer: B

$$298) \lim_{x \rightarrow 2^+} \frac{5}{x^2 - 4}$$

A) $-\infty$

B) 1

C) 0

D) ∞

Answer: D

$$299) \lim_{x \rightarrow 1^-} \frac{4}{x^2 - 1}$$

A) 0

B) ∞

C) 1

D) $-\infty$

Answer: D

$$300) \lim_{x \rightarrow 0} \frac{1}{x^{2/3}}$$

A) $-\infty$

B) 2/3

C) 0

D) ∞

Answer: D

$$301) \lim_{x \rightarrow (\pi/2)^+} \tan x$$

A) 0

B) $-\infty$

C) 1

D) ∞

Answer: B

$$302) \lim_{x \rightarrow (-\pi/2)^-} \sec x$$

A) $-\infty$

B) 1

C) 0

D) ∞

Answer: D

$$303) \lim_{x \rightarrow 0^+} (1 + \csc x)$$

A) 1

B) 0

C) ∞

D) Does not exist

Answer: C

$$304) \lim_{x \rightarrow 0} (1 - \cot x)$$

A) ∞

B) 0

C) $-\infty$

D) Does not exist

Answer: D

$$305) \lim_{x \rightarrow 0^+} \frac{x^2}{2} - \frac{1}{x}$$

A) Does not exist

B) ∞

C) ∞

D) 0

Answer: C

$$306) \lim_{x \rightarrow \sqrt[3]{2}} \frac{x^2}{2} - \frac{1}{x}$$

A) ∞

B) 0

C) ∞

D) $2\sqrt[3]{2}$

Answer: B

$$307) \lim_{x \rightarrow -1^-} \frac{x^2 - 5x + 4}{x^3 - x}$$

A) ∞

B) 0

C) ∞

D) $-\frac{3}{2}$

Answer: D

$$308) \lim_{x \rightarrow 0} \frac{x^2 - 3x + 2}{x^3 - x}$$

A) 2

B) ∞

C) ∞

D) Does not exist

Answer: D

$$309) \lim_{x \rightarrow 0^+} \left(\frac{1}{x^{1/5}} + 2 \right)$$

A) 2

B) ∞

C) Does not exist

D) ∞

Answer: D

$$310) \lim_{x \rightarrow 0^-} \left(\frac{1}{x^{2/5}} + 1 \right)$$

A) Does not exist

B) ∞

C) ∞

D) 1

Answer: B

311)

$$\lim_{x \rightarrow 3^+} \left(\frac{1}{x^{4/5}} - \frac{1}{(x-3)^{1/5}} \right)$$

A) ∞

B) ∞

C) Does not exist

D) 0

Answer: A

$$312) \lim_{x \rightarrow 5^-} \left(\frac{1}{x^{4/5}} - \frac{1}{(x-5)^{4/5}} \right)$$

A) ∞

B) 0


C) ∞

D) Does not exist


Answer: C

Graph the rational function. Include the graphs and equations of the asymptotes.


313) $f(x) = \frac{x}{x-1}$




A) asymptotes: $x = -1, y = 1$




B) asymptotes: $x = 1, y = 0$



C) asymptotes: $x = 1, y = 1$




D) asymptotes: $x = -1, y = 0$




Answer: C


314) $f(x) = \frac{x}{x^2 + x + 1}$




A) asymptotes: $x = 1, x = -1$




B) asymptote: $y = 1$



C) asymptote: $y = 0$




D) asymptote: $y = 0$




Answer: C


315) $f(x) = \frac{x^2 + 3}{x^3}$




A) asymptotes: $x = 0, y = 0$




B) asymptote: $y = 0$



C) asymptotes: $x = 0, y = 0$




D) asymptotes: $x = 0, y = 0$




Answer: D


316) $f(x) = \frac{1}{x+1}$




A) asymptotes: $x = -1, y = 0$




B) asymptotes: $x = -1, y = 0$



C) asymptotes: $x = 1, y = 0$




D) asymptotes: $x = 1, y = 0$




Answer: B


317) $f(x) = \frac{x - 1}{x + 1}$




A) asymptotes: $x = -1, y = 1$




B) asymptotes: $x = -1, y = -1$



C) asymptotes: $x = 1, y = 1$




D) asymptotes: $x = 1, y = -1$




Answer: A


318) $f(x) = \frac{1}{(x+2)^2}$




A) asymptotes: $x = 2, y = 0$




B) asymptotes: $x = -2, y = 0$



C) asymptotes: $x = 2, y = 0$




D) asymptotes: $x = -2, y = 0$




Answer: B


319) $f(x) = \frac{2x^2}{4 - x^2}$




A) asymptotes: $x = -2$, $x = 2$, $y = 0$




B) asymptotes: $x = -2$, $x = 2$, $y = 2$



C) asymptotes: $x = -2$, $x = 2$, $y = -2$




D) asymptotes: $x = -2$, $x = 2$, $y = 0$




Answer: C


320) $f(x) = \frac{2 - x^2}{2x + 4}$




A) asymptotes: $x = -4, y = -\frac{1}{8}x + \frac{1}{2}$




B) asymptotes: $x = -2, y = -\frac{1}{4}x + \frac{1}{2}$



C) asymptotes: $x = -2, y = -x + 2$




D) asymptotes: $x = -2, y = -\frac{1}{2}x + 1$




Answer: D


321) $f(x) = \frac{2 - 2x - x^2}{x}$




A) asymptotes: $x = 0, y = -x - 4$




B) asymptotes: $x = 0, y = -x + 2$



C) asymptotes: $x = 0, y = -x - 2$



D) asymptotes: $x = 0, y = -x$




Answer: C


SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

Sketch the graph of a function $y = f(x)$ that satisfies the given conditions.


$$322) f(0) = 0, f(1) = 3, f(-1) = -3, \lim_{x \rightarrow \infty} f(x) = -2, \lim_{x \rightarrow -\infty} f(x) = 2.$$




Answer: Answers may vary. One possible answer:




323) $f(0) = 0$, $f(1) = 6$, $f(-1) = 6$, $\lim_{x \rightarrow \pm\infty} f(x) = -6$.




Answer: Answers may vary. One possible answer:




324) $f(0) = 6, f(1) = -6, f(-1) = -6, \lim_{x \rightarrow \pm\infty} f(x) = 0.$




Answer: Answers may vary. One possible answer:



325) $f(0) = 0$, $\lim_{x \rightarrow \pm\infty} f(x) = 0$, $\lim_{x \rightarrow 4^-} f(x) = -\infty$, $\lim_{x \rightarrow 4^+} f(x) = -\infty$, $\lim_{x \rightarrow 4^-} f(x) = \infty$, $\lim_{x \rightarrow 4^+} f(x) = \infty$.




Answer: Answers may vary. One possible answer:




Find a function that satisfies the given conditions and sketch its graph.


$$326) \lim_{x \rightarrow \pm\infty} f(x) = 0, \quad \lim_{x \rightarrow 1^-} f(x) = \infty, \quad \lim_{x \rightarrow 1^+} f(x) = \infty.$$




Answer: (Answers may vary.) Possible answer: $f(x) = \frac{1}{|x - 1|}$.




327) $\lim_{x \rightarrow \pm \infty} f(x) = 0$, $\lim_{x \rightarrow 3^-} f(x) = \infty$, $\lim_{x \rightarrow 3^+} f(x) = \infty$.




Answer: (Answers may vary.) Possible answer: $f(x) = \frac{1}{|x + 3|}$.




328) $\lim_{x \rightarrow \infty} g(x) = -5$, $\lim_{x \rightarrow \infty} g(x) = 5$, $\lim_{x \rightarrow \theta^+} g(x) = 5$, $\lim_{x \rightarrow \theta^-} g(x) = -5$.




Answer: (Answers may vary.) Possible answer: $f(x) = \begin{cases} 5, & x > 0 \\ -5, & x < 0 \end{cases}$




329) $\lim_{x \rightarrow \infty} f(x) = 0$, $\lim_{x \rightarrow 0^+} f(x) = \infty$.




Answer: (Answers may vary.) Possible answer: $f(x) = \frac{1}{\sqrt{x}}$.



330) $\lim_{x \rightarrow \infty} f(x) = 1$, $\lim_{x \rightarrow \theta^+} f(x) = -1$, $\lim_{x \rightarrow \theta^-} f(x) = -1$, $\lim_{x \rightarrow \theta^-} f(x) = 1$



Answer: (Answers may vary.) Possible answer: $f(x) = \begin{cases} 1, & x < 0 \\ -1, & x > 0 \end{cases}$



MULTIPLE CHOICE. Choose the one alternative that best completes the statement or answers the question.

Find the limit.

331) $\lim_{x \rightarrow \infty} (4x - \sqrt{16x^2 - 5x + 6})$

A) $\frac{5}{8}$

B) ∞

C) 0

D) -12

Answer: A

332) $\lim_{x \rightarrow \infty} (\sqrt{x^2 + 12x} - x)$

A) 6

B) 0

C) 12

D) ∞

Answer: A

333) $\lim_{x \rightarrow \infty} (\sqrt{3x^2 + 7} - \sqrt{3x^2 - 3})$

A) ∞

B) 0

C) $\frac{1}{2\sqrt{3}}$

D) $\sqrt{3}$

Answer: B

334) $\lim_{x \rightarrow \infty} \sqrt{x^2 + 2x} - \sqrt{x^2 - 7x}$

A) does not exist

B) $-\frac{5}{2}$

C) $\frac{9}{2}$

D) 9

Answer: C

Provide an appropriate response.

335) Which of the following statements defines $\lim_{x \rightarrow x_0} f(x) = \infty$?

I. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 - \delta < x < x_0 + \delta$.

II. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 < x < x_0 + \delta$.

III. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 - \delta < x < x_0$.

A) III

B) II

C) I

D) None

Answer: C

336) Which of the following statements defines $\lim_{x \rightarrow (x_0)^-} f(x) = \infty$?

I. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 - \delta < x < x_0 + \delta$.

II. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 < x < x_0 + \delta$.

III. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 - \delta < x < x_0$.

A) II

B) III

C) I

D) None

Answer: B

337) Which of the following statements defines $\lim_{x \rightarrow (x_0)^+} f(x) = \infty$?

I. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 - \delta < x < x_0 + \delta$.

II. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 < x < x_0 + \delta$.

III. For every positive real number B there exists a corresponding $\delta > 0$ such that $f(x) > B$ whenever $x_0 - \delta < x < x_0$.

A) III

B) II

C) I

D) None

Answer: B

338) Which of the following statements defines $\lim_{x \rightarrow x_0} f(x) = -\infty$?

- I. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 - \delta < x < x_0 + \delta$.
- II. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 < x < x_0 + \delta$.
- III. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 - \delta < x < x_0$.

A) II B) III C) I D) None

Answer: C

339) Which of the following statements defines $\lim_{x \rightarrow (x_0)^+} f(x) = -\infty$?

- I. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 - \delta < x < x_0 + \delta$.
- II. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 < x < x_0 + \delta$.
- III. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 - \delta < x < x_0$.

A) I B) II C) III D) None

Answer: B

340) Which of the following statements defines $\lim_{x \rightarrow (x_0)^-} f(x) = -\infty$?

- I. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 - \delta < x < x_0 + \delta$.
- II. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 < x < x_0 + \delta$.
- III. For every negative real number B there exists a corresponding $\delta > 0$ such that $f(x) < B$ whenever $x_0 - \delta < x < x_0$.

A) III B) II C) I D) None

Answer: A

341) Which of the following statements defines $\lim_{x \rightarrow \infty} f(x) = \infty$?

- I. For every positive real number B there exists a corresponding positive real number N such that $f(x) > B$ whenever $x > N$.
- II. For every positive real number B there exists a corresponding negative real number N such that $f(x) > B$ whenever $x < N$.
- III. For every negative real number B there exists a corresponding negative real number N such that $f(x) < B$ whenever $x < N$.
- IV. For every negative real number B there exists a corresponding positive real number N such that $f(x) < B$ whenever $x > N$.

A) II B) IV C) I D) III

Answer: A

SHORT ANSWER. Write the word or phrase that best completes each statement or answers the question.

342) Use the formal definitions of limits to prove $\lim_{x \rightarrow 0} \frac{4}{|x|} = \infty$

Answer: Given $B > 0$, we want to find $\delta > 0$ such that $0 < |x - 0| < \delta$ implies $\frac{4}{|x|} > B$.

Now, $\frac{4}{|x|} > B$ if and only if $|x| < \frac{4}{B}$.

Thus, choosing $\delta = 4/B$ (or any smaller positive number), we see that

$|x| < \delta$ implies $\frac{4}{|x|} > \frac{4}{|\delta|} \geq B$.

Therefore, by definition $\lim_{x \rightarrow 0} \frac{4}{|x|} = \infty$

343) Use the formal definitions of limits to prove $\lim_{x \rightarrow 0^+} \frac{5}{x} = \infty$

Answer: Given $B > 0$, we want to find $\delta > 0$ such that $x_0 < x < x_0 + \delta$ implies $\frac{5}{x} > B$.

Now, $\frac{5}{x} > B$ if and only if $x < \frac{5}{B}$.

We know $x_0 = 0$. Thus, choosing $\delta = 5/B$ (or any smaller positive number), we see that

$x < \delta$ implies $\frac{5}{x} > \frac{5}{\delta} \geq B$.

Therefore, by definition $\lim_{x \rightarrow 0^+} \frac{5}{x} = \infty$